BAM Site - Field Survey Form Site Sheet no: **Survey Name** Plot Identifier Recorders 17 b 22 Date HHIP 56 V102 Zone Datum **IBRA** region Photo# Zone ID Easting Northina Orientation of midline **Dimensions** from the 0 m point. Confidence: **Vegetation Class** H M L Confidence: HU833 - WSP- RP2.1 **Plant Community Type** EEC: H M L Record easting and northing from the plot marker. If applicable, orient picket so that perforated rib points along direction of midline. Dimensions (Shape) of 0.04 ha base plot inside 0.1 ha FA plot should be identified, magnetic bearing taken along midline. BAM Attribute (20 x 50 m plot) # Tree Stems Count **BAM Attribute** Sum values Record number of (400 m² plot) dbh Non Fuc Hollows[†] Fuc* living eucalypt* (Euc*) and living Trees large trees for native non-eucalypt 2 2 Euc* & Non Euc cm (Non Euc) stems Shrubs separately 50 - 79 cm Z Count of Grasses etc. * includes all species Native of Eucalyptus, 30 - 49 cmRichness **Forbs** Corymbia, Angophora, Lophostemon and **Ferns** 20 - 29 cm Syncarpia Other †Record total 10 - 19 cm 21 number of stems by Trees size class with hollows (including 5 - 9 cm n/a Sum of Shrubs dead stems/trees) Cover of native Grasses etc. < 5 cm n/a vascular plants by **Forbs** total Length of logs (m) arowth (≥0 cm diameter, >50 cm form group Ferns 30 in length) Counts must apply to each size class when the number of living tree stems within the size class is ≤10. Other Estimates can be used when the number of living tree stems within a class is > 10. Estimates should draw from the number series: 10, 20, 30..., 100, 200, 300 High Threat Weed cover For a multi-stemmed tree, only the largest living stem is included in the count/estimate. For hollows count only the presence of a stem containing hollows, not the count of hollows in that stem. Only count as 1 stem per tree where tree is multi-stemmed. The hollow-bearing stem may be a dead stem. BAM Attribute (1 x 1 m plots) Litter cover (%) Bare ground cover (%) Cryptogam cover (%) 7060 70 Subplot score (% in each) 60 70 Average of the 5 subplots 66 Litter cover is assessed as the average percentage ground cover of litter recorded from five 1 m x 1 m plots located on alternate sides and 5 m from the plot midline at the locations 5, 15, 25, 35, and 45 m along the midline. Litter cover includes leaves, seeds, twigs, branchlets and branches (less than 10 cm in diameter). Within these 1 m x 1 m plots assessors may also record the cover of rock, bare ground and cryptogam soil crusts. Collection of these data is optional - the data do not currently contribute to assessment scores, they hold potential value for future vegetation integrity assessment attributes and benchmarks, and for enhancing PCT description Physiography + site features that may help in determining PCT and Management Zone (optional) Morphological Landform Landform Microrelief Type Element Pattern Soil Surface Soil Soil Lithology Texture Colour Depth Distance to nearest water and type Slope Aspect Site Drainage Severity Age Observational evidence: Plot Disturbance Clearing (inc. logging) Cultivation (inc. pasture) Soil erosion Firewood / CWD removal Grazing (identify native/stock) Fire damage Storm damage

Severity: 0=no evidence, 1=light, 2=moderate, 3=severe

Age: R=recent (<3yrs), NR=not recent (3-10yrs), O=old (>10yrs)

Weediness Other

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date		V1 02	SC

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratu m	vouch er
	Angophara Costata		20	10		
	Conymbia gummfera		ιG	6		
	Eucalyph's capitellate		5	3		
	Barksia Spirulosa		25	50		
	Pittosporn undlatin		10	20		
	Macrozania Communis		2	16		
	Kartlerdeg latifolia		2	16		
	Hibbertia aspera		16	50		
	Mirbelia rubfolia		\$	50		
	Lesto spermy m transcring		10	20		
	Resource linearis		2	16		
	Acacia myrtifolia		1	(6		
	Py Hanaen spp.			20		
	Longada obligua		S	100		
	Imperata Cylindrica		5	100		
	Phyllonthis Shitells		0.2	20		
	Themeda triandra		10	500		
	Pterdum esculertum		5	50		
	Acacia terminali		0.1	5		
	Epacris mic-ophylla		0.2	20		
	Gooder a heterofila		0.2	50		
	Philothrit destra		01	20		
	Pultanaea netusa		0.5	50		
	Allocasygrina littoralis		8.5	2_		
	Dignella 1.0		2	160		
	Billerdieria scanderi		0-2	16		
	Dodonala triguetra		2	20		
	Rytidosperna pallidin		5	100		
	Comatia silatfolia		1	5		
	Pterosofilis sop. whens		0.1	10		
	Entolasia Ifricta		5	100		
	Acaeia Melfolia		1	10		
	Daviesia Vicifolia		1	10		
	Cassytha puberces		0.5	5		
	Polystias sambuccafola		a.5	5		
	Podolobium , licitate		0.5	5		
	Personia levis		6.1	2		-
	38					
	39					
	40					
L	de: see Growth Form definitions in Annendix 1 N: native F: exotic HTE: high threa		GF – c			

GF Code: see Growth Form definitions in Appendix 1 **N:** native, **E:** exotic, **HTE:** high threat exotic **GF - circle code** if 'top 3'. **Cover:** 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); **Note:** 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m **Abundance:** 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

BAM Site - Field Survey Form Site Sheet no: 1 of **Plot Identifier** Survey Name Recorders 6 22 Date V1 64 Zone Datum **IBRA** region Photo # Zone ID Easting Northing Orientation of midline **Dimensions** from the 0 m point. Confidence: **Vegetation Class** M Confidence: HU 841 - WSP-RP2J **Plant Community Type** EEC: M I Record easting and northing from the plot marker. If applicable, orient picket so that perforated rib points along direction of midline. Dimensions (Shape) of 0.04 ha base plot inside 0.1 ha FA plot should be identified, magnetic bearing taken along midline. BAM Attribute (20 x 50 m plot) # Tree Stems Count **BAM Attribute** Record number of Sum values (400 m² plot) Non Euc Hollows[†] living eucalypt* (Euc*) and living Trees 80 + large trees for Euc* & Non Euc native non-eucalypt cm (Non Euc) stems Shrubs separately 50 - 79 cm 10 7_ Count of Grasses etc. * includes all species Native of Eucalyptus, Richness Forbs 30 - 49 cm Corymbia. 20 Angophora. **Ferns** Lophostemon and 20 - 29 cm Svncarpia Other † Record total 10 - 19 cm number of stems by Trees size class with hollows (including 5 - 9 cm n/a Sum of Shrubs dead stems/trees) Cover of native Grasses etc. < 5 cm n/a vascular plants by **Forbs** total Length of logs (m) growth (≥0 cm diameter, >50 cm form group Ferns in length) Counts must apply to each size class when the number of living tree stems within the size class is ≤10. Other Estimates can be used when the number of living tree stems within a class is > 10. Estimates should draw from the number series: 10, 20, 30..., 100, 200, 300 High Threat Weed cover For a multi-stemmed tree, only the largest living stem is included in the count/estimate. For hollows count only the presence of a stem containing hollows, not the count of hollows in that stem. Only count as 1 stem per tree where tree is multi-stemmed. The hollow-bearing stem may be a dead stem. BAM Attribute (1 x 1 m plots) Litter cover (%) Bare ground cover (%) Cryptogam cover (%) Rock cover (%) Subplot score (% in each) 60 70 30 70 Average of the 5 subplots 59 Litter cover is assessed as the average percentage ground cover of litter recorded from five 1 m x 1 m plots located on alternate sides and 5 m from the plot midline at the locations 5, 15, 25, 35, and 45 m along the midline. Litter cover includes leaves, seeds, twigs, branchlets and branches (less than 10 cm in diameter). Within these 1 m x 1 m plots assessors may also record the cover of rock, bare ground and cryptogam soil crusts. Collection of these data is optional - the data do not currently contribute to assessment scores, they hold potential value for future vegetation integrity assessment attributes and benchmarks, and for enhancing PCT description Physiography + site features that may help in determining PCT and Management Zone (optional) Morphological Landform Landform Microrelief Pattern Type Element Soil Surface Lithology Texture Colour Depth Distance to nearest Site Drainage Slone Aspect water and type Severity Age Observational evidence: Plot Disturbance code Clearing (inc. logging) Cultivation (inc. pasture) Soil erosion Firewood / CWD removal Grazing (identify native/stock) Fire damage Storm damage Weediness Other

Severity: 0=no evidence, 1=light, 2=moderate, 3=severe

Age: R=recent (<3yrs), NR=not recent (3-10yrs), O=old (>10yrs)

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date		VI 04	SC

GF Code		I, E or HTE	Cover	Abund	stratu m	vouch er
	Angeshera costata		5	4		
	Angophora costata Eccolopts piperta		15	7		
	(organtia gummifera		5	3		
	E. globoidea		5	3		
	Syntaraia alomelifera		1	i.		
	30donoec triggetia		10	50	mundadada a proposada da de	
	lterdum excles him		10	50	A STATE OF THE STA	
	alochidian fordinandii		,	5		
	Banksia spinulosa		5	20		
	Leptespernon polygal folion		ζ	200		
	In para to extende ca var marco		16	(00		
	Kanthorrea lat		1	5		
	Entalasia Sticta		10	500		
	Vanandea longifelia		i	20		
	Lomand-a multiflara		1	50		
•	Maerozamia Communis		,	5		
	Lindsea microphy 1/9		0.2	100		
•			0.2	100		
	Personia lous		,	10		
			0.1	50		
			,	200		
	Dianella c.p Hildourtia aspera		0.2	50		
<u> </u>						
	Eistrephis latifolia		0.5	20		
	Goodenia Neteropylla			100		
	Pesmodium rytidophy 11 a		0.5	16		
	Pandenes paindora &		0.1	20		
	Frana piperesces		0.2			
	Brunoniella avstralis		0.5			
			0.2			
	Polyscias sambiceatiless		0.2			
	Heacia longifolia	· · · · · · · · · · · · · · · · · · ·	l l	5	-	-
	Breynia delongifolia		1	5		
	Lorhatia siliafolia		0-2	5		
	Microlaena etipoides		2	100		
	Parsons 19 straining	······	0.2	(8)		
	Personia levis		0.5	18		
	Pterostylis spp. (Ntans)		0.1	10		
	Advantum aethiopicum		0.2	30		
	Smilax australis		0.2	20		
	Calachlaena dubin			50		
	de: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat e		GF – c			

GF Code: see Growth Form definitions in Appendix 1

N: native, E: exotic, HTE: high threat exotic

GF - circle code if 'top 3'.

BAM Site - Field Survey Form Site Sheet no: Plot Identifier **Survey Name** Recorders Date 6 22 V103 SC 1HHIP Zone Datum IBRA region Photo # Zone ID Easting Northing Orientation of midline **Dimensions** from the 0 m point. Confidence: **Vegetation Class** М WSP-RP2 Confidence: **Plant Community Type** EEC: M L Record easting and northing from the plot marker. If applicable, orient picket so that perforated rib points along direction of midline. Dimensions (Shape) of 0.04 ha base plot inside 0.1 ha FA plot should be identified, magnetic bearing taken along midline.

	Attribute m² plot)	Sum values
	Trees	
	Shrubs	
Count of Native	Grasses etc.	
Richness	Forbs	
	Ferns	
	Other	
	Trees	
Sum of	Shrubs	
Cover of native		
of native	Grasses etc.	
of native vascular plants by	Grasses etc.	
of native vascular		
of native vascular plants by growth	Forbs	

BAM Attribute	(20 x 50	m plot)	# Tree	Stems Count	Record number of
dbh		Euc*	Non Euc	Hollows [†]	living eucalypt*
large trees for Euc* & Non Euc	80 + cm	(Merc A	: : : : : : : : : : : : : : : : : : : :	(Euc*) and living native non-eucalypt (Non Euc) stems
50 –	79 cm	6		(* includes all species
30 – 49 cm	C	7			of Eucalyptus, Corymbia, Angophora,
20 – 29 cm	1.	2			Lophostemon and Syncarpia
10 – 19 cm	2	2			† Record total number of stems by size class with
5 – 9 cm	10	f		n/a	hollows (including dead stems/trees)
< 5 cm	10	ĺ		n/a	
Length of logs (≥ 0 cm diameter, in length)				16	total 80

Counts must apply to each size class when the number of living tree stems within the size class is \leq 10. Estimates can be used when the number of living tree stems within a class is > 10. Estimates should draw from the number series: 10, 20, 30..., 100, 200, 300

For a multi-stemmed tree, only the largest living stem is included in the count/estimate. For hollows count only the presence of a stem containing hollows, not the count of hollows in that stem. Only count as 1 stem per tree where tree is multi-stemmed. The hollow-bearing stem may be a dead stem.

BAM Attribute (1 x 1 m plots)	Litter cover (%)	Bare ground cover (%)	Cryptogam cover (%)	Rock cover (%)
Subplot score (% in each)	35 60 80 36 20			
Average of the 5 subplots	45			

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1 m x 1 m plots located on alternate sides and 5 m from the plot midline at the locations 5, 15, 25, 35, and 45 m along the midline. Litter cover includes leaves, seeds, twigs, branchlets and branches (less than 10 cm in diameter). Within these 1 m x 1 m plots assessors may also record the cover of rock, bare ground and cryptogam soil crusts. Collection of these data is optional - the data do not currently contribute to assessment scores, they hold potential value for future vegetation integrity assessment attributes and benchmarks, and for enhancing PCT description

Physiography + site features that may help in determining PCT and Management Zone (optional) Morphological Landform Landform Microrelief Element Pattern Soil Surface Soil Lithology Texture Colour Depth Distance to nearest Slope Aspect Site Drainage water and type

Plot Disturbance	Severity code	Age code	Observational evidence:
Clearing (inc. logging)			
Cultivation (inc. pasture)	No.		
Soil erosion	SECONO DE LA CONTRACTOR		
Firewood / CWD removal			
Grazing (identify native/stock)			
Fire damage			
Storm damage			
Weediness			
Other			

Severity: 0=no evidence, 1=light, 2=moderate, 3=severe

Age: R=recent (<3yrs), NR=not recent (3-10yrs), O=old (>10yrs)

400 m ² plot: Sheet _ of	_ S	Survey Name	Plot Identifier	Recorders
Date				

GF ode	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratu m	voucl er
	Angophera Costata		15	8		
	Caryonbia gumon. fore		10	4		
	Esc capitellate		10	4		
	Banksia spiriosa		26	50		
	E. operta		/	1		
	Dodonala trigretra		10	25		
			2	1		
	Exocaras cuprescitoross Allocasiarina torolosa		e-	6		
	Pterdium exception		10	50		
	Kemeda trandra		10	200		
	Entologia Stricta		10	200		
***************************************	Xa-thornea lahfola		(15		
			2	50		
	RHangey evchila Mirbelia ribifolia			50		
	Comanda obligia		10	100		
	Podolobium licitalium		/	5		
	Rytidosperma pollidom		10		}	
	Imperata cylindrica			100		
	Pratia purpurescens		0.2	700		
	Retaneare retisa		2	20		
	Macrozania Commis		6-1	2		
	Davieria elicifolia		0,2	5		
	Acaera licifola		-	20		
	Comepena evicinion		(200		
	Goodenca hederacea		0.2	20		
	Leptospernum trinervium			20		
	Personia leuts		(15		ļ
	Diarella Caerlea podite		6.5	50		
	Hibbetia asper		0.5	20		
	Dianella Caerlea & productes Hibberta aspera Billerdiera scanders		0.2	(0		-
	Daviesia Squarasa		1	10		
	Comertia sialifola		/	10		
	Pseuderantenum variobile		0.2	20		
	Acacia ferminalis		0.5	5		
	Acacia ferminalis Parsonsia strammen	<u> </u>	0.2	3		
	26					
	37					
	38					
	25 · · · · · · · · · · · · · · · · · · ·					
	40					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'. **Cover:** 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); **Note:** 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...