

Watt Street Commercial Centre Level 1, 45 Watt Street Newcastle NSW 2300 Telephone: +61 2 4903 5500 Fax: +61 2 4929 5363

19 June 2020

Reference: 0470861

www.erm.com

Javier Canon
Senior Policy Officer
Department of Planning, Industry & Environment
4 Parramatta Square, 12 Darcy Street
PARRAMATTA NSW 2150

ERM

Dear Javier Canon

Subject: Bonshaw Solar Farm (SSD 9438) - Response to RFI Dated 1 May 2020

1. INTRODUCTION

GAIA Australia Pty Ltd (GAIA) are seeking approval to develop a large scale photovoltaic (PV) generation facility with a capacity of 200 megawatts (MW) and associated infrastructure, including a Lithium-ion Energy Storage System (ESS/Li-ion). The proposed development of the Bonshaw Solar Farm (the 'Project') is located at Bonshaw, within the Inverell Local Government Area in New South Wales (NSW) (refer to **Figure 1-1**). The Project is located approximately 66 km north of Inverell, with site access fronting Bruxner Highway.

1.1 Project Progress

The Project's Environmental Impact Statement (EIS) was exhibited over a 29 day period, from 6 November 2019 to 4 December 2019. There were 13 submissions received from government agencies and two public submissions during this period. A Response To Submissions (RTS) was submitted to the Department of Planning, Industry and Environment (DPIE) to address these comments.

Since exhibition and subsequent submission of the RTS, the Project was amended through the submission of an Amendment Report (AR) to provide details of the overhead connection.

In addition, on 8 April 2020 the DPIE had requested further information on five (5) matters relating to the Project updates. These matters were addressed through the submission of a letter dated 16 April 2020.

1.2 Purpose of Letter


On 1 May 2020, DPIE provided a 'Request for Additional Information', seeking further details to effectively address eight (8) items raised, relating to traffic, biodiversity assessment, flooding and water supply. Upon submitting an initial response on 22 May 2020, DPIE requested additional information on three (3) other items on 27 May 2020. These matters are collectively outlined in **Table 2-1** below, with consideration for the three additional matters referred to in the 'Item' column as a sub-reference identified by the inclusion of the letter 'a' attached to the relevant matter.

The purpose of this letter is to provide details on items raised, to assist with the Department's assessment and determination of the Project. Details to address the items raised by the Department is provided in **Section 2** below.

In addition, this letter also addresses comments raised by the Biodiversity and Conservation Division (BCD) in relation to the location of AHIMS #11-3-0083 (refer to **Section 2.1**).

Page 1 of 19

Offices worldwide



2. PROVISION OF ADDITIONAL INFORMATION

Each of the eleven collective items raised by DPIE have been outlined in **Table 2-1** below, with supporting details provided in the right column as a response to the request for information.

Table 2-1 Request for Information Details

Table 2-1 Request for Information Details				
Item	DPIE Comments	Details		
Traffic	raffic and Access Route			
1	provide an assessment of the traffic impacts of the all phases of the development on all proposed access routes and key intersections identifying any improvements required to maintain road safety and associated assessment of the impact of any additional road improvements (including biodiversity and heritage);	The Traffic Impact Assessment (TIA) prepared by Seca Solution has identified the impacts during construction, when there will be the biggest impact, and has determined that no road upgrades on the external road network is required other than proposed upgrades to the site access point. All the roads used for heavy vehicle access to the Site currently carry B-doubles and regional / state and local traffic demands. The overall traffic flows along these roads are well within acceptable capacity limits and through the major urban centres the road capacity is increased and monitored by the road authority to ensure adequate capacity is provided. During the operational phase of the project, the traffic volumes are significantly lower than during construction with a maximum of 10 staff working on site and allowance for all staff to drive will generate 10 inbound vehicles in the morning and 10 outbound in the afternoon. For the decommissioning stage of the project, which could occur in 20-30 years time, the expected traffic demands would be less than those associated with the construction. The possible decommissioning stage of the development has been discussed with TfNSW who agreed that this would be more appropriate to review if the Project Site is decommissioned, due to a wide range of factors that could impact the Project at that stage. An updated Traffic Impact Assessment has been provided in Attachment A below.		
1a	Provide an assessment of the proposed site access, considering heavy traffic safe turning requirements, sight distances, swept paths and safe deceleration requirements, and confirming any road upgrades required as a result of the assessment. The assessment of impacts of any road improvements must include a desktop biodiversity and heritage addendum.	Access to the site will be via the existing access provided to the electric substation adjacent to the subject site. This access was constructed to allow for the construction activities associated with the electric substation and is now used for maintenance purposes only for the substation. This access will be upgraded to provide a left turn deceleration lane and shoulder widening to provide an auxiliary right turn treatment, both in accordance with Austroads Guidelines (refer to Figure 2-1).		

Item	DPIE Comments	Details
		An assessment of the road upgrade impacts on biodiversity and heritage matters is included in Attachment B.
	provide swept paths at key intersections for the largest vehicle that will traverse the proposed access routes, including the Bruxner Highway access;	Any swept path analysis can only be completed via aerial photography, which is not as clear or accurate in this regional area. Therefore, this would not provide an accurate demonstration of the layout of the intersections.
2		All of the external traffic routes for the Project construction are approved B-double routes and as such currently safely accommodate the movement of these vehicles. The construction will use semi-trailers only for access which require less road pavement than a B-double for turning movements.
		The site access, which will utilise the access road for the Dumaresq substation adjacent to the Project site, was designed and built to accommodate the swept path movement of the heavy vehicles associated with the construction of the electrical substation and these included over size / over mass vehicles associated with the delivery of the substation unit. This would have used a vehicle that was larger than a B double and as such the site access will accommodate the swept path of a semi-trailer entering and exiting the site.
	update the project layout to include onsite parking and laydown areas;	The "Ancillary infrastructure" area is described broadly in the EIS as including a substation, battery storage and operations. Section 2.2.5 of the EIS also details the inclusion of the maintenance shed and car park area. The car park and laydown areas form part of the Ancillary Infrastructure area, previously indicated on relevant Project Figures. The car park area within this prescribed area, is shown in the snip below and in Figure 2-2 .
3		CAR PARK WORKSHOP SITE OFFICE - x SITE ACCESS POINT x x x x x x x x x x x x x x x x x x x

ERM

Bonshaw Solar Farm (SSD 9438) Response to RFI – May 2020 Page 5 of 8

Item	DPIE Comments	Details
		Similarly, the ancillary infrastructure area will be used as a laydown area, although during construction, a temporary car park and laydown area will be utilised in the northern portion of the Site (refer to Figure 2-2).
3a	Please include the site access, car park and laydown areas and Bruxner Highway label as per the Updated Project Layout (Figure 2.1 in letter dated 22 May 2020) into the current project layout (Figure 5.1 in the Submissions report, Figure 2.2. in Amendment Report) and provide a consolidated project layout.	Site layout figures have been updated to include the site access, car park and proposed laydown areas (refer to Figure 1-1).
4	confirm if the access road from the site access point to the proposed site buildings and ancillary infrastructure area is wholly located within the project boundary (including additional landowners consent if required);	Site access will be provided from the existing access road leading to the Dumaresq substation. A connecting road will then join at the northwestern corner of the Project Site to connect to the internal access road network. The internal road network will run around the parameter of the solar array, including running from the access point in the north-western corner, parallel to the existing Dumaresq access road, down to the ancillary infrastructure area. Landowner consent from Crown Lands, as owner of the site access road being utilised to gain access to the Project Site is provided in Attachment A of the AR. A second site access point is included as part of the Project at the Ancillary Infrastructure area in the western portion of the Site. This has been provided as an alternative emergency exist point.
5	provide the number of heavy vehicles required during operation;	Under standard operation, there will be no need for heavy vehicles. In the event that panels or an inverter requires replacing, a single B-Double truck would be required for delivery of materials. It is estimated that under a worst-case scenario, this may be required 2 to 3 times in a year.
Biodiv	ersity Assessment	
6	provide an addendum to the BDAR to include a species polygon and appropriate justification for the four species that will be impacted by the proposal, or justification for the species credits (the Department notes that only one species polygon has been prepared as per Figure 5.3 in the BDAR, and the area of impact to calculate the species credits is not consistent for all species)	The Biodiversity Development Assessment Report (BDAR) has been updated to address this comment (refer to Attachment C) and revised species polygons and credit reports are provided in Appendix L of the BDAR.
6a	Provide an addendum to the BDAR, addressing:	The Biodiversity Development Assessment Report (BDAR) has been updated to address these comments (refer to Attachment C) and

Item DPIE Comments

■ The species polygons for the Masked Owl and Barking Owl must be expanded to include each of the recorded hollow bearing trees (green dots) within the development site as identified in Figure 5.3 of the BDAR, because hollow bearing trees are an important part of the breeding and foraging habitat for these owls. The expanded species polygons do not need to include extensive areas of disturbed grassland.

- The Bristle-faced Free-tailed bat species polygon currently covers the entire development site, including the mapped disturbed grassland areas. We note the precautionary approach of including large areas of disturbed grassland for potential foraging habitat. However, due to the large area of disturbed grassland, we would also accept a reduced species polygon, that aims to capture the more viable bat habitat components, such as hollow bearing trees, similar to that described above for the owls.
- The amended species polygons above will need to also be reflected in the Biodiversity Assessment Method (BAM) calculator. We note the above changes will reduce the overall species credit requirement for the development.

Details

revised species polygons are provided in Appendix L of the BDAR.

- The updated species polygon for the Bristlefaced Free-tailed Bat conservatively includes all woodland PCT plus a 50m buffer to all hollow-bearing trees recorded within the Subject Land. Disturbed grasslands have been excluded from the updated species polygon as provided in Figure L-1.
- The updated species polygon for the Barking Owl includes a 100m buffer to all mapped hollow-bearing trees that contained hollows of a suitable size (i.e. >20cm wide) and height above the ground (i.e. >4m) in accordance with breeding habitat constraints provided in BAM-C. The updated species polygon for the Barking Owl is provided in Figure L-2.
- The updated species polygon for the Masked Owl includes a 100m buffer to all mapped hollow-bearing trees that contained hollows of a suitable size (i.e. >40cm wide). The updated species polygon for the Masked Owl is provided in Figure L-3.

As both the Masked Owl and Barking Owl are dual credit species, potential foraging habitat is addressed within the ecosystem credit requirement for PCTs associated with this species. Therefore, the updated species polygons have been revised to include potential breeding sites only as required in the BAM. This is still considered to represent a conservative overestimate of actual/potential breeding habitat in that all hollow-bearing trees of a suitable size have been included rather than a single confirmed nest tree.

Based on the updated mapping of the threatened species polygons, impacts to habitat of Threatened Species requiring offsets now include:

- Habitat for the Bristle-faced Free-tailed Bat requiring 827 species credits;
- Foraging habitat for Eastern Cave Bat requiring 1,030 species credits;
- Potential (assumed) breeding habitat for Masked requiring 104 species credits; and
- Potential (assumed) breeding habitat for Barking Owl requiring 283 species credits.

A total of 598 ecosystem credits are also required to offset the impacts of the project.

HК	M

Bonshaw Solar Farm (SSD 9438) Response to RFI – May 2020 Page 7 of 8

Item	DPIE Comments	Details		
Floodi	Flooding			
	provide further evidence that the project will not impact overland flows (including further consideration of drop-down fencing);	The Flood Impact Assessment prepared for the Project concluded that a minor increase to peak flood levels within the Site immediately upstream of the proposed vehicle crossings were observed, however, these impacts are contained within the channel and the Site boundary. This minor impact was modelled under the 1 in 100 Annual Exceedance Probability (1% AEP) flood event scenario.		
		There is no proposed filling of the floodplain, except for works associated with the vehicle crossings and the installation of fencing along the Project boundary.		
7		The fencing of the Project boundary will either utilise the existing fencing if appropriate, upgrade existing fencing or install new chain-mesh fencing.		
		Where fencing intersects with waterways, drop- down fencing will be adopted to mitigate any potential impact to overland flows. To ensure there is no debris build-up at these points, regular checks of the drop-down fencing will be incorporated into the Site's Environmental Management Plans.		
		Figure 2-3 below provides an indication of the location of drop-down fencing for the Site.		
Water	Water Supply			
8	provide further evidence of the ability of identified potential water suppliers for the project to meet the necessary water demands.	Water supply for the Project will be provided through a licensed water supplier. GAIA have been in discussions with three potential suppliers – Trident Water or Wade's Water in Warwick and Vital Water Service in Casino.		
O		Discussions are ongoing with water providers, however at this stage Trident Water have provided confirmation of their ability to supply the necessary water demands for the Project (refer to Attachment D)		

2.1 AHIMS #11-3-0083 Location Amendment

Comments were raised in the assessment process by BCD in relation to the 2018 survey effort being unsuccessful in locating AHIMS #11-3-0083. Although a concerted effort was originally made to locate AHIMS #11-3-0083, the survey was thought to be unsuccessful in relocating AHIMS #11-3-0083, with a number of newly recorded sites located in nearby areas.


Through the assessment process of the project, BCD undertook a technical assessment of all the information recorded regarding AHIMS #11-3-0083. The result of this assessment





Bonshaw Solar Farm (SSD 9438) Response to RFI – May 2020 Page 8 of 8

suggests that a technical error in the recording of coordinates has resulted in a misunderstanding and error about the location of AHIMS #11-3-0038. Having projected the original coordinates, BCD has determined that the area was actually identified during the 2018 survey and given a new site identifier.

Accordingly, all figures showing the location of AHIMS #11-3-0083 have been updated to reflect the true location of AHIMS #11-3-0083. The Aboriginal Cultural Heritage Assessment has been updated is provided in **Attachment E**.







3. **SUMMATION**

This response has been prepared to address items raised in the Department's request for information, dated 1 May 2020 and 27 May 2020. This response provides various updated details associated with the Bonshaw Project.

We trust the information provided is sufficient to address details requested by DPIE. Should you require any further details, please do not hesitate to contact the undersigned on (02) 4903 5535 or via email <u>Lachlan.Giles@erm.com</u>.

Yours sincerely,

Lachlan Giles

Clydes

Environmental Planner

Michael Rookwood Senior Planner

ERM		Bonshaw Solar Farm (SSD 9438) Response to RFI – May 2020 Page 13 of 8
ATTACHMENT A	UPDATED TRAFFIC IMPACT	ASSESSMENT



Proposed Solar Farm,
Bruxner Highway,
Bonshaw
ERM

Traffic Impact Assessment and Management Plan

Updated Access

Final Report

3rd June 2020



Solar Farm Project, Bonshaw NSW Traffic Assessment Report- access update

Author: Sean Morgan

Client: ERM

Issue: Final/3.06.2020

Reference: P1302

3 June 2020

Quality Review and Document History

Version	Date	Description	Prepared By	Checked By
Ver01	21/01/2020	Draft	S Morgan	C Thomas
Ver02	14/02/2020	Final	S Morgan	C Thomas
Ver03	26/03/2020	Final	S Morgan	C Thomas
Ver04	2/6/2020	Final	S Morgan	C Thomas



Ground Floor, 161 Scott Street, Newcastle NSW 2300 Ph (02) 4032 7979 www.secasolution.com.au

© Seca Solution Pty Ltd 2020

intended solely for the use of the client for the purpose for which is has been prepared. Use or copying of this document in whole or in part without the written permission of Seca Solution constitutes an infringement of copyright. The intellectual property contained in this document remains the property of Seca Solution.



Contents

1 Int	troduction	2
1.1	Consultation and Authority Requirements	6
2 Ex	xisting Road Network and Local Characteristics	9
2.1	Traffic Volumes and Road Operation	10
2.2	Road Safety	10
2.3	Accident History	19
2.4	Mitigation Measures	19
3 Cc	onstruction Activities	20
3.1	Timing	20
3.2	Working Hours	20
3.3	Construction staff numbers	21
3.4	Heavy vehicle requirements	21
3.5	Vehicle movements	23
4 Tr	affic Management Assessment	25
4.1	Impact Assessment	27
4.2	Cumulative impacts	29
4.3	Impacts on road pavement	30
5 Sı	ummary and Conclusion	32
Append	lix A. Safe Construction Actvities	33
Append	lix B. Drivers Code of Conduct	37
1.1	General Requirements	37
1.2	Vehicle Speeds	37
1.3	Driver Fatigue	37
1.4	Operating Hours	39
1.5	Transport Routes	39
1.6	Vehicle Departure and Arrival	41
1.7	Overtaking	41
1.8	Breakdowns and Incidents	42
1.9	Penalties and Disciplinary Action	42
1.10	Emergency Contact Numbers	42
1.11	Driver Declaration	43
Append	lix C. Accident Data	44
Append	lix D. Traffic Control Plan for Works at Site Access on Bruxner Highway	46
Append	lix E. Construction Traffic Management Plan (CTMP)	47



1 Introduction

Seca Solution has been commissioned by Environmental Resources Management (ERM) Pty Ltd to review the traffic impacts associated with the construction and operational phase of a new Solar Farm development and to determine traffic management measures associated with the construction activities for the project. The project involves construction, operation and eventual decommissioning of a 200 megawatt (MW AC) solar farm located in Bonshaw, to the west of Tenterfield in NSW.

The traffic impact has been previously assessed by Seca Solution and has been reviewed by the Road Authority (Inverell Shire Council), the RMS (now Transport for NSW) and the Department of Planning, Industry and Environment. From this review, the access route has been altered in accordance with the Council and RMS comments and additional information provided to satisfy the requirements of these various stakeholders.

The following works and infrastructure would be required to support the construction and operation of the solar farm:

- Construction of a main access road for all access and egress for the Site
- Installation of Electrical infrastructure including:
- A 132kV Substation including two transformers and associated 132kV switchgear.
- Inverters to collect and convert DC to AC.
- BESS/battery storage with up to 300 MW (AC);
- Cabling and other electrical infrastructure (e.g. security systems).
- A maintenance compound and buildings.
- Fencing, landscaping and environmental works.
- Construct site access from the existing access road leading to the Dumaresg substation. A connecting road will join to the access road, connecting to the internal access road network at the north-western corner of the Project Site.
- installation of an overhead grid connection to Dumaresq Substation

The operational life of the solar farm is expected to be 30 years at which point the panels are either replaced and operations continue, or the infrastructure is removed, and the site is decommissioned and rehabilitated.

Construction of the site will take approximately 12 months.

Power generated by the facility will be transmitted via existing 330kV transmission lines, in an easement owned by TransGrid that traverses the Site.

As part of the development consent, and prior to work on site, a Construction Traffic Management Plan will need to be approved by the road authorities (Inverell Shire Council and Transport for NSW). The busiest period associated with the development with regards to traffic is during construction, with the operational phase of the project only requiring between 6-10 staff on site for the majority of the time. Seca Solution has prepared this Construction Traffic Management Plan (CTMP) for the project to ensure traffic issues can be safely and efficiently managed during the construction activities on site.

This CTMP has been developed for the construction activity for the project and the potential decommissioning element for the project, which may occur in 30 years' time. The potential decommissioning of the project site will require a similar level of activity, although will probably require less staff and would be completed over a shorter timeframe. The requirements and protocols for the decommission stage of the project will be as per the construction





phase, although it is acknowledged these may need to be reviewed and altered in 30 years to suit the road conditions at that time as well as the work requirements.

The site is located within the locality of Bonshaw, west of Tenterfield and is shown in Figure 1-1 and 1-2 to follow.

The site is currently arable land and has road frontage to the Bruxner Highway along its northern boundary.




Figure 1-1: Site Location within the greater road network





Figure 1-2 – Location of Sunnyside Platform Road

SECAsolution




Figure 1-3 – Detailed site access and layout



Consultation and Authority Requirements

As part of the project, there has been consultation with the Department of Planning and Environment by the project manager and SEARs have been issued. A summary of the SEARs as they relate to traffic and access issues is presented below and the response is provided within this table.

OF A D	D /0 () (
SEARs issue	Response / Section of report
An assessment of the peak and average traffic generation including over-dimensional vehicle and construction worker transportation	The volume of traffic has been assessed for both the construction and operational phase. Construction: Peak staffing levels will require 65 light vehicles and 15 heavy vehicle inbound movements per day and similar outbound. For the sub-station installation there will be 3 over sized vehicles. Operational: 10 light vehicles per day inbound and outbound. Infrequent heavy vehicle for specific maintenance work only Distribution: Heavy vehicles via the designated heavy vehicle route to connect with the Bruxner Highway and New England Highway to east or west to Goondiwindi. Refer Section 2.3, 2.4 3.1.1
An assessment of the likely transport impacts to the site access route (including Bruxner Highway, Glenrock Road and Rocky Creek Road), site access point, rail safety issues, any Crown Land, particularly in relation to capacity and condition of the road	Heavy vehicles approaching the site will travel via the Bruxner Highway and New England Highway. For heavy vehicles exiting the site, they shall travel via the Bruxner Highway, then south through Ashford via Inverell – Bonshaw Road to Inverell then along the Gwydir Highway to the New England Highway at Glen Innes. No heavy vehicle access via Glenrock Road or Rocky Creek Road. Refer Section 3, Figure 3.1 Existing traffic flows on New England Highway, Bruxner Highway and Gwydir Highway are low and well within acceptable limits. Minimal impact created by traffic during construction and operations. Refer Section 4.1
A cumulative impact assessment of traffic from nearby developments (including cumulative impacts from Sundown Solar Farm, Sapphire Solar Farm and White Rock Solar Farm)	Sundown Solar Farm located off Gwydir Highway and currently EIS being prepared. Limited interaction with the Bonshaw site on Bruxner Highway. Sapphire Solar Farm constructed and operational. Limited interaction with Bonshaw site on Bruxner Highway. White Rock Solar farm under construction. Limited interaction with Bonshaw site on Bruxner Highway. Refer Section 4.2
A description of any proposed road upgrades developed in consultation with the relevant road authority and rail authorities (if required)	Access provided on Bruxner Highway via the existing access to the electrical substation in this location. This access will be upgraded to provide a left turn deceleration lane (AUL (s)) and a widened shoulder for eastbound traffic (BAR treatment) in accordance with Austroads Guidelines. No other road upgrades proposed.





A description of the measures that would be implemented to mitigate any transportation impacts during construction

Map of route for heavy vehicles provided Refer Section 3 Figure 3.1.

All drivers will sign code of conduct which specifies all road rules must be obeyed including driving through school zones - Refer Appendix B

All staff and delivery drivers will be inducted to site and sign a driver code of conduct – Refer Appendix B
The contractor on site shall establish a complaint

handling process and resolution process.

During construction activities nearby properties along the local haulage route on Bruxner Highway will be notified via a letter drop of construction work and timeframe – Refer Appendix A.

Transport for NSW consultation

Post exhibition of the project, discussion has been held with Greg Schiffe (GS) on 20th December 2019 from the RMS (now Transport for NSW) and the following points are noted:

- Agreed that the right turn out of Bruxner Highway onto the New England Highway is not suitable for the
 project. However, the left turn into this road is considered safe and appropriate. The right turn into the
 Bruxner Highway is also not considered appropriate for the project construction work;
- GS stated that the use of Sunnyside Platform Road is considered appropriate for the project, with this intersection providing a safe layout for vehicles including trucks turning right into Sunnyside Platform Road for access from the north.
- Accident data review is required for access between the site and the regional road network (being the New England Highway);
- If exit truck movements are to head south via Ashford to Inverell then this route needs to be reviewed and assessed for the impact of these trucks on this route
- Require clarification on the access location and GS indicated access via the existing access to the electric substation is the preferred option
- Require clarification on access routes for light and heavy vehicles. GS indicated that directing light vehicles to use the route via Ashford is not feasible due to the additional travel distance / time and directing light vehicles via Sunnyside Platform Road is appropriate.

RMS Consultation

Consultation has been held via a phone conversation with Andrew McIntyre, manager Land Use Assessment, Western Region (September 2017) with regard to a number of solar farms proposed to be constructed across rural NSW. The relevant outcome of the discussion with Andrew McIntyre is provided below:

- The critical phase for the assessment is the construction activities as this involves heavy vehicle access to the site along regional and local roads as well as a high number of workers;
- Consideration to the movement of staff to and from the site must be given. In remote areas where the
 solar farms are constructed, there are a large number of staff who can be drive in/drive out re-locating for
 temporary work from the established east coast centres such as Sydney and Newcastle. This requires
 staff to drive a long-distance home after working on the site for long hours for a week or more –
 consideration to controls for staff driving home after working on site should be considered;
- Provide details on the access routes to the site for heavy vehicles and the size / number of heavy vehicle movements associated with the construction and operation of the site;



- Provide details on the operational characteristics of the project it is recognised that the staff levels and traffic volumes for the operational stage of the project are low;
- Provide comment with regard to the decommissioning stage of the project and the potential traffic impacts;
- Prepare a driver code of conduct for the project to control vehicle access and maintain safety;
- Assess impacts on road safety, including pedestrians and cyclists and any bus routes impacted
- Review alternative transport options for the site including pedestrians, cyclists and bus use
- Provide details on any road upgrades identified as part of the project and include a Road Safety Audit as required





Existing Road Network and Local Characteristics

The Bruxner Highway is a state classified road, which runs to the north of the subject site with an east-west orientation providing connection between the New England Highway to the east and Boggabilla to the west where it connects with the Newell Highway. The south, east and west boundaries of the subject lands are defined by neighbouring agricultural lots with some sections of unnamed, unsealed rural roads. The Bruxner Highway is sealed (refer Photo 1 below) and provides a width of approximately 6 metres passing the site allowing for 2-way traffic movements as required. It operates under the posted speed limit of 100 km/h.

The Bruxner Highway connects with the New England Highway to the east of the site at a four way give way controlled intersection with the New England being the priority road and Old Ballandean Road being the opposite minor road.



Photo 1 – View along Bruxner Highway in the vicinity of the project site

The New England Highway is a state classified road that is a key freight route in NSW and forms part of the road network designated by the Roads and Maritime to carry oversize, over mass vehicles. It typically provides a single lane of travel in both directions and operates under the posted speed limit of 110 km/h outside of the urban areas where the alignment permits. As part of the state road network, the New England Highway carries a mixture of local, regional and inter-state traffic with a significant number of trucks including B-double combinations. The Cunningham Highway operate in a similar manner providing key transport routes between Ipswich and the New England Highway at Warwick.

To the south of the site is the **Gwydir Highway**, which is a state classified road that is a key freight route in NSW and forms part of the road network designated by Transport for NSW to carry oversize, over mass vehicles. It typically provides a single lane of travel in both directions and operates under the posted speed limit of 100 km/h outside of the urban areas where the alignment permits. As part of the state road network, the Gwydir Highway carries a mixture of local, regional and inter-state traffic with a significant number of trucks including B-double combinations.



Bonshaw Road is a local road managed by Inverell Shire Council, located to the west of the site. It is a sealed two-way road with an overall width in the order of 7 metres. It intersects with the Bruxner Highway via a simple give way controlled intersection with the Bruxner Highway being the priority road. This road continues south and connects with Ashford Road in Ashford to provide a road link through to Inverell. This route provides a consistent road standard and forms part of the approved B-double road network in NSW. Both of these road provide a single lane of travel in both directions and operate under the posted speed limit of 100 km/h.

Sunnyside Platform Road is a local road providing an approved B-double connection between the New England Highway and Bruxner Highway. It provides a pavement width in the order of 7 metres wide permitting 2-way traffic movements.

As part of the project, it is proposed that all heavy vehicles will travel via the roads identified above. Local supplies could be sourced from Goondiwindi or Tenterfield as well as accommodation for workers associated with the project.

Traffic Volumes and Road Operation

Traffic volumes in the immediate vicinity of the subject site are very low, reflective of the rural environment. The Bruxner Highway carries relatively low traffic flows, reflective of its rural setting with a mixture of local traffic as well as regional traffic demands. Observations on site during a typical morning period (Tuesday 11th September 2018) shows that the current road network in the vicinity of the subject site operates very well with no delays. The route proposed to be used for the project carries low traffic flows and operates with no delays except for those associated with drivers slowing down to observe traffic flows on the approaches to the various intersections and negotiating the intersections.


Transport for NSW webpage provides traffic data on the Bruxner Highway at Mingoola (station Id 91170), approximately 15 kms east of the subject site. The traffic data from 2011 shows that the daily traffic flow was 213 vehicles per day with around 23% heavy vehicles, reflective of rural demands in this location. It is considered that there has been limited growth in traffic since this time and as such the current daily traffic flows are considered to be similar.

The same web page shows that in 2011 the daily traffic flow on the New England Highway to the immediate north of Tenterfield (station Id 91577) was 2421. It is considered that there has been limited growth in this area since 2011 and as such the daily traffic flows would be similar. In 2012 the traffic flows on Bonshaw Road were 232 vehicles per day northbound. Assuming southbound flows to be the same would give daily flows in the order of 500 vehicles per day. It is considered that these flows would not have altered much since 2012.

For the Gwydir Highway, the same web page shows that the daily flows to the east of Inverell were 1,317 in 2019 whilst to the west of Glen Innes the flows on the Gwydir Highway in 2019 were 1,442.

2.2 Road Safety

It is recognised that as part of the project work, there will be a significant number of heavy vehicle movements associated with construction which may impact the local road network. All inbound heavy vehicle access to the project site will be via the Bruxner Highway and the New England Highway. For traffic from Goondiwindi, the route will be via the Newell Highway to town and along the Bruxner Highway. Inbound items such as the solar panels will be via the New England Highway (northbound from the Port of Newcastle) or via the Cunningham Highway connecting with the Port of Brisbane. No alternative route for inbound heavy delivery vehicles has been considered as this route is appropriate.





For outbound empty truck movements, movements to Goondiwindi would be the same, as would truck movements to the Brisbane area which shall use Sunnyside Platform Road to then turn left on the New England Highway. However, for trucks heading back towards Newcastle, the right turn out of the Bruxner Highway onto the New England Highway is not considered appropriate due to its poor visibility and intersection layout it represents a road safety risk for the project. As such, trucks wishing to head south towards Newcastle shall use the route via Inverell - Bonshaw Road to connect to Inverell and the Gwydir Highway. This is via the approved B-double road network and shall have an acceptable impact upon road safety.



Photo 2 – Sheltered turn lane for right turning vehicles into Sunnyside Platform Road from the New England Highway. Note straight alignment that allow for good visibility





Figure 2-1– aerial photo showing layout of New England Highway and Sunnyside Platform Road showing straight alignment of New England Highway and sheltered turn lane (source: google maps).

The major road safety impact is associated with the delivery trucks accessing the site and their impact upon the operation of the intersections. The trucks carrying the solar panels and other specialist materials will be accessing the site from either the Port of Newcastle or the Port at Brisbane, to which the solar panels shall be shipped. The trucks will then access Bonshaw via the regional road network which will include the New England Highway and the Cunningham Highway if from Brisbane. These regional roads currently provide a high standard of road and allow for the movement of local, regional and national road freight and carry B-double trucks. It is considered that the additional truck movements associated with the construction activities for the project will have a minimal and acceptable impact upon road safety along these roads.

For the sub-station installation there is a requirement for 3 over sized vehicles to access the site. These will require a separate approval and permit through the National Heavy Vehicle Regulator with Transport for NSW approval. Safety requirements and impacts are assessed as part of the permit application and the necessary safety controls applied which can include escort vehicles and Police vehicle support.

The nominated vehicle route currently caters for a large number of heavy vehicles including B-double combinations. This route provides a wide road pavement and in major built up areas cater for kerb side parking and the safe 2-way movement of trucks along the road. The intersections along this route are well laid out and provide good visibility in all directions to allow for the safe turning movements of vehicles. It is considered that this route through the various towns can safely accommodate the additional traffic movements associated with the project.





For the local traffic impacts, consideration has been given to the existing alignment of the road, intersection layouts, current traffic flows and existing users along the Bruxner Highway. Observations on site with regard to road safety are summarised below:

- Existing traffic flows on the local road are very low
- The sealed width of the road allows for two-way traffic movements
- The alignment of the road is generally good
- There are a number of sub-standard curves where there are advisory signs provided in advance with 55 and 65 km/h speed limit guidance signs
- A number of heavy vehicles were observed travelling along the Bruxner Highway during the site work, associated with local farm requirements as well as general deliveries in the area. These included semitrailers.

These routes are provided below (Figure 2-1) and will be included within the Driver's Code of Conduct which will form part of the project inception meeting for the project for all staff and drivers.




Figure 2-2 – Designated Heavy Vehicle route to project site






Figure 2-3 – Detail for truck connection from north between New England Highway and Bruxner Highway. Allows for inbound and outbound connection

Intersection of Bruxner Highway and New England Highway

The intersection of the Bruxner Highway and New England Highway is 4-way cross road with the New England Highway being the priority road. Give way signs delineate the controls for the side road. This intersection has been upgraded with new line marking that provides a sheltered right turn lane for traffic turning off the New England Highway into the Bruxner Highway from the north as well as a right turn lane for traffic from the south turning into Old Ballandean Road. It allows for all turning movements and as part of the main transport route in this area caters for the swept path of B-double combinations. This intersection is located within a posted speed zone of 100 km/h. The sight line requirements for drivers approaching the intersection have been assessed against the requirements of Austroads Guidelines. For the posted speed limit of 100 km/h the sight distance requirement is 248 metres. To the right, looking south this sight line is available for drivers. However, to the left (north) the sight line is restricted due to the vertical alignment of the road, which creates a safety concern. In addition, for a truck turning right out of this road onto the New England Highway, there is a single lane southbound which creates further safety concerns for the project. A truck turning right out of the Bruxner Highway could then be a hazard for a southbound vehicle leading to rear end accidents. In a 100 km/h speed zone this is not considered to be acceptable.

For the right turn into the Bruxner Highway off the New England Highway this visibility restriction also creates concerns with potential for rear end type accidents.





Photo 3 – View to left for a driver exiting the Bruxner Highway onto the New England Highway.



Photo 4 – View northbound on the New England Highway on approach to intersection with the Bruxner Highway. Note good forward visibility and dual lanes to allow a through vehicle movement to pass a left turning vehicle into the Bruxner Highway.



For all other movements at the intersection of the Bruxner Highway and the New England Highway the existing intersection controls are considered to be appropriate for the traffic movements associated with the project site. For traffic from the south on the New England Highway, the forward visibility is good allowing a driver to adjust their vehicle speed to cater for a vehicle in front turning left onto the Bruxner Highway. The traffic flows on the New England Highway in this location are relatively low and as such this will create minimal delays for other drivers.

With the proposed restrictions in turn movements for the construction traffic at this intersection, no upgrade works are required at this intersection to accommodate the traffic movements associated with the proposed solar farm (both construction and operation phase).

To mitigate the issue associated with the right turn out of the Bruxner Highway into the New England Highway, all trucks leaving the site wishing to head south shall instead turn left out of the site onto the Bruxner Highway, then turn left onto Bonshaw Road to head south to Ashford continuing south on Ashford Road to Inverell via the approved B-double route. These vehicles shall then travel along the Gwydir Highway to Glen Innes to connect to the New England Highway. This will eliminate right turn movements allowing for left turn movements only.

For all other intersections along the routes, the intersections are well laid out and provide good visibility for road users. As such these other routes, which are all approved for use by B-doubles, will provide a safe and appropriate route for heavy and light vehicles associated with the project site,

2.2.2 Site access on Bruxner Highway

The access to the site will be via the existing access provided to the electric substation adjacent to the subject site. This access was constructed to allow for the construction activities associated with the electric substation and is now used for maintenance purposes only for the substation. This access will be upgraded to provide a left turn deceleration lane (AUL(s)) and shoulder widening to provide an AUR treatment, both in accordance with Austroads Guidelines.

The design of this upgrade will be reviewed with the road authority through the approval process.



Photo 5 View on Bruxner Highway showing layout of access connection to existing electric substation.







During the construction work, there will be some heavy vehicles arriving to the site from Goondiwindi, west of the site. These movements would be associated with the supply of equipment for the construction work. For traffic approaching from the west, the forward sight distance to the site access is approximately 700 metres. Austroads Guidelines require a sight distance of 248 metres for the posted speed limit of 100 km/h and 300 metres for a speed limit of 110 km/h. With the forward sight distance available, a driver can adjust their vehicle speed if required on the approach to the site access if there is a vehicle propped waiting to turn right into the site. However, given the very low traffic flows on the Bruxner Highway, it is not considered that vehicles turning right into the site will need to prop on the highway with the only delays created by vehicles slowing down to negotiate the site access driveway. Safety will be maintained with a Traffic Control Plan to reduce vehicles speeds on the Bruxner Highway during this construction work with reduced vehicle speeds and truck turning advisory signs.

Trucks exiting the site will all turn left onto the Bruxner Highway. With the low traffic flows on the Bruxner Highway, this can safely occur as there are large gaps in the through traffic movements in this location. For trucks turning left out of the site, the sight distance available to the right (east) exceeds 700 metres allowing a driver to safely exit the site and travel along the highway. No left turn acceleration lane is required for this movement.



Photo 6 – View to right along Bruxner Highway for drivers exiting the project site





Photo 7 – view to left for vehicles exiting the access to the electric substation




Figure 4 – AUR type right turn treatment proposed at site access (same as old BAR type control)




Figure 5 – AUL type left turn treatment proposed at site access







2.3 Accident History

A review of accident data provided by TfNSW (Appendix C) for the period July 2014 to June 2019 has determined the vast majority of accidents along the proposed heavy vehicle route involved single vehicles (74%). Speeding was also determined as a contributing factor for over half of the accidents recorded, with the most common crash type being vehicles going off-road. In reviewing the data, it can be seen that speeding is a major factor in the number of accidents recorded. As discussed above the route and major intersections have been assessed on road, with no inherent road safety issue identified.

2.4 Mitigation Measures

From the details above the following mitigation measures are proposed.

- Provide a temporary TCP on the site frontage on the Bruxner Highway, adjacent to the site access, for
 construction work associated with upgrading the access and for traffic entering and exiting the site. This
 TCP shall only be in place during construction and signs shall be removed or covered outside of
 construction activities on the site. Once the construction work is complete this TCP shall be fully removed.
 This TCP will be prepared in accordance with "Traffic Control at work sites" published by the RMS dated
 July 2018. This TCP is provided in Appendix D to this report;
- Provide regular community updates for residents along the Bruxner Highway in the vicinity of the site to advise of construction activities and increased heavy vehicle movements along this road;

2.4.1 Light Vehicle Route

For light vehicles associated with workers, the proposed access route will be via the designated heavy vehicle route shown in Figure 2-1 above. This route provides a safe and acceptable route for light vehicles which can safely and conveniently access the site. For light vehicles leaving the site and heading towards Tenterfield or Glen Innes (via the New England Highway), these drivers will be required to turn left onto Sunnyside Platform Road and then turn right from this road onto the New England Highway, to avoid the existing safety concerns identified at the intersection of the Bruxner Highway and New England Highway. This shall be documented through the Driver Code of Conduct for the construction staff.

Light vehicle access could also be from Goondiwindi associated with construction workers via the Bruxner Highway.

The project will be utilising workers local to the site from the main centres e.g. Tenterfield, Goondiwindi, Inverell who will use this route as well as other local roads to connect between Inverell and the site. Additional specialist staff may be required, and these staff members would be located in accommodation in Tenterfield, Inverell and Goondiwindi.



Construction Activities

The construction and commissioning phase is expected to last approximately 12 months with expected commencement in late 2020. The main construction activities would include:

- Site preparation: geotechnical investigations to confirm ground conditions; site survey to confirm allotment boundary, riparian zone, and infrastructure positioning and placement; installation of fencing, internal access tracks, establishment of foundations and hardstands; office and car parking area;
- Construction activities: including installation of mounting structures and tracker tubes; securing PV modules to tracker tubes; installation of cabling and switching station, establishment of BESS / battery storage and maintenance compounds and associated site infrastructure; and testing and commissioning;
- Plant and Equipment: will include earthmoving plant and equipment for site preparation and clearing; cable trenching and laying equipment; pile drive equipment; forklifts and cranes; water truck for dust suppression and machinery equipment for construction of BESS / battery storage and associated facilities.

The project does not require any concrete footings to be provided for the solar panels construction.

A site office and compound will be established on site for the duration of the works with temporary access tracks provided to allow for access as required across the site.

Staffing levels at peak construction activities is expected to be 190 personnel on site maximum. The demand for staff numbers will be lower at the commencement of the project and shall increase overtime to the maximum of 190 staff. This demand is also expected to decrease towards the end of the project.

All staff vehicles will be able to park within the site adjacent to the site office with no external parking demands. The car park area will allow for up to 80 vehicles to park within this compound area. The size of the overall site footprint however will allow for all construction staff vehicles to park on site. As part of the project construction it is proposed to maximise the local workers content and car-pooling will be encouraged and supported as part of these trips. With 2 or 3 people arriving in a single vehicle it can be seen that the parking demands can be contained within the site.

The access to the site will be via the existing access constructed for the adjacent electric sub-station. This access road was previously used for the construction works associated with this substation and is now only used for access for maintenance work on the sub-station. With the vast majority of the heavy vehicles accessing the site from the east (via the New England Highway) it is proposed to provide a left turn deceleration lane for the site access to allow for a safe entry for the construction traffic. With limited demand for access from the west of the site, there is no requirements for a sheltered right turn lane to be provided at the access point.

3.1

The construction of the solar farm is expected to commence in 2020 and be completed within a 12 month timeframe.

Working Hours

Construction hours are in accordance with the Interim Construction Noise Guidelines (DECC 2009) (ICNG) with standard construction hours being

- 7:00am and 6:00pm Monday to Friday
- 8.00 AM to 1.00 PM on a Saturday







No construction work is to be carried out on a Sunday or public holiday.

No construction work, upgrading or decommissioning activities will be undertaken outside of these hours with the exception of:

- The delivery of material as requested by the NSW Police Force to other authorities for safety reasons; or
- Emergency work to avoid the loss of life, property and / or material harm to the environment.

3.3 Construction staff numbers

Peak demand levels for the construction work will vary with a peak of 190 people for a 6 month duration and a lower level outside of this peak period. The staff will be sourced locally where appropriate with some specialist and project management staff being temporarily located in Tenterfield, Inverell and potentially Goondiwindi. Staff will be encouraged and supported to carpool as appropriate with other staff transferred to and from the site via mini coaches to reduce vehicle demands. Due to the size of the site footprint, these same vehicles will also be used on site to move staff across the site.

With a peak of 190 staff, a vehicle occupancy rate of 3 people per vehicle has been assumed based upon carpooling and the use of a mini bus e.g. Toyota Hiace. This would give around 65 vehicle movements inbound and outbound for staff movements during this peak construction activity. Either side f this peak the staff levels will be lower and hence light vehicle numbers will correspondingly decrease.

All light construction vehicles will be able to park on site within the office compound or across the site as required. The parking on site will be informal and located across the site to suit the construction work. Due to the overall footprint of the site the vehicles on site will be required to carry staff and equipment across the site to the construction activity area (which varies across the project) and as such the vehicles are not all parked in one location for the duration of the project.

Given the overall size of the project site all parking demands can be managed on site with no impact upon the external road network.

3.4 Heavy vehicle requirements

The number of heavy vehicles accessing the site will vary across the project timeframe. At the beginning of the project there will be a requirement for some earth moving equipment to construct the access road and some minor earthworks across the site as required. This may require a scraper or bulldozer which will be transported to site on a low loader. This machinery will remain on site for the duration of the earthworks portion of the project construction work.

While extensive earthworks are not proposed, some land forming (including localised cut and fill areas) may be undertaken to achieve more consistent gradients beneath the PV modules. Additionally, earthworks are required for trenching works.

In total:

- Approximately 15,000 m³ of gravel would be required to cap the access road
- Approximately 15,000 m³ of sand (subject to detailed design) would be required for the bedding of cables that are to be buried throughout the site

Should any excavated material not be suitable for reuse or additional fill material required, the maximum amount of fill is estimated to be 12,000 m³.

Once the earthworks have been completed, the balance of the construction work will commence requiring machinery including:

• Pile driver (20)



- Piling rig
- All terrain fork-lift (20)
- All terrain utility vehicles (10)
- Backhoe (10)
- Flatbed trucks (10)
- Mobile crane (1)

Other equipment if required may include an elevated work platform, scraper, roller and winches. All of the plant will be located on site and will therefore be only required to access the site once for the construction works.

The solar panels are expected to be delivered from either the Port of Newcastle or Port of Brisbane. Other specialist equipment is generally sourced from Newcastle or Brisbane as required whilst consumables such as concrete and general material supplies will be sourced locally from the Tenterfield area.

A summary of the expected vehicle movements associated with the construction work is provided below and shows the full movements for the duration of the project. These movements are spread out across the project, with the site set up and earthworks commencing at the beginning of the project. Once this work is complete, the balance of the construction work will commence with deliveries of the specialist equipment etc along with the import of backfill material being over a number of weeks to suit the construction timeframe.

These traffic numbers are based on the concept design work for the project and could alter through the detailed design phase of the project.





3.5 Vehicle movements

A summary of the vehicle movements is provided below.

Phase	Purpose	Vehicle Type / Trailer Type	No. of one-way vehicle movements
	Portacabin delivery and removal	Low loader	20
	Skip delivery and removal	Low loader	40
Site Set-Up and	Generator delivery and removal	Semi-trailer	4
Demobilisation	General deliveries	Semi-trailer	40
	Crane mobilization and demobilization	Crane	4
	Water tank delivery and removal		4
	Delivery of imported capping for road laydowns and crane hardstands	Truck and dog	500
Roads and	Plant delivery and removal: excavators, compactors drill rig	Low loader	40
hardstands	Concrete deliveries for maintenance container hardstands	Concrete agitator	120
	Tool container delivery and removal	Low loader	4
Concreting	Module deliveries Semi-trailer		2000
Generating	Mounting structure and pile deliveries	te deliveries for maintenance container hardstands agitator Intainer delivery and removal Ideliveries	
Equipment	Inverter Station deliveries	Low loader	3
	DC cabling trays and combiner boxes	Semi-trailer	400
AC Cable Installat	AC Cable delivery	Semi-trailer	400
ion	Backfill material delivery	Dump Truck	1800
Plant delivery and r emoval	Telescopic handler and excavator	Low loader	50
	Conductor delivery	Semi-trailer	25
Overhead Line	Pole deliveries	RAV	6
	Pole dressing delivery	Semi-trailer	2
	Miscellaneous deliveries	Light vehicle	40
Other	Monitoring equipment fibre SCADA servers etc	Truck	2
Otriei	Waste Collection	Truck	400
	Consumables (Oil and Fuel)	Truck	40
		TOTAL	7,544



In summary, typical vehicle movements during the peak construction period (over 6 months) are in the order of 65 light and 20 heavy vehicles two-way (65/20 inbound, 65/20 outbound) per day. For the light vehicles, the vast majority of these will be inbound movements in the morning bringing workers to the site with these vehicles then remaining on site for the full working day before leaving at the end of the working day. It is expected that there will be limited light vehicle movement outside of these periods, other than support staff e.g. office staff or the occasional visitor to the site.

For the heavy vehicles, these will typically be spread across the working day. For the solar panel deliveries, these trucks are arriving from either the Port of Brisbane or the Port of Newcastle and the journey length will be over 5 or 7 hours respectively, seeing a spread of these vehicles not all arriving at the same time. Allowing for each truck to be emptied on site one at a time, the outbound movements will also be spread out and not all leave at the same time. All other heavy vehicles will also be spread out across the normal working day with no concentration of heavy movements expected.

Outside of the peak period of construction, the staff levels will be lower and the daily light vehicle numbers will be less than 65 inbound and outbound per day. The heavy vehicle numbers will also be lower outside of the peak construction activity and less than 20 vehicles inbound and outbound per day.





4 Traffic Management Assessment

The proposed traffic management measures allow for all access off the Bruxner Highway only. The access to be used will be for the construction traffic movements as well as the future on-site operational demands. This access is to be provided in accordance with the requirements for the site operations (including swept path requirements for delivery vehicles) and take into account the design requirements of Inverell Shire Council.

The designated access route to the site will be used by both light and heavy vehicles.

All vehicle movements in and out of the site are as shown below in Figure 4-1.




Figure 4-1 – Heavy and Light Vehicle access route to subject site






Figure 4-2 – detail for connection between New England Highway and Bruxner Highway



4.1 Impact Assessment

The project will require the delivery of the solar panels and other specialist equipment from Newcastle or Brisbane with the access route via:

- Newcastle metropolitan regional road network
- Hunter Expressway / New England Highway
- New England Highway to Bruxner Highway turn off
- Bruxner Highway to site access
- Bonshaw-Inverell Road
- Gwydir Highway
- Brisbane metropolitan regional road network
- Cunningham Highway
- New England Highway to Bruxner Highway.

These roads all form part of the road freight routes within the State road network and all are approved for heavy vehicle movements including B-double access for the full length of the routes. These routes will be documented as the Haulage Route for all delivery vehicles associated with haulage of the solar panels and other specialist materials for the project site.

These roads carry a high number of heavy vehicles, including B-doubles associated with local and regional agricultural demands. These agricultural demands are seasonal in nature and occur 24 hours a day often involving night travel and operations. There are a number of farms in the general locality of the project site as well as in the wider area that use these local and regional roads during these seasonally high demand periods. Due to the seasonal nature of this work and the requirement for quick turnaround of crop deliveries it is considered that it is not appropriate to limit truck movements for these farms. Similarly, it is considered that it is not appropriate to limit truck movements to and from the project site at these times as the traffic movements on the local roads will continue to remain low.

The roads proposed to be used for trucks accessing the site all currently accommodate the swept path requirements for B doubles to access the site. The site access on the Bruxner Highway was previously used for constructing the electric sub station on the site and accommodates the swept path movements of B doubles as well as Over Size Over Mass (OSOM) vehicles that were required as part of the construction for this electric substation.

For the regional road network e.g. New England Highway / Hunter Expressway, Bruxner Highway, Cunningham Highway the total traffic flows will remain well within acceptable limits and as such will continue to operate to a good level of service and accommodate all road users.

The traffic flows along the local roads giving access for the heavy and light vehicle movements associated with the project are currently very low based on-site observations. Therefore, during the peak construction period and peak staff / material demands, the additional 65 light vehicle movements associated with the staff movements and 20 daily truck movements (per direction) will have a minimal and acceptable impact upon the operation of these local roads during construction. Once operational, the traffic movements are much lower with a maximum of 10 staff on site per day and as such the impact will be negligible.

There is minimal background traffic growth in this location. Transport for NSW count data from the station located between Tenterfield and Glen Innes (Station I.D. DNDSTC) shows traffic flows of 2,201 in 2012 and 2,091 in 2019, with minor fluctuations up and down for the intervening years. Other counts along the regional road network show similar or lower increases in values. For the assessment of the future impacts in 10 years-time, it can be seen that the site at that time will be operational with 10 staff located on the site. The impact of these ten staff will be negligible on the local and regional road network.



The site is expected to be operational for more than 10 years so that the impact of the decommissioning of the site cannot be assessed in detail at this stage. The site could remain operational beyond 10 years and the impact will remain low beyond the 10 year design horizon.

There will be no public vehicle access within the work site during the construction works, with a fence provided at the commencement of the project along the entire site boundary. This fence will remain once the project is constructed for security purposes with a locked gate to be provided at the site access off the Bruxner Highway.

There will be no pedestrian access to the site for the general public. There are no pedestrian paths in the locality of the site nor expected demands in this remote rural area so there will be no impacts for pedestrians created by the project works.

There is a school bus that runs along the Bruxner Highway to Bonshaw in the morning with the return trip in the afternoon. There may be some deliveries occurring in the morning and afternoon during the school bus operation, depending on the stage of the construction work and travel requirements for the deliveries. All staff will be on site prior to the morning bus run and will depart site after the afternoon bus run. As such it is considered that there is very limited interaction with the school bus. On the regional and state road network all school zones will be delineated in accordance with RMS Guidelines with reduced speed limits in accordance with normal NSW road rules. All drivers associated with the project construction work will adhere to the road rules as applicable and will be advised of the school bus operation on the Bruxner Highway.

There will be no impact upon public transport services with no diversions required. There are no bus stops impacted upon by the proposal.

There will be minimal impact for emergency vehicles and heavy vehicles with no diversions required.

There will be minimal impact upon any other development within the locality of the site.

There will be minimal impact upon adjoining Council areas. Traffic routes in and out of the locality will be along the arterial road network which will experience minimal impacts due to the works.

There are no residential dwellings in the immediate locality of the site access that will be impacted upon by the project and construction work. There are a number of residences along the heavy and light vehicle access route. The residents along the Bruxner Highway will be notified in writing of the construction works and the activities as required.

Construction vehicle movement on internal roads could lead to dust generation. A water truck will be used for dust suppression to minimise the production of dust, with the amount of water spreading adjusted accordingly to respond to the conditions. Additionally, any significant deposits of dirt and other construction materials will be promptly removed from public roadways.

Post construction, the traffic numbers generated by the project are very low, with staffing levels varying daily with a maximum on-site workforce of 10 people on any one day. There will not be any need for regular heavy vehicle access to the site once the solar farm is operational except for the occasional heavy vehicle for emergency repairs or irregular maintenance.







4.2 Cumulative impacts

A search of the Major Projects Register on the DPE website was undertaken together with the requirements of the SEARs for the project. The following projects are in the council area that may add to cumulative impacts.

- Sundown Solar Farm
- Sapphire Solar Farm
- White Rock Solar Farm

Project	Cumulative construction Impacts	Cumulative operational Impacts
Sundown Solar Farm (EIS currently being prepared)	Site located off Gwydir Highway between Inverell and Glen Innes. No overlap with construction along Gwydir Highway. If constructed at same time cumulative impact along New England Highway. As a state highway there is adequate capacity to accommodate these vehicles movements.	Operational traffic would be expected to be less than 10 light vehicles per day in the morning and afternoon and no impact with traffic for Bonshaw Solar Farm site.
Sapphire Solar Farm	Site located off Gwydir Highway and currently under constructed. This site will be fully constructed before the Bonshaw Solar Farm commences construction.	Operational traffic would be expected to be less than 10 light vehicles per day in the morning and afternoon and no impact with traffic for Bonshaw Solar Farm site.
White Rocks Solar Farm	Construction complete.	Operational traffic would be expected to be less than 10 light vehicles per day in the morning and afternoon and no impact with traffic for Bonshaw Solar Farm site.

4.2.1 Delivery vehicles

The majority of the deliveries for the project will be via 19 metres semi-trailers and B-doubles. The access routes along the regional / state road network to the site are all along approved B double routes.

Delivery vehicles would be required throughout the project period. The travel time between the ports (Newcastle or Brisbane) and the site for the solar panels is approximately 5 to 7 hours and these deliveries will be spaced out over the construction period, to minimise the impact upon the road network and to reduce the need to store the panels on site. Other deliveries will include the metal structures for the solar panels, sand and gravel for the foundations and internal tracks and cabling. There will also be some deliveries of specialist equipment such as photovoltaic boxes or skids and delivery stations.

The trucks associated with the delivery of the supplies will all travel along the State and regional road network. There are a number of schools located along these routes, however all have marked school zones and speed limit restrictions as per State guidelines. As these routes are all on the State and regional road network it can be seen that heavy vehicles currently operate on these roads safely. It is considered that the additional truck movements associated with the construction work will result in no noticeable impact upon road safety adjacent to these schools.

There is no requirement to divert traffic as part of this construction work.



The substation will require 3 over sized vehicle deliveries to the site. These will require a separate approval and permit through the National Heavy Vehicle Regulator with RMS approval. Safety requirements and impacts are assessed as part of the permit application and the necessary safety controls applied which can include escort vehicles and Police vehicle support.

4.2.2 Construction staff movements

For the construction work, the staffing levels will peak at 190 on site and as part of the project, staff will be encouraged and supported to carpool as part of the Code of Conduct for the project and use mini buses provided to allow for shared trips to the site from shared accommodation in Tenterfield, Glen Innes and Goondiwindi. There could be 65 vehicles inbound in the morning associated with on-site staff and a similar number departing at the end of the working day. Either side of the peak demand for construction staff the light vehicle demands associated with staff movements will be less than 65 inbound and outbound per day.

The site is in a rural location well away from the local towns and as such it is considered that there will be no pedestrian or cyclists accessing the site in conjunction with staff movements.

The vehicle numbers associated with the construction work are relatively low and it is considered that the movement of vehicles in and out of the site for construction works can occur in a safe manner. No limitation on truck access times is considered appropriate for the project. Given the journey length between the port and the subject site, the vehicles as they are approaching the site will be spread out reducing the impact of the arrivals. With unloading of vehicles taking 30 minutes or more, trucks exiting the site will also be spread out.

Impacts on road pavement

A protocol will be provided for both undertaking dilapidation surveys and making any necessary repairs following construction to Bruxner Highway to within 200 metres to both sides of the site access. It is considered that there could be some impact here due to the turning movements and braking / accelerating of trucks. Beyond these limits it is considered that the impact on the state road will be minor and shall not impact upon the overall pavement construction.

With regards to any emergency repairs required within the above zone, the contractor on site would contact the relevant authorities and will ensure the road is safe. Repairs will be made in accordance with the relevant authority standard.








Figure 4-3 Dilapidation Assessment Protocol



Summary and Conclusion

The project allows for the construction of a solar farm in the locality of Bonshaw in northern NSW located off the Bruxner Highway. The construction work is planned to commence in late 2020 and will take approximately 12 months to construct. During construction there will be a requirement for a significant number of trucks to access the site for material delivery as well as light vehicles associated with construction staff. All vehicle access will be directly off the Bruxner Highway with the majority of deliveries expected to be via the New England Highway, with the solar equipment arriving via the Port of Newcastle or Port of Brisbane.

The trucks accessing the site will all travel along the regional and state road network, which currently carries heavy vehicle movements including B-doubles. The site is located in a rural setting and as such the hourly and daily traffic flows along the Bruxner Highway and the New England Highway in this location are relatively low. As such there is considerable spare capacity to cater for the additional traffic movements associate with the project construction stage. During peak construction activities, the project will generate 65 inbound light vehicle movements in the morning associated with the construction staff with a similar number of vehicles leaving the site in the late afternoon. At these peak activity times, the site will also generate approximately 20 trucks inbound and outbound per day, associated with the delivery of material and specialist equipment to the site. Outside of the peak construction activity the light and heavy vehicle movements will be lower per day.

Post construction, the operational traffic demands are very low with around 10 people working on the site. There will also be the requirement for occasional heavy vehicle movements, associated with on-going maintenance for the facility.

The access to the site will be via the existing access constructed for the adjacent electric sub-station. This access road was previously used for the construction works associated with this substation and is now only used for access for maintenance work on the sub-station. With the vast majority of the heavy vehicles accessing the site from the east (via the New England Highway) it is proposed to provide a left turn deceleration lane (AUL(s)) for the site access to allow for a safe entry for the construction traffic. Shoulder widening will also be provide to cater for the limited demand for right turning traffic into the site with a AUR type upgrade.

A review of the access route shows that the layout of the intersection of the Bruxner Highway and the New England Highway does not safely cater for the right turn out movement. The sight distance for this right run is restricted and the width of the New England Highway in this location does not allow for a run off area for vehicles. It is proposed that as part of the construction traffic management plan, southbound empty trucks leaving the site shall turn left onto the Bruxner Highway then proceed south via Bonshaw-Inverell Road to Inverell then along the Gwydir Highway to connect to the New England Highway via Glen Innes.

All access routes for the heavy vehicles associated with the delivery of materials to the site are approved for Bdouble use and carry B-doubles associated with local rural demands.

A Traffic Control Plan will be in place during construction work at the site access to ensure safety for road users and construction workers is managed in an appropriate manner. A Construction Traffic Management Plan has been prepared for the project (see below) to ensure that road safety is managed for all workers associated with the project as well as existing road users.

The overall conclusion for the project shows that the construction traffic can safely and efficiently access the site with minimal impact for existing road users. The management plan for the construction traffic access ensures that the trucks accessing the site shall have an acceptable impact on the road network and safety concerns at the intersection of the Bruxner Highway and the New England Highway are address through the drivers code of conduct. Once operational, the traffic demands are minimal and shall have little impact upon the local road network.





Appendix A. Safe Construction Activities

The contractor on site is responsible for the management of all traffic in connection with its activities and the construction works conducted on the site. The Contractor will provide all traffic management, safety warnings and signage including such persons as necessary to direct traffic, as required by AS 1742:2009 – Manual of uniform traffic control devices.

External traffic movements

The Contractor will:

Ensure traffic management controls are established, maintained and monitored to underpin the safety of workers, other personnel and the general public

Establish traffic management controls in consultation with relevant stakeholders

Ensure traffic management controls comply with regulatory and legislative requirements

Ensure traffic management controls comply with the contract

Ensure traffic management controls maintain the flow of traffic within the site and on surrounding public roads

Reinstate any areas affected by the temporary construction access requirements to their original condition

The primary drivers for determining the traffic management controls during the construction period are:

- Safety of personnel, the general public and construction workers
- Minimising impact (if any) on operations
- Contractual requirements (including site access)
- Road traffic authority and local government requirements
- OHS requirements in relation to the movement of all vehicular traffic and pedestrians either within or adjacent to sites
- Environmental management requirements
- The impact construction traffic has on the local community in the surrounding area, and
- The need to meet construction requirements (including any schedule and cost constraints)

The traffic management controls will be communicated to appropriate stakeholders which will include the local community in the site vicinity via a letter box drop.

The Contractor will ensure:

Any significant deposit of dirt and other materials caused by construction traffic and other operations (in relation to the works) will be promptly removed from existing public roadways

Suitable precautions are taken to ensure no rock is dislodged onto any roadway from construction vehicles

Construction plant and equipment do not park on or within the pavement or shoulders of any existing trafficked roadway

Construction vehicles (when loaded) comply with the mass, loading and access requirements of the road traffic authority

Construction traffic will cause the least possible obstruction to public and other traffic



Directional signage will be installed to direct construction traffic and warn other motorists of construction traffic.

This signage is positioned in accordance with the approved Traffic Control Plans.

All drivers will be provided with a copy of the access routes to and from the site as part of their induction for the project;

A Vehicle Movement Strategy has been developed to eliminate the impact on local roads arising from additional construction traffic (e.g. solar panel delivery vehicles). The Vehicle Movement Strategy directs all drivers to access the site from the Bruxner Highway to eliminate the impact on the local roads. There is no requirement to restrict the direction of flow and/or time of day for movements.

The Contractor will comply with any client or Road Traffic Authority signage requirements for traffic control. Where construction work is to be undertaken either on or adjacent to a public roadway that is open to traffic, the work must be undertaken in accordance with all regulatory and legislative requirements that govern the movement of vehicles and pedestrians on any public roadway.

Within the Worksite

All employees, subcontractors, suppliers and any other persons connected with the project must adhere to all such Statutory Requirements and comply with all lawful directions. Any breach of such requirements may result in disciplinary action of the persons concerned.

The maximum speed limits within the Worksite are:

- 40 kph on formed roads
- 20 kph during foggy/dusty conditions with headlights on
- 10 kph when passing pedestrians

The Contractor will manage access to and from the site by all employees, subcontractors, suppliers and any other persons connected with its activities and the works; and all occupants within the worksite and through each area of the site.

The Contractor shall provide for safe and continuous operation of normal pedestrian and vehicular traffic along all roads, pedestrian paths and vehicular access to the worksite and must provide and maintain all necessary watchmen, lights, barriers, notices and signs.

The Contractor will not unnecessarily obstruct any side road, branch track, drain or watercourse and will not break down or remove any fences or gates without prior notification to the client. If unavoidable, the Contractor will remove such obstruction or repair such breakage as soon as possible, or as directed by the Client.

A Vehicle and Traffic Management Procedures briefing will be included in the Project Site Induction.







Pedestrian Traffic

The Contractor may encounter pedestrian traffic at and near to the site. The Contractor will ensure that sites are appropriately isolated and secured from unauthorised entry; and that the Site is appropriately sign-posted and controlled. Given the location of the site it is considered that any pedestrian activity will be negligible.

Site Construction Traffic

Traffic within the Site will be managed in accordance with the Site Management Plan. The Sites Layout Plans will indicate site access and egress points and detail any required separation of construction plant and personnel. These plans will be communicated during Tool Box Meetings and/or Daily Pre-start Meetings.

The Site Layout Plan will incorporate details of parking arrangements for the site construction workers, speed limits within the construction works or through access roads established for vehicular and plant construction traffic.

The Sites Layout Plan will detail traffic management controls that are appropriate within each site.

Traffic controls shall be regularly reviewed for effectiveness and will be amended to maintain or improve a safe work environment. Traffic management controls established for sites will be inspected at weekly intervals to verify that a safe work environment is being maintained. Records of inspections shall be maintained.

Access Roads and Site Movement

Unless sign-posted otherwise, load limits on public roads adjoining the sites apply within them.

If required the Contractor shall request approval from the client prior to any over-dimensional load, or load in excess of load limits entering the site, or using the roads within the site.

All workers must travel to and from the site via the nominated access roads.

Parking

All workers must park in the Designated Parking Areas as specified in the Site Management Plan. The Contractor shall ensure no persons (in connection with its activities) parks in any other area of the site or in any other area without prior written consent.

Monitoring, Measurement and Review

The purpose of Monitoring and Measurement is to ensure that all construction works, including subcontracted activities, are being performed in accordance with the contract requirements, statutory requirement and in a controlled and safe environment. Ongoing monitoring and audit of Traffic Management procedures and the worksite implementation of traffic control shall be conducted.

Audits of the Traffic Control measures under differing operating conditions are to be carried out including during overcast and rainy weather, at night or at any other restrictive times where conditions may change in accordance with the requirements of AS1742.3.

Results of audits, inspections and improvements are to be reported in the reporting cycle of the contract to enable assessment of the adequacy of the implementation of the Traffic Control within contract performance and system review meetings.



Inspection and Auditing of Traffic Control Plan (TCP)

Regular Site Inspections by designated supervisory and field staff of worksite protection are to be arranged on a daily frequency depending on the complexity of traffic control on the site.

Site Inspections will be carried out and the following Traffic Management Forms completed:

Traffic Control Daily Checklist Traffic Control Weekly Checklist

A daily record of the inspections should be kept. This should include:

- When traffic controls were erected
- When changes to controls occurred and why the changes were undertaken
- Any significant incidents or observations associated with the traffic controls and their impacts on road users or adjacent properties
- Where significant changes to the work or traffic environment or adverse impacts are observed, the controls should be reviewed as a matter of urgency.

The monitoring program should generally incorporate inspections:

- Before the start of work activities on site
- During the hours of work
- Closing down at the end of the shift period

The inspection program shall be adjusted to suit changing circumstances and/or risk environment such as during times of increased traffic flows or speeds, contra-flow arrangements or when changed controls are introduced.

The Audits of the implemented Traffic Management features will be undertaken following setup in accordance with the TCP and prior to the TCP being put into service.





Appendix B. Drivers Code of Conduct

1.1 General Requirements

All vehicles / drivers accessing the site must:

- i) Be registered and hold a valid driver's licence for the class of vehicle being operated;
- ii) Operate the vehicle in a safe and appropriate manner whilst travelling to / from the site or when operating within the site. This includes obeying all New South Wales state road rules.
- iii) ALL heavy vehicles must adhere to the designated heavy vehicle routes as far as practical;
- iv) NO trucks shall turn right from the Bruxner Highway onto the New England Highway.
- v) ALL trucks heading south shall turn left out of the site then proceed via Bonshaw-Inverell Road to Inverell then travel along the Gwydir Highway to Glen Inness to connect with the New England Highway.
- vi) Comply with the directions of authorised personnel when operating within the site and obey any relevant signage installed along the internal roads.
- vii) Not use a mobile phone while operating any vehicle.
- viii) Must always wear a seatbelt when operating any vehicle.

1.2 Vehicle Speeds

Drivers shall observe the posted speed limit along the designated transport route and adjust their vehicle speed as required to suit the road environment and prevailing weather conditions. Vehicle speeds must be appropriate to ensure the safe movements of the vehicle with consideration to the vehicle configuration.

Maximum speeds limits within the project site shall be as follows:

- i) 40 km/hr along formed roads.
- ii) 20 km/hr during foggy / dusty conditions. Headlights must be on.
- iii) 10 km/hr when passing pedestrians or any plant equipment.

1.3 Driver Fatique

Drivers shall not be permitted to operate a vehicle or plant equipment when impaired by fatigue. If you suspect that you or someone else is experiencing fatigue, please inform your supervisor.

Operators of heavy vehicles shall be aware of the requirements relating to fatigue as outlined in the Heavy Vehicle National Law. Drivers shall also be aware of their adopted fatigue management scheme (shown below) and ensure that they are operating within its requirements.

- i) Standard Hours of Operation
- ii) Basic Fatigue Management (BFM)
- iii) Advanced Fatigue Management (AFM)



Basic Fatigue Management (single driver)

Time	Work	Rest
In any period of	A driver must not work for more than a maximum of	And must have the rest of that period off work with at least a minimum rest break of
6 1/4 hours	6 hours work time	15 continuous minutes rest time
9 hours	8 1/2 hours work time	30 minutes rest time in blocks of 15 continuous minutes
12 hours	11 hours work time	60 minutes rest time in blocks of 15 continuous minutes
24 hours	14 hours work time	7 continuous hours stationary rest time*
7 days	36 hours long/night work time**	No limit has been set
14 days	144 hours work time	24 continuous hours stationary rest time taken after no more than 84 hours work time and 24 continuous hours stationary rest time and 2 x night rest breaks taken on consecutive days.

Advanced Fatigue management:

The seven principles are grouped into three categories:

Work-related rest breaks (such as short rest breaks):


- 1. Reduce the time spent continuously working in the work opportunity
- 2. The more frequent breaks from driving, the better

Recovery breaks (such as major rest breaks):

- 1. Ensure an adequate sleep opportunity in order to obtain sufficient sleep
- 2. Maximise adequate night sleep
- 3. Minimise shifts ending between 00:00-06:00
- 4. Minimise extended shifts

Reset breaks (such as long periods of rest or extended leave):

- 1. Prevent accumulation of fatigue with reset breaks of at least 30hrs (and include two night periods, 00:00 06:00) between work sequences
- ALL details relating to fatigue management for delivery vehicles are covered by the National Heavy Vehicle Regulator





1.4 Operating Hours

Construction

Construction is to be in completed in accordance with the *Interim Construction Noise Guideline* (DECC 2009) which defined standard construction work hours as:

- Monday to Friday: 7am to 6pm
- Saturday: 8am to 1pm
- Sunday and Public holidays: No work

The following construction, upgrading and decommissioning activities may be undertaken outside these hours without the approval of the secretary:

- The delivery of materials as requested by the NSW Police Force or other authorities for safety reasons;
- Emergency work to avoid loss of life, property and / or material harm to the environment.

Vehicle movements shall be undertaken during standard construction hours (or just before to allow workers to get to site). Oversize vehicles up to 26 metres long may require access to the site after hours however this would be subject to the requirements of Roads and Maritime, Dubbo Regional Council or NSW Police.

Normal Operations

Daily operations and maintenance by site staff would be undertaken during standard working hours:

- Monday to Friday: 7am to 6pm
- Saturday: 8am to 1pm
- Sunday and Public holidays: No work

During normal operations, all vehicle movements shall be undertaken during the standard operating hours (or just before to allow workers to get to site). There may be a requirement for vehicles to access the site after hours during an emergency however these would be infrequent.

Vehicles which arrive at the site prior to commencement of working hours shall have the engine turned off to minimise noise impacts on surrounding residences.

1.5 Transport Routes

All vehicles must travel to and from the project site via the route as shown below (Figure 1).






Figure 1 - Transport route to/from the site for ALL 19 m semi-trailers or greater








Figure 2 - Transport route to/from the site for ALL 19 m semi-trailers or greater (detail for northbound via New England Highway to use Sunnyside Platform Road)

1.6 Vehicle Departure and Arrival

Heavy vehicles departing the site shall have a minimum 5 minute separation to reduce the impacts upon the local road network.

Always maintain a minimum separation of at least 50 metres between vehicles when travelling within the site.

Drivers must contact the site supervisor upon arrival and await further instructions or direction before proceeding.

Drivers must also report to the site supervisor prior to departure.

All vehicles must enter and exit the site in a forward direction. Vehicles are to have clean tyres upon exiting the site to prevent dirt being tracked onto the public road network.

ALL 19 metres semi-trailer or greater must turn left out of the site and proceed via Bonshaw-Inverell Road to Inverell then via the Gwydir Highway to Glen Innes and the New England Highway when heading south.

1.7 Overtaking

Overtaking shall not be permitted within the site unless the intention to overtake has been communicated to the driver of the leading vehicle and consent to overtake granted.



1.8 Breakdowns and Incidents

Heavy Vehicles

In the case of a breakdown, the vehicle must be towed to the nearest breakdown point as soon as possible. All breakdowns must be reported to Transport for NSW Transport Management Centre on 131 700 and the vehicle protected in accordance with the Heavy Vehicle Drivers Handbook. The relevant shift manager on site shall also be notified.

If a breakdown occurs on-site please remain inside your vehicle, notify the shift manager of your location and await further instruction.

If you are involved in an accident, please notify the shift manager immediately and contact emergency services if required.

Light Vehicles

In the case of a breakdown, ensure that the vehicle is secure, notify the shift manager of your location and await further instruction.

If you are involved in an accident, please notify the shift manager immediately and contact emergency services if required.

1.9 Penalties and Disciplinary Action

Any driver who fails to comply with the above requirements will have their details recorded and may be subject to disciplinary action.

1.10 **Emergency Contact Numbers**

i)	RMS Transport Management Centre	131 700
ii)	Queensland Traffic	131 940
iii)	Inverell Shire Council	(02) 6728 8288
iv)	NSW Police Service	(02) 6722 0599
v)	Site Office	
vi)	Shift Manager on Duty	





1.11 Driver Declaration

I, the undersigned, hereby agree to abide by this Driver Code of Conduct for the transport of equipment or personnel to / from the Bonshaw Solar Farm, located off Bruxner Highway in the general locality of Bonshaw, NSW. I have read and understand the requirements outlined in the attached document and will, to the best of my ability, comply and assist with their implementation, requirements or ongoing administration.


The subject document to which this declaration relates is included as part of this overall document and signing of this declaration confirms that the signee has read and understood their requirements as outlined throughout.

Driver Details				
Full Name				
Organisation				
Signature				
Date				

Full Name Signature Date



Appendix C. Accident Data





Transport NSW **Summary Crash Report** for NSW # Crash Type Crash Movement CRASHES 42 CASUALTIES 36 **Contributing Factors** 24 57.1% Intersection, adjacent approaches 2.4% Fatal 0 0.0% Killed 0 0.0% Car Crash Speeding 22 52.4% **Light Truck Crash** 21.4% Head-on (not overtaking) 2 Serious ini. 14 33.3% Seriously ini. 14 38.9% 14.3% Fatigue 6 Rigid Truck Crash 4.8% Opposing vehicles; turning Moderate inj. 10 23.8% Moderately inj. 12 33.3% U-turn Minor/Other inj. 16.7% Minor/Other ini. Articulated Truck Crash 9.5% 2.4% 10 27.8% (6) (14.3%) Weather Rear-end 3 7.1% Uncategorised inj. 0 'Heavy Truck Crash 0.0% Uncategorised inj. 0 0.0% Non-casualty 11 26.2% Fine 36 85.7% Lane change 0 Unrestrained 2.8% **Bus Crash** 0 0.0% 0.0% "Heavy Vehicle Crash (6) (14.3%) Rain 4 9.5% Parallel lanes; turning 0 0.09 Belt fitted but not worn. No restraint Self Reported Crash 8 19.05% fitted to position OR No helmet wom Overcast 2 Emergency Vehicle Crash 0 0.0% 4.8% Vehicle leaving driveway 2.4% Crashes Casualties Fog or mist 0 0.0% Overtaking; same direction Motorcycle Crash 16.7% 2.4% Time Group % of Day Other 0 2018 Pedal Cycle Crash 2.4% 0.0% Hit parked vehicle 0 0.0% 8 g 00:01 - 02:59 2.4% 12.5% 6 9 2017 Pedestrian Crash 2.4% Hit railway train 0.0% Road Surface Condition 03:00 - 04:59 2.4% 8.3% Rigid or Artic. Truck " Heavy Truck or Heavy Bus 10 2016 6 Hit pedestrian 2.4% 05:00 - 05:59 2.4% 4.2% Wet 8 19.0% # These categories are NOT mutually exclusive 8 2015 10 Permanent obstruction on road 0.0% 34 06:00 - 06:59 0.0% 4.2% 81.0% Dry 7 2014 Location Type Hit animal 2.4% 07:00 - 07:59 2.4% 4.2% 0 Snow or ice 0.0% *Intersection 13 31.0% Off road, on straight 2 4.8% 08:00 - 08:59 2.4% 4.2% Non intersection 29 69.0% Off road on straight, hit object 14.39 **Natural Lighting** 09:00 - 09:59 9.5% 4.2% 4 Out of control on straight 0.0% * Up to 10 metres from an intersection 10:00 - 10:59 7.1% 4.2% 3 Dawn 2 4.8% Off road, on curve 14.3% 11:00 - 11:59 3 7.1% 4.2% Daylight 32 Collision Type 76.2% Off road on curve, hit object 11 26.2% 12:00 - 12:59 7.1% 4.2% 3 Single Vehicle 31 73.8% Dusk 4 9.5% Out of control on curve 2 4.89 13:00 - 13:59 2 4.8% 4.2% Multi Vehicle McLean Periods % Week 11 26.2% Darkness 4 9.5% Other crash type 4 9.5% 14:00 - 14:59 4.8% 4.2% 17.9% 4.8% Speed Limit 9.5% 4.2% 15:00 - 15:59 Road Classification 4.8% 7.1% 40 km/h or less 4 9.5% 80 km/h zone 4 9.5% 16:00 - 16:59 14.3% 4.2% Freeway/Motorway 0 0.0% 11 26.2% 17.9% 50 km/h zone 2.4% 0 0.0% 90 km/h zone 17:00 - 17:59 2 4.8% 4.2% State Highway 3 7.1% 2.4% 3.5% 22 60 km/h zone 11 26.2% 100 km/h zone 52.4% 18:00 - 18:59 2.4% 4.2% Other Classified Road 37 88.1% 5 11.9% 3.6% 70 km/h zone 0 0.0% 110 km/h zone 0 0.0% 19:00 - 19:59 9.5% 4.2% Unclassified Road 4.8% 7 16.7% 10.7% 20:00 - 21:59 3 7.1% 8.3% ~ 07:30-09:30 or 14:30-17:00 on school days ~ 40km/h or less 9 21.4% G 16.7% 7.1% ~ School Travel Time Involvement 22:00 - 24:00 0 0.0% 8.3% 11.9% Day of the Week 7.1% 12.5% Street Lighting Off/Nil 2.4% % of Dark Monday 6 14.3% Wednesday 4 9.5% Friday 9 21.4% Sunday 9 21.4% WEEKEND 13 31.0% 10.7% 1 2.4% 29 69.0% Tuesday 6 14.3% Thursday 4 9.5% Saturday 4 9.5% WEEKDAY 4 in Dark 75.0% #Holiday Periods 2 4.8% New Year 0 0.0% Easter 0 0.0% Queen's BD 0 0.0% Christmas 1 2.4% Easter SH 0 0.0% Sept./Oct. SH 1 2.4% Aust. Day 0 0.0% Anzac Day 0 0.0% Labour Day 2 4.8% January SH 1 2.4% June/July SH 3 7.1% December SH Crashid dataset SECA Crashes Bruxner Way New Enland to Gwydir Highway 1.7.14 to 30.6.19

Reporting yrs 1996-2004 & 2019 Q3 onwards contain uncategorised inj crashes.

Percentages are percentages of all crashes. Unknown values for each category are not shown on this report.


Note: Crash self reporting, including self reported injuries began Oct 2014. Trends from 2014 are expected to vary from previous yrs. More unknowns are expected in self reported data.



Appendix D. Traffic Control Plan for Works at Site Access on Bruxner Highway

Access controls for on-site construction (upon completion of construction of site access). Distance D = 100 metres

TCP 195







Appendix E. Construction Traffic Management Plan (CTMP)



Proposed Solar Farm,
Bruxner Highway,
Bonshaw
ERM

Construction Traffic Management Plan

> Final Report February 2020



Bonshaw Solar Farm, Bonshaw NSW Construction Traffic Management Plan

Author: Sean Morgan Client: ERM Australia

Issue: Ver02

Reference: P1302 14 February 2020

Quality Review and Document History

Version	Date	Description	Prepared By	Checked By
Ver01	20/01/2020	Draft	S.Morgan	C.Thomas
Ver02	14/2/2020	Final	S.Morgan	C.Thomas





Contents

1	lr	troduction	1	2
	1.1	Exis	ting Road Network and Local Characteristics	3
	1.2	Traf	fic Volumes and Road Operation	5
2	С	onstructio	n Works	6
	2.1	Timi	ng	6
	2.2	Wor	king Hours	7
	2.3	Con	struction staff numbers	7
	2.4	Con	struction Traffic	7
3	S	ite Operat	ions	8
4	Т	raffic Man	agement	g
	4.1	Traf	fic Management Objectives	g
	4.2	Veh	cle Movement Plan	g
	4.3	Traf	fic Diversions	11
	4.4	Traf	fic Impacts	11
5	Е	nvironme	ntal Conditions	13
6 Incident Procedures			13	
7 Complaints		13		
		cess	13	
9	С	ompliance	Documentation	13
10 Traffic Control Plan		trol Plan	14	
	10.1	Gen	eral	14
	10.2	Exis	ting Traffic Conditions	14
	10.3	Cyc	ists and Pedestrians	14
	10.4	Gen	eral Traffic Control Considerations	14
	10.5	Traf	fic Control – Signage and Line Marking	14
	10.6	Com	pliance	14
	10.7	Dail	/ Checklist	15
	10.8	Con	tractors Contact Details	16
	10.9	TCF	Approval	16
Αį	ppen	dix A. S	te Plan	17
Αį	ppen	dix B. T	CP at site access on the Bruxner Highway	18
A	ppen	dix C. D	river Code of Conduct	19
	Driv	er Declara	ıtion	25



Introduction

The impact of the proposed Bonshaw Solar Farm has been assessed and submitted for review by the authorities. From this review Transport for NSW have requested a Construction Traffic Management Plan (CTMP) be prepared for the project. This CTMP has been prepared based upon the information available at the time of this submission, however, may be altered by the construction company for the future construction. Any changes to this CTMP in the future will be submitted to the road authorities for review and approval.

Due to the nature of works involved, the CTMP includes details of the required Traffic Control Plans (TCPs) to ensure that the safety for road users is preserved throughout the construction works.

The following CTMP has been prepared allowing for the construction of the Bonshaw Solar Farm project. Details of traffic management associated with decommissioning of the site has not been considered and shall be addressed prior to the decommissioning of the site to reflect the road and traffic conditions at this future time.

Bonshaw Solar Farm is located off the Bruxner Highway in the general locality of Bonshaw and approximately 65 kms west of Tenterfield, as shown in Figure 1-1.




Figure 1-1 - Site Location and Road Network







Figure 1-2 - Location of Sunnyside Platform Road

1.1 Existing Road Network and Local Characteristics

The **Bruxner Highway** is a state classified road, which runs to the north of the subject site with an east-west orientation providing connection between the New England Highway to the east and Boggabilla to the west where it connects with the Newell Highway. The south, east and west boundaries of the subject lands are defined by neighbouring agricultural lots with some sections of unnamed, unsealed rural roads. The Bruxner Highway is sealed (refer Photo 1 below) and provides a width of approximately 6 metres passing the site allowing for 2-way traffic movements as required. It operates under the posted speed limit of 100 km/h.

The Bruxner Highway connects with the New England Highway to the east of the site at a four way give way controlled intersection with the New England being the priority road and Old Ballandean Road being the opposite minor road.





Photo 1 – View along Bruxner Highway in the vicinity of the project site

The New England Highway is a state classified road that is a key freight route in NSW and forms part of the road network designated by the Roads and Maritime to carry oversize, over mass vehicles. It typically provides a single lane of travel in both directions and operates under the posted speed limit of 110 km/h outside of the urban areas where the alignment permits. As part of the state road network, the New England Highway carries a mixture of local, regional and inter-state traffic with a significant number of trucks including B-double combinations. The Cunningham Highway operates in a similar manner providing key transport routes between Ipswich and the New England Highway at Warwick.

To the south of the site is the **Gwydir Highway**, a state classified road that is a key freight route in NSW and forms part of the road network designated by Transport for NSW to carry oversize, over mass vehicles. It typically provides a single lane of travel in both directions and operates under the posted speed limit of 100 km/h outside of the urban areas where the alignment permits. As part of the state road network, the Gwydir Highway carries a mixture of local, regional and inter-state traffic with a significant number of trucks including B-double combinations.

Bonshaw Road is a local road managed by Inverell Shire Council, located to the west of the site. It is a sealed twoway road with an overall width in the order of 7 metres. It intersects with the Bruxner Highway via a simple give way controlled intersection with the Bruxner Highway being the priority road. This road continues south and connects with Ashford Road in Ashford to provide a road link through to Inverell. This route provides a consistent road standard and forms part of the approved B-double road network in NSW. Both of these roads provide a single lane of travel in both directions and operate under the posted speed limit of 100 km/h.

Sunnyside Platform Road is a local road providing an approved B-double connection between the New England Highway and Bruxner Highway. It provides a pavement width in the order of 7 metres wide permitting 2-way traffic movements.

As part of the project, it is proposed that all heavy vehicles will travel via the roads identified above. Local supplies could be sourced from Goondiwindi or Tenterfield as well as accommodation for workers associated with the project.







1.2 Traffic Volumes and Road Operation

Traffic volumes in the immediate vicinity of the subject site are very low, reflective of the rural environment. The Bruxner Highway carries relatively low traffic flows, reflective of its rural setting with a mixture of local traffic as well as regional traffic demands. Observations on site during a typical morning period (Tuesday 11th September 2018) shows that the current road network in the vicinity of the subject site operates very well with no delays. The route proposed to be used for the project carries low traffic flows and operates with no delays except for those associated with drivers slowing down to observe traffic flows on the approaches to the various intersections and negotiate the intersections.

Transport for NSW webpage provides traffic data on the Bruxner Highway at Mingoola (station Id 91170), approximately 15 kms east of the subject site. The traffic data from 2011 shows that the daily traffic flow was 213 vehicles per day with around 23% heavy vehicles, reflective of rural demands in this location. It is considered that there has been limited growth in traffic since this time and as such the current daily traffic flows are considered to be similar.

The same web page shows that in 2011 the daily traffic flow on the New England Highway to the immediate north of Tenterfield (station Id 91577) was 2421. It is considered that there has been limited growth in this area since 2011 and as such the daily traffic flows would be similar. In 2012 the traffic flows on Bonshaw Road were 232 vehicles per day northbound. Assuming southbound flows to be the same would give daily flows in the order of 500 vehicles per day. It is considered that these flows would not have altered much since 2012.

For the Gwydir Highway, the same web page shows that the daily flows to the east of Inverell were 1,317 in 2019 whilst to the west of Glen Innes the flows on the Gwydir Highway in 2019 were 1,442 vpd.



Construction Works

The construction and commissioning phase is expected to last approximately 12 months with expected commencement by late 2020. The main construction activities would include:

- Site establishment and preparation for construction:
 - Installation of security measures including fencing.
 - o Establishment of site compound and material layout areas. This shall include the provision of suitable on site parking for vehicles and workers.
 - Ground preparation.
- Installation of environmental controls:
 - o A detailed Construction Environmental Management Plan (CEMP) would outline the environmental controls required.
- Minor vegetation clearing (grasses, shrubs and isolated trees):
 - Targeted clearance of low laying vegetation around trenching areas to steel post installation to minimise disturbance to existing ground cover.
 - Establishment of tree and vegetation protection measures as required.
 - Clearance of larger vegetation such as bushes and isolated trees.
 - Establishment of additional sedimentation and erosion controls as required.
- Preliminary civil works including:
 - Drainage works
 - Setting up foundations for the substation
 - Earthing works (see below)
- Installation of steel post and rail foundation system for the solar panels.
- Installation of PV panels and DC wiring beneath the panels.
- Installation of underground cabling (trenching) and installation of inverter stations.
- Testing of electrical infrastructure
- Removal of temporary construction facilities and rehabilitation of disturbed areas.

The project does not require any concrete footings to be provided for the solar panels construction.

A site office and compound will be established on site for the duration of the works with temporary access tracks provided to allow for access across the site as required. Internal roads shall be constructed as all-weather roads.

All staff vehicles will be able to park within the site adjacent to the site office and across the site as required. The construction site is relatively large and the vehicles will be parked across the site located near the active work site, with no external parking demands created by the project. As part of the project construction it is proposed to maximise the local workers content (from Tenterfield and Glen Innes as well as potentially Goondiwindi) and carpooling will be supported as part of these trips. With 3 or 4 people arriving in a single vehicle it can be seen that the parking demands can be contained within the site.

2.1 **Timing**

Construction of the solar farm will commence in 2020 (subject to approval) requiring approximately 12 months to complete.





2.2 Working Hours

Construction hours are in accordance with the *Interim Construction Noise Guidelines* (DECC 2009) (ICNG) with standard construction hours being

- 7:00am and 6:00pm Monday to Friday
- 8.00 AM to 1.00 PM on a Saturday
- No construction work is to be carried out on a Sunday or NSW public holiday.

No construction work, upgrading or decommissioning activities will be undertaken outside of these hours with the exception of:

- The delivery of material as requested by the NSW Police Force to other authorities for safety reasons; or
- Emergency work to avoid the loss of life, property and / or material harm to the environment.

2.3 Construction staff numbers

Peak demand levels for the construction work will vary with a peak of 190 people for a 6 month duration and a lower level outside of this peak period. The staff will be sourced locally where appropriate with any specialist and project management staff from outside of the local area to be housed in Tenterfield and Glen Innes as well as potentially Goondiwindi.

Details of staff accommodation and employment shall be addressed within an Accommodation and Employment Strategy to be provided to Inverell Shire Council.

2.4 Construction Traffic

2.4.1 Light Vehicles

A shuttle bus service shall be provided to transport staff housed in Tenterfield and Glen Innes as well as potentially Goondiwindi to the site and to discourage private vehicle use. Shuttle buses will typically be 12 or 22 seater mini buses which will arrive at the start of the and leave at the end of the day. The route for these shuttle buses shall allow for collection at local hotels / motels to be determined once workers have been housed.

Specialist trades which require the use of their own vehicle for the transportation of tools and other equipment shall be encouraged to car pool where appropriate.

Local staff who reside outside of Tenterfield and Glen Innes as well as potentially Goondiwindi shall be permitted to drive to the site, however shall be encouraged and supported to car pool where appropriate to minimise demands for light vehicle travel to the site.

With a peak of 190 staff, a vehicle occupancy rate of 3 people per vehicle has been assumed based upon carpooling and the use of a mini bus e.g. Toyota Coaster. This would give 65 vehicle movements inbound and outbound for staff movements.

All construction light vehicles will be able to park on site as required.

2.4.2 Heavy Vehicles

The demands for heavy vehicles accessing the site will vary throughout the construction phase. At the beginning of the project there will be a requirement for some earthwork moving equipment to construct the access and some minor earthworks across the site as required. This may require a scraper or bull dozer which will be transported to site on a low loader. This machinery will remain on site for the duration of the earthworks portion of the project construction work.

While extensive earthworks are not proposed, some land forming (including localised cut and fill areas) may be undertaken to achieve more consistent gradients beneath the PV modules. Additionally, earthworks are required for trenching works.



In total, approximately:

- Approximately 15,000 m³ of gravel would be required to cap roads within the site
- Approximately 13,000 m³ of sand (subject to detailed design) would be required for the bedding of cables that are to be buried throughout the site

Should any excavated material not be suitable for reuse or additional fill material is required, the maximum amount of fill is estimated to be 12.000 m³.

Once the earthworks have been completed, the balance of the construction work will commence requiring machinery including:

- Pile driver
- Piling rig
- All terrain fork-lift
- All terrain utility vehicles
- Backhoe
- Flatbed trucks
- Mobile crane

Other equipment if required may include an elevated work platform, scraper, roller and winches. All of the plant will be located on site and will therefore be only required to access the site once for the construction works.

The solar panels are expected to be all delivered from the Port of Newcastle or Port of Brisbane. Other specialist equipment is generally sourced from Newcastle or Greater Brisbane as required whilst consumables be local from the Tenterfield and Glen Innes as well as potentially Goondiwindi.

Typical vehicle movements during the peak construction period (over 6 months) are in the order of 65 light and 20 heavy vehicles two-way (65/20 inbound, 65/20 outbound) per day. For the light vehicles, the vast majority of these will be inbound movements in the morning bringing workers to the site with these vehicles then remaining on site for the full working day before leaving at the end of the working day. It is expected that there will be limited light vehicle movement outside of these periods, other than support staff e.g. office staff or the occasional visitor to the site.

For the heavy vehicles, these will typically be spread across the working day. For the solar panel deliveries, these trucks are arriving from either the Port of Brisbane or the Port of Newcastle and the journey length will be over 5 or 7 hours respectively, seeing a spread of these vehicles not all arriving at the same time. Allowing for each truck to be emptied on site one at a time, the outbound movements will also be spread out and not all leave at the same time avoiding platooning of heavy vehicles on the road network. All other heavy vehicles will also be spread out across the normal working day with no concentration of heavy movements expected.

Outside of the peak period of construction, the staff levels will be lower and the daily light vehicle numbers will be less than 65 inbound and outbound per day. The heavy vehicle numbers will also be lower outside of the peak construction activity and less than 20 vehicles inbound and outbound per day

The construction company on site shall keep records of the number of heavy and over-dimensional entering and leaving each day.

Site Operations

Once constructed the site will operate for approximately 25 years at which point the site will either be decommissioned, or panels replaced to extend the length of operations. The site is expected to provide employment for no more than 10 staff with traffic demands associated with the ongoing operations to be significantly lower than those during construction.





4 Traffic Management

4.1 Traffic Management Objectives

The objectives of the traffic management plan are to:

- Minimise disruptions and ensure the safety of traffic on the external road network.
- Provide suitable access to the site for construction workers, heavy and over-dimensional vehicles.
- Ensure the safety and protection of workers working within the public domain.

4.2 Vehicle Movement Plan

The proposed traffic management measures allow for all access off the Bruxner Highway only, via the existing access to the adjacent electric substation. This access shall be used by the construction traffic movements as well as the future on-site operational demands.




Figure 4-1 – Detailed site layout showing access point via existing access to electric sub-station

All heavy and over-dimensional vehicles associated with the development shall travel to and from the site via the Bruxner Highway. Due to safety concerns at the intersection of the Bruxner Highway and the New England Highway trucks heading south from the site shall travel via Ashford and Inverell. The proposed heavy vehicle routes are shown in Figure 4-2 below.






Figure 4-2 – Heavy and Over-Dimensional Vehicle Route








Figure 4-3 - Detail for connection between New England Highway and Bruxner Highway using Sunnyside Platform Road

The application is required to obtain relevant permits under Heavy Vehicle National Law (NSW) for the use of overdimensional vehicles on the road network.

4.3 Traffic Diversions

There are no diversions or road closures associated with the construction or ongoing operation of the Bonshaw Solar Farm.

4.3.1 Proposed Speed Zone

No reduced speed zones shall be implemented during construction of the solar farm.

4.4 Traffic Impacts

The potential impacts of construction and operational traffic on the surrounding road network have been assessed and documented and included the cumulative impacts of other State Significant Projects within the general locality.

As there are no road closures or diversions required during the various stages of the project, there shall be only minor delays associated with the proposed on-site construction works on the Bruxner Highway.



There will be no public vehicle access within the work site during the construction works, with a fence provided at the commencement of the project along the entire site boundary. This fence will remain once the project is constructed for security purposes with a locked gate to be provided at the site access off the Bruxner Highway.

All vehicles are to be loaded and unloaded within the site only and will enter and exit in a forward direction.

Deliveries are to be scheduled to avoid platooning of heavy vehicles travelling together when accessing or departing the site. Two trucks travelling together should leave a suitable gap to enable other drivers to overtake as required.

4.4.1 Construction Parking Demands

Parking of vehicles will be accommodated as required within the site. Management policies shall be implemented to discourage construction workers living in Tenterfield and Glen Innes as well as potentially Goondiwindi driving to the site (unless required for the transportation of specialist tools and equipment).

No vehicles shall be permitted to park on the Bruxner Highway or other local roads.

4.4.2 Public Transport and School Buses

There is school bus that runs along the Bruxner Highway, with one bus in the morning and one bus in the afternoon. The majority of the nominated heavy vehicle routes does not form part of the school bus route with the only interactions being along the Bruxner Highway and the New England Highway to Tenterfield.

Whilst there may be some deliveries occurring in the morning during the school bus operation, very limited deliveries are expected in the afternoon period. All staff will be on site prior to the morning bus run and will depart site after the afternoon bus run. As such it is considered that there is very limited interaction with the school bus.

On the regional and state road network all school zones will be delineated in accordance with Transport for NSW Guidelines with reduced speed limits in accordance with normal NSW road rules. All drivers associated with the project construction work will adhere to the road rules as applicable and will be advised of the school bus operation on the Bruxner Highway.

There will be no impact upon public transport services with no diversions required. There are no bus stops impacted upon by the proposal.

4.4.3 Pedestrians and Cyclists

Given the rural location, demands for pedestrians and cyclists are very low. No pedestrian or cyclist diversions are required for the project work.

4.4.4 Emergency Services

There will be minimal impact for emergency vehicles with no diversions required.

4.4.5 Local Residents

There are no residential dwellings in the immediate locality of the site access that will be impacted upon by the project and construction work. There are a number of residences along the heavy and light vehicle access routes and these residents will be notified in writing of the construction works as necessary including any works which may impact on the local road network.

4.4.6 Other Developments

There will be minimal impact upon any other development within the locality of the site.

4.4.7 Adjoining Council Areas

There will be minimal impact upon adjoining Council areas. Traffic routes in and out of the locality will be along the arterial road network which will experience minimal impacts due to the works.





5 Environmental Conditions

Noise has been considered in the approved working times.

Construction vehicle movement on internal roads could lead to dust generation. A water truck will be used for dust suppression to minimise the production of dust, with the amount of water spreading adjusted accordingly to reflect the conditions. Additionally, any significant deposits of dirt and other construction materials will be promptly removed from public roadways.

Vehicles departing the site are to be in a clean condition to ensure dirt is not tracked into the public road network. A suitable wash bay shall be provided adjacent to the access point for the cleaning of vehicles wheels as required. The access to the sub-station provides a length of seal to prevent the tracking of dirt onto the Bruxner Highway.

The rural location of the site presents a number of environmental hazards such as fog or wildlife along the haulage route. Drivers accessing the site are to drive to suit the road conditions including slowing down in fog or adverse weather conditions. Incidents involving wildlife are to be reported to the site and relevant authorities as appropriate.

6 Incident Procedures

In the case of an incident on site, emergency services will be notified and given priority access to the site as required.

7 Complaints

Any complaints received from the community in relation to the construction and operational traffic for Bonshaw Solar Farm shall be directed to the nominated Health Safety Environment and Community Manager for the project. Details shall be confirmed.

8 Review Process

The Construction Traffic Management Plan and relevant Traffic Control Guidance Plans (TCGPs) will be monitored regularly to ensure their effectiveness and applicability and updated accordingly.

9 Compliance Documentation

Refer to Legislative and Jurisdiction compliance requirements, company policies and procedures as appropriate.



10 Traffic Control Guidance Plan

10.1 General

This TCGP has been prepared to meet the requirements of the RMS Traffic Control at Work Sites Manual 2018. The plans cover the access requirements to the site on the Bruxner Highway.

At all times the Roads and Maritime Service's Traffic Control at Work Sites guidelines must be adhered to. Please refer to the RMS guidelines for traffic control matters not listed in this report.

10.2 Existing Traffic Conditions

- 100 km/hr posted speed limit on the Bruxner Highway.
- Daily traffic volumes on the Bruxner Highway are less than 300 vehicles per day.

10.3 Cyclists and Pedestrians

Given the rural location, demands for pedestrians and cyclists are zero.

10.4 General Traffic Control Considerations

The factors that have been considered in preparing the TCGP are:

- Duration of the works and minimising potential disruptions to local traffic.
- Provision of safe access for heavy and over-dimensional vehicles off the Bruxner Highway.
- Existing traffic volumes and road conditions.

The location and nature of the work will NOT require safety barrier to be installed along the Bruxner Highway.

10.5 Traffic Control – Signage and Line Marking

The TCGP provides Work Site definition. Temporary signage required as part of the works are included due to the nature of the passing traffic, access for construction traffic to the site and the location and nature of the works.

All signs shall be placed on the verge adjacent to the road and made secure against wind and shall be covered when not in use and removed outside of working hours. The signs shall be uncovered before any trucks access the site.

A copy of this TCP must always be on site during the construction work.

10.6 Compliance

The traffic impact assessment prepared by Seca Solution demonstrates that the construction and operation will not have an unacceptable impact on the surrounding roads and intersections.

b) MANAGEMENT PLAN DETAILS:

- Statutory Requirements: Prepare in accordance with Transport for NSW and Council, identify measures that are to be implemented to ensure road safety, detail heavy vehicle routes, identify driver code of conduct, minimise road impacts and environmental risks, notification of any upcoming major works to the local community.
- Limits & Performance Measures: No traffic control is required to manage traffic throughout construction works. Refer to the above report for further details and below for indicators.





- iii) Specific Performance Indicators None required except installation of TCGP
- c) DESCRIPTION OF MEASURES USED TO COMPLY WITH STATUTORY REQUIREMENTS:
 - Licenced traffic consultant and traffic controllers engaged to complete the works
 - Traffic control signage and plan to be maintained throughout the works
 - Maintain through access on affected roads
- d) PROGRAM FOR MONITORING & REPORTING To be managed by the Construction Contractor
- e) CONTINGENCY PLAN:

If the traffic control plan does not maintain the safety of road users and construction workers, the TCP is to be modified by a suitably qualified and accredited person and changes recorded within the register.

f) PROGRAM TO IMPROVE PERFORMANCE

The traffic management plan will be regularly reviewed to ensure its suitability for the works being completed.

- g) PROTOCOL FOR MANAGING & REPORTING INCIDENTS & COMPLAINTS
 - i) All incidents and / or non-compliances that may arise will be documented and the HSEC manager notified immediately on the same business day. This will be documented within the site diary and / or raised at weekly disruption meetings.
 - ii) Complaints shall be directed to the nominated HSEC manager and appropriate actions taken to address any concerns raised.
- h) PROTOCOL FOR PERIODIC REVIEW

Periodic review of the plan is captured above.

10.7 Daily Checklist

In accordance with the Roads and Maritime Services of New South Wales 'Traffic Control at Worksites' guidelines, the site foreman / manager should complete a daily traffic control checklist and this checklist should be filed for future reference.

Details on this checklist can be found at https://www.rms.nsw.gov.au/business-industry/partners-suppliers/documents/technical-manuals/traffic-control-at-worksites-manual.pdf





10.8 Contractors Contact Details

Project Manager: **TBC**

Telephone: **TBC**

Mobile: **TBC**

E-mail: **TBC**

10.9 TCP Approval

This TCGP will be submitted to the road authority for review and approval.

Details for lodging this TCGP and the Construction Traffic Management Plan are:

Inverell Shire Council:

Inverell Shire Council Administration Offices, P O Box 138, Inverell NSW 2360

Transport for NSW (formerly RMS):

Transport for NSW, RMS, P O Box 576, Grafton NSW 2460

This Traffic Control Plan has been prepared and reviewed by suitable qualified professionals in accordance with the RMS Traffic Control at Work Sites Manual 2018 edition.

Sean Morgan (PWZTMP 0051749238)


Director

15/



SECAsolution


Appendix A. Site Plan





Appendix B. TCP at site access on the Bruxner Highway

TCP 195







Appendix C. Driver Code of Conduct

This Drivers Code of Conduct has been established to minimise the impact of our construction on the environment and the local community whilst providing our customers with a high quality, reliable and safe service.

All persons traveling to and from the Bonshaw Solar Farm must adhere to the rules detailed in this document.

Please ensure all persons attending the construction site read, understand and sign this document and return it to the nominated site representative.

General	All vehicles / drivers accessing the site must: i) Be registered and hold a valid driver's licence for the class of vehicle being operated.
	ii) Operate the vehicle in a safe and appropriate manner whilst travelling to / from the site or when operating within the site. This
	includes obeying all New South Wales state road rules. Comply with the directions of authorised personnel when operating within the site and obey any relevant signage installed along the internal roads.
	iv) Not use a mobile phone while operating any vehicle.
	v) Must always wear a seatbelt when operating any vehicle.
Times	Construction
	Construction is to be in completed in accordance with the <i>Interim Construction Noise Guideline</i> (DECC 2009) which defined standard construction work hours
	as: Manday to Friday 7am to 6am
	Monday to Friday: 7am to 6pm Catuaday 9am to 1am
	Saturday: 8am to 1pm
	Sunday and NSW Public holidays: No work
	The following construction, upgrading and decommissioning activities may be undertaken outside these hours without the approval of the secretary: • The delivery of materials as requested by the NSW Police Force or other authorities for safety reasons; or
	Emergency work to avoid loss of life, property and / or material harm to the environment.
	Vehicle movements shall be undertaken during standard construction hours (or just before to allow workers to get to site). Oversize vehicles up to 26 metres long may require access to the site after hours however this would be subject to the requirements of Transport for NSW, Inverell Shire Council or NSW Police.
	Normal Operations Daily operations and maintenance by site staff would be undertaken during standard working hours:
	Monday to Friday: 7am to 6pm
	Saturday: 8am to 1pm
	Sunday and NSW Public holidays: No work
	During normal operations, all vehicle movements shall be undertaken during the standard operating hours (or just before to allow workers to get to site).

	There may be a requirement for vehicles to access the site after hours during an emergency however these would be infrequent.
Access	All heavy vehicles must adhere to the designated heavy vehicle routes as nominated in Figure 1 below.
	Drivers must ensure they enter and exit the site in a forward direction and are not permitted to load or unload on the Bruxner Highway.
Vehicle Departure and Arrival	Heavy vehicles departing the site shall have a minimum 5 minute separation to reduce the impacts upon the local road network.
	Always maintain a minimum separation of at least 50 metres between vehicles when travelling within the site.
	Drivers must contact the site supervisor upon arrival and await further instructions or direction before proceeding. Drivers must also report to the site supervisor prior to departure.
	All vehicles must enter and exit the site in a forward direction. Vehicles tyres are to be washed down and in a clean condition upon exiting the site to prevent dirt being tracked onto the public road network
Illegal and Unsafe Truck Parking	No vehicles are permitted to park on the Bruxner Highway except in designated parking bays.
	All parking within the site is to be as directed.
Heavy Vehicle Compression Braking	Compression braking by heavy vehicles is a source of irritation to the community. Brakes must be applied so as not to create excessive noise that could disturb residents along the haulage routes, particularly within the township of Tenterfield.
	Compression braking should only be used if required for safety reasons.
Load Covering and Restraint (as per NHVL and CoR Regulations)	All trucks arriving to or departing the site are required to have an effective cover over their load for the duration of the trip.
regulationey	Drivers must ensure that following loading or unloading that all gates and tailgates are secured and locked before leaving the site.
	Drivers are to ensure that their loads are secured as per the NTC Load Restraint Guide 2018 and that checks are completed on restraint equipment such as tailgates, chains, straps, ratchet dogs, tarps etc on a regular basis to ensure they are functioning correctly.
	Drivers must exit their vehicle to inspect the above every time they are loaded, prior to arrival at the site and following unloading at the site, prior to departure. The security of your load, your life and the life of others relies upon proper load restraint practices.
Mass and Dimension (as per NHVL and CoR Regulations)	Drivers should be aware that:



Catigue and License	All drivers should be aware that:
Fatigue and License Requirements (as per NHVL	All drivers should be aware that:
and CoR Regulations)	a person must not drive a heavy vehicle on a road while impaired by fatigue
	 managing driver fatigue is a shared responsibility by all parties in the chain
	 parties must take all reasonable steps to ensure a person does not drive the heavy vehicle on a road while impaired by fatigue.
	Drivers shall not be permitted to operate a vehicle or plant equipment when impaired by fatigue. If you suspect that you or someone else is experiencing fatigue, please inform your supervisor.
	Operators of heavy vehicles shall be aware of the requirements relating to fatigue as outlined in the Heavy Vehicle National Law. Drivers shall also be aware of their adopted fatigue management scheme (shown below) and ensure that they are operating within its requirements. i) Standard Hours of Operation ii) Basic Fatigue Management (BFM)
	iii) Advanced Fatigue Management (AFM)
	All drivers accessing the site must be registered and hold a valid driver's licence for the class of vehicle being operated
Vehicle Speeds	Drivers shall observe the posted speed limit along the designated transport route and adjust their vehicle speed as required to suit the road environment and prevailing weather conditions. Vehicle speeds must be appropriate to ensure the safe movements of the vehicle with consideration to the vehicle configuration.
	Maximum speeds limits within the project site shall be as follows:
	i) 40 km/hr along formed roads.
	ii) 20 km/hr during foggy / dusty conditions. Headlights must be on. iii) 10 km/hr when passing pedestrians or any plant equipment.
Overtaking	Overtaking shall not be permitted within the site unless the intention to overtake has been communicated to the driver of the leading vehicle and consent to overtake granted.
Breakdowns and Incidents	Heavy Vehicles
	In the case of a breakdown, the vehicle must be towed to the nearest breakdown point as soon as possible. All breakdowns must be reported to the Transport for NSW Transport Management Centre on 131 700 and the vehicle protected in accordance with the Heavy Vehicle Drivers Handbook. The relevant shift manager on site shall also be notified.
	If a breakdown occurs on-site please remain inside your vehicle, notify the shift manager of your location and await further instruction.
	If you are involved in an accident, please notify the shift manager immediately and contact emergency services if required. Light Vehicles
	In the case of a breakdown, ensure that the vehicle is secure, notify the site manager of your location and await further instruction.
	If you are involved in an accident, please notify the shift manager immediately and contact emergency services if required.
Signs	Comply with the directions of authorised personnel when operating within the site and obey any relevant signage installed along the internal roads.



Personal Protective	The mandatory Protective Clothing requirement on this site is TO BE
Equipment (PPE)	CONFIRMED
1 1 (/	Hard Hat
	Steel cap work shoes
	Safety Glasses
	High-Vis clothing
	Ear Protection (as required)
	Gloves (as required)
Mobile Phone Usage	A mobile phone is not to be used while operating any vehicle.
	The use of a mobile phone while operating machinery or undertaking site activities is STRICTLY PROHIBITED. Driving, operating or undertaking high risk activities must be temporarily stopped to take a call.
	When a call or text message is received while driving and a hands-free kit is not available or the hands free kit is not voice activated, the driver MUST pull to the side of the road when safe to do so prior to taking the call or checking the message.
	Where a conversation is complex, technical or requires notes to be taken is mandatory to pullover when safe to do so to continue the conversation.
	Any person contravening this Policy will be subject to the Company's disciplinary procedure which may include termination of Employment/Contractor contract.
Alcohol and Drug Policy	All persons are to be in a "fit for work" state. This means must not be affected by alcohol or other drugs, (including prescription medication if inhibiting ability to operate plant and equipment safely) whilst at work. It is prohibited for any person to possess, use, sell or work under the influence of Alcohol or other Drugs.
Contacts	Transport for NSW Transport Management Centre 131 700 Site Supervisor
	Emergency 000






Figure 1 - Heavy and Over-Dimensional Vehicle Route






Figure 2 - Transport route to/from the site for ALL 19 m semi-trailers or greater (detail for northbound via New England Highway to use Sunnyside Platform Road)



Driver Declaration

I, the undersigned, hereby agree to abide by this Driver Code of Conduct for the transport of equipment, product or personnel to / from the construction site to the Bonshaw solar farm located off the Bruxner Highway, Bonshaw, NSW. I have read and understand the requirements outlined in the attached document and will, to the best of my ability, comply and assist with their implementation, requirements or ongoing administration.

The subject document to which this declaration relates is included as part of this overall document and signing of this declaration confirms that the signee has read and understood their requirements as outlined throughout.

Driver Details

Full Name	
Organisation	
Signature	
Date	

Representative of:

Full Name	
Signature	
Date	

Disciplinary action will be taken against drivers who do not adhere to this Code of Conduct.

ERM		Bonshaw Solar Farm (SSD 9438) Clarification Letter
		Page A1
ATTACHMENT B	SITE ACCESS ROAD UPGRAD HERITAGE ASSESSMENT	DE – BIODIVERSITY &

1. SITE ACCESS ROAD UPGRADE

1.1 Introduction

Access to the site will be via the existing access provided to the electric substation adjacent to the subject site. This access was constructed to allow for the construction activities associated with the Dumaresq substation, and is now used for maintenance purposes only for the substation.

This access will be upgraded to provide a left turn deceleration lane and shoulder widening to provide an auxiliary right turn treatment, both in accordance with Austroads Guidelines (refer to **Figure 1-1** below)

Environmental impacts of these road upgrades are considered herein, namely associated with biodiversity and heritage matters. However, it is highlighted that both the construction of the Bruxner Hwy and the use of the Travelling Stock Reserve (TSR) immediately south of the Bruxner Hwy have added to the highly disturbed nature of this area.

1.2 Biodiversity Impact


The proposed road upgrades will occur within the existing road corridor, on the shoulders of the Bruxner Hwy. This area has been cleared and is now dominated by introduced grass species. There are no anticipated impacts to biodiversity matters.

1.3 Heritage Impact

The original heritage surveys have identified Aboriginal artefacts in the northern part of the development footprint, including BSF1, a significant artefact scatter with potential archaeological deposit. This has been mapped as extending to the Bruxner Hwy corridor. However, this area is located further to the east of the site access point and does not extend to the road upgrade area.

The road shoulder is highly disturbed from the original Bruxner Hwy construction and is unlikely to contain Aboriginal heritage matters.

In the event that an Aboriginal artefact is uncovered during construction, an unexpected finds protocol will be implemented for the upgrade works, forming part of the Construction Environmental Management Plan (CEMP).



ERM	Bonshaw Solar Farm (SSD 94: Response to RFI – May 2020 Page 17 of 8
	I
ATTACHMENT C	UPDATED BIODIVERSITY ASSESSMENT REPORT





GAIA Bonshaw Solar EIS

Revised Biodiversity Development Assessment Report

18 May 2020

Project No.: 0470861



Document details	
Document title	GAIA Bonshaw Solar EIS
Document subtitle	Revised Biodiversity Development Assessment Report
Project No.	0470861
Date	18 May 2020
Version	4.0
Author	Matt Jenkins North Coast Ecology BAAS 18029
Client Name	GAIA Australia

Document history

				ERM approva	I to issue	
Version	Revision	Author	Reviewed by	Name	Date	Comments
Draft	01	Adriana Corona Mothe	Joanne Woodhouse	Paul Douglass	1 July 2019	Draft for client review
Final	01	Adriana Corona Mothe	Joanne Woodhouse	Paul Douglass	26 July 2019	Final to support EIS submission
Final	02	Matt Jenkins	Joanne Woodhouse	David Dique	20 February 2020	Revision based on BCD review and comment
Final	03	Matt Jenkins	Joanne Woodhouse	Henrietta Jukes	20 March 2020	Addition of overhead transmission corridor
Final	04	Matt Jenkins	Joanne Woodhouse	Henrietta Jukes	15 May 2020	Updated based on DPIE RFI

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

Signature Page

18 May 2020

GAIA Bonshaw Solar EIS

Revised Biodiversity Development Assessment Report

Matt Jenkins North Coast Ecology

Michael Rookwood Project Manager

Joanne Woodhouse Principal Ecologist

M.Woodhouse.

Henrietta Jukes Partner

Environmental Resources Management Australia Pty Ltd Level 1, 45 Watt Street Newcastle NSW 2300

PO Box 803

Newcastle NSW 2300

© Copyright 2020 by ERM Worldwide Group Ltd and / or its affiliates ("ERM"). All rights reserved. No part of this work may be reproduced or transmitted in any form, or by any means, without the prior written permission of ERM

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

CONTENTS

EXE	CUTIV	E SUMMA	ARY	I		
STA	GE 1: I	BIODIVER	RSITY ASSESSMENT	1		
1.	INTR	INTRODUCTION				
	1.1 1.2 1.3 1.4 1.5	Assessr The Pro Site Des Consult	ment Requirements			
2.	LEGI	SLATIVE	CONTEXT	15		
3.	LANI	DSCAPE I	FEATURES	18		
4.	NATI	NATIVE VEGETATION				
	4.1	Vegetation Assessment Methodology				
		4.1.1 4.1.2 4.1.3 4.1.4	Review of Existing Information	22 23 24		
	4.2		dentified in the Subject Land			
		4.2.1	Cleared and Disturbed Land			
	4.3	•	tion Integrity Assessment			
		4.3.1 4.3.2	Vegetation Zones and Survey Effort			
5.	THRI	EATENED	SPECIES			
	5.1	Fauna H	Habitat Assessment	44		
		5.1.1 5.1.2	Important Habitat Mapping Onsite Habitat Assessment	44		
	5.2	Rivers,	Streams and Wetlands			
		5.2.1 5.2.2	Creek LinesFarm Dams			
	5.3		tem Credit Species			
	5.4 5.5		S Credit Species			
	5.6		nal Species Consideredate Species Surveyedate Species Surveyed			
	5.7		d Survey Results			
		5.7.1 5.7.2 5.7.3 5.7.4 5.7.5 5.7.6 5.7.7	Flora Species Recorded Fauna Species Recorded Grey-crowned Babbler Microchiropteran Bats Owls Cicadabird Threatened Aquatic Species			
	5.8	Limitatio	ons To Data	75		
STA	GE 2: I	MPACT A	ASSESSMENT (BIODIVERSITY VALUES)	81		
6.	IMPA	CT ASSE	ESSMENT (BIODIVERSITY VALUES)	82		
	6.1 6.2 6.3	Avoiding	g and Minimising Impacts On Native Vegetation and Habitat g and Minimising Prescribed Biodiversity Impactsthat Cannot be Avoided	83		

		6.3.1 6.3.2 6.3.3	Direct Impacts	88	
	6.4		Prescribed Impacts		
7.	• • •	-	NT AND MITIGATION MEASURES		
8.	SERIO	US AN	D IRREVERSIBLE IMPACTS (SAII)	99	
	8.1		ial Serious and Irreversible Impact Entities		
		8.1.1	Threatened Ecological Communities		
		8.1.2	Threatened Species	100	
		8.1.3	Additional Potential Entities		
	8.2 Assessment of SAII				
9.	REQUI	REME	NT TO OFFSET	101	
	9.1	Impact	s Requiring an Offset	101	
		9.1.1	Impacts on Native Vegetation – Ecosystem Credits	101	
		9.1.2	Impacts on Threatened Species – Species Credits		
		9.1.3	Offsets Required Under the EPBC Act		
	•		s Not Requiring an Offsetary of Offset Credits Required		
	9.4		ersity Offset Framework		
10.	CONCI	_USIO	NS	105	
11.	REFER	REFERENCES10			
APPE	ENDIX A	. E	BOSET REPORT		
APPENDIX B NSW BIONET AT			ISW BIONET ATLAS RECORDS		
APPE	ENDIX C	ı.	MATTERS OF NATIONAL SIGNIFICANCE ASSESSMENT – PMST REPO	RT	
APPE	ENDIX D	L	IKELIHOOD OF OCCURRENCE AND RISK ANALYSIS		
APPE	ENDIX E	Т	EST OF SIGNIFICANCE AND SIGNIFICANT IMPACT ASSESSMENT		
APPE	NDIX F	S	SURVEY METHODS AND EFFORT		
APPE	ENDIX G	Е	BAM PLOTS DATA AND PHOTOGRAPHS		
APPE	ENDIX H	F	LORA AND FAUNA SPECIES LISTS		
APPE	ENDIX I	N	MICROCHIROPTERAN BAT CALL REPORT		
APPE	ENDIX J	Е	BAM REPORTS – VEGETATION INTEGRITY ASSESSMENT		
APPE	ENDIX K	E	SAM CALCULATOR CREDIT REPORT		
APPE	ENDIX L	P	PERSONNEL		

Client: GAIA Australia

18 May 2020

List of Tables

Table 1.1	SEARs requirements and how they have been addressed.	2
Table 1.2	Government agency requirements	3
Table 1.3	Key Features of the Subject Land	9
Table 1.4	Summary of Consultation with BCD during preparation of the BDAR	10
Table 2.1	Applicable Legislation, Plan and Guidelines	15
Table 3.1	Summary of Landscape Features	18
Table 4.1	Mapped Vegetation Communities (SVTM VIS 4681) within the Buffer Area and Sul	oject
Land (not gro	ound-truthed)	22
Table 4.2	Plant Community Types Occurring within the Subject Land	26
Table 4.3	PCT 594: Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brig	alow
Belt South B	ioregion and Nandewar Bioregion	26
Table 4.4	PCT 596: Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark	
shrubby woo	dland mainly in the northern Nandewar Bioregion	29
Table 4.5	PCT 516: Grey Box grassy woodland or open forest of the Nandewar Bioregion an	ıd
New England	d Tableland Bioregion	31
Table 4.6	Summary of Vegetation Integrity Scores	40
Table 5.1	Additional Habitat Features Considered	44
Table 5.2	Ecosystem Credit Species (BAM Calculator)	52
Table 5.3	Species Credit Species	55
Table 5.4	Candidate Species Surveyed	62
Table 5.5	Summary of Fauna Species Recorded	66
Table 5.6	Habitat Summary for Microchiropteran Bat Species Credit Species	69
Table 6.1	Avoidance and Minimisation of Impacts	83
Table 6.2	Direct Impacts to Biodiversity during the Construction and Operational Phases	87
Table 6.3	Current and Future Vegetation Integrity Scores for each Vegetation Zone	88
Table 6.4	Potential Indirect Impacts Associated with the Proposed Development	89
Table 6.5	Identification of Prescribed Impacts associated with the Proposed Development	91
Table 6.6	Assessment of the Project in relation to EPBC Act	93
Table 7.1	Recommended Mitigation Measures for Residual Impacts	94
Table 8.1	Potential SAII within the Development Site	99
Table 9.1	Summary of Ecosystem Credits	101
Table 9.2	Summary of Impacts on Vegetation Not Requiring Offsets	103
Table B.1	BioNet Atlas results	1
List of Figur	res	
Figure 1.1	Site Map	7
Figure 1.2	Indicative Development Layout	
Figure 3.1	Areas listed as high biodiversity value	
Figure 3.2	Location Map	
Figure 4.1	State Vegetation Type Community (VIS 4681)	41
Figure 4.2	Vegetation Zones within the Development Site	
Figure 4.3	PCTs and BAM Plot locations	
Figure 5.1a	Targeted Survey Locations	
Figure 5.2	Threatened Species Recorded on the Subject Land	
Figure 5.3	Species Polygons for Relevant Species Credit Species	
Figure 5.4	Threatened Aquatic Species	

Acronyms and Abbreviations

Name	Description		
AoS	Assessment of Significance (under the EPBC Act)		
AOBV	Areas of Outstanding Biodiversity Value		
APZ	Asset Protection Zone		
Assessment Area Assessment Area refers to the 1,500m buffer area surrounding the Subject La other areas requiring consideration as part of the biodiversity values assessm Stage 1 of the BAM			
BAM	Biodiversity Assessment Method		
BC Act	NSW Biodiversity Conservation Act 2016		
BCD	Biodiversity Conservation Division of the Environment, Energy and Science Group in the NSW Department of Planning, Industry and Environment		
BDAR	Biodiversity Development Assessment Report		
BIA	Biodiversity Impact Assessment		
BOSET Biodiversity Offset Scheme Entry Threshold			
BSA	Biodiversity Stewardship Agreement		
Buffer area	All land within 1500 m of the Subject Land.		
BV Map	Biodiversity Values Map		
Candidate species	Threatened species requiring assessment other than 'ecosystem credit species' and 'species credit species'		
Development Site	The area of land that is subject to either temporary or permanent impact. At the proposed Bonshaw solar farm, the development site includes the land proposed to insta the solar arrays, perimeter fence, access roads, transmission line and ancillary construction areas. The development site is approximately 149.24 ha.		
Ecosystem credits	Is a measurement of the value of threatened ecological communities, threatened specie habitat for species that can be reliably predicted to occur with a PCT and PCTs generally. Ecosystem credits measure the loss in biodiversity values at a development site.		
Ecosystem Credit Species	Are threatened species identified as such in the Threatened Biodiversity Data Collection and whose selection was based on the likelihood of occurrence of the species or elements of the species' habitat can be predicted by vegetation surrogates and landscape features, or for which targeted survey has a low probability of detection.		
EIS	Environmental Impact Statement		
EP&A Act	Environmental Planning and Assessment Act 1979		
EPBC Act	Commonwealth Environment Protection and Biodiversity Conservation Act 1999		
ERM	Environmental Resources Australia Pty Ltd		
FM Act	NSW Fisheries Management Act 1994		
GAIA	GAIA Australia		
На	Hectares		
HBT	Hollow Bearing Tree		
IBRA Bioregion	A bioregion identified under the Interim Biogeographic Regionalisation for Australia (IBRA) system, which divides Australia into bioregions on the basis of their dominant landscape-scale attributes		

Client: GAIA Australia

EXECUTIVE SUMMARY

Environmental Resources Australia Pty Ltd (ERM) was commissioned by GAIA Australia (GAIA) to prepare a Biodiversity Development Assessment Report (BDAR) to support an Environmental Impact Statement (EIS) for a proposed solar farm located within Lot 2 DP1039185, Bonshaw NSW.

The proposed development will involve the construction, operation and maintenance of a solar PV generation facility and associated infrastructure with a capacity of up to 200 MW and associated infrastructure, including a Lithium-ion Energy Storage System, supplying electricity to the national electricity grid. The Development Site includes all supporting infrastructure and site access points.

This BDAR was prepared in accordance with the requirements of the NSW *Biodiversity Conservation Act 2016*, including the Biodiversity Assessment Method (BAM) and Biodiversity Offsets Scheme (BOS). The BDAR also considers relevant matters associated with the Commonwealth *Environment Protection and Biodiversity Conservation Act 1999* and the addresses the Secretary's Environmental Assessment Requirements (SEARs) provided for the Project.

Flora and fauna surveys were undertaken at the Development Site across four survey periods (11-13 September 2018 (spring), 11-14 December 2018 (summer), 25-28 March 2019 (autumn) and 21-22 January 2020 (summer)). Flora and habitat surveys included vegetation mapping along random meander transects and over 100 vegetation community observation points and 44 BAM plots. Targeted fauna surveys included amphibian call payback (three surveys), bird surveys (eight surveys), reptile surveys (eight surveys), arboreal/tree hollow dependent fauna surveys (eight camera trap over 28 trap nights), microchiropteran bat call recording (eight SongMeters totalling 28 trap nights) and fauna habitat observations (hollow bearing trees, ant/termite mounds, rocky areas, bird nests and creek lines).

The Subject Land is comprised of highly disturbed land whose current and historical land uses include livestock grazing and cropping. Several creeks (first, second and third order), representing tributaries of the Dumaresq River, occur within the Project Boundary along with eight farm dams. The majority of the Development Site consists of cleared grazing land (72%), with the remainder comprising disturbed native woodland vegetation (28%).

A total of 143 flora species in 47 families were recorded within the assessment area. This included a total of 111 native (78%) and 32 exotic (22%) species. No threatened flora species were recorded on the Subject Land. Vegetation occurring on the Subject Land is represented by the following four Plant Community Types (PCTs):

- PCT 594 Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion. Two vegetation zones of this PCT are present: moderate and low condition;
- PCT 596 Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion. Four vegetation zones of this PCT are present: moderate condition, low condition, very low condition and derived grassland;
- PCT 544 Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest/woodland of the Nandewar Bioregion and New England Tableland Bioregion. One vegetation zone of this PCT is present: low condition; and
- PCT 516 Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion. Three vegetation zones of this PCT are present: low condition, very low condition and derived grassland.

PCT 544 is considered to represent the following Threatened Ecological Community, which is listed as Endangered under the NSW *Biodiversity Conservation Act 2016:*

White Box Yellow Box Blakely's Red Gum Woodland

The occurrence of this TEC on the Subject Land is not considered to satisfy the minimum condition criteria for the Commonwealth listing of this TEC. This determination has been made on the basis of very small patch sizes occurring on the Subject Land, low diversity of native understorey species and absence of "important species" as described in the Threatened Species Scientific Committee listing advice and National Recovery Plan for the TEC.

A total of 75 fauna species were recorded within the Project Boundary, including thirteen threatened species. Threatened species included eight vulnerable microchiropteran bats (Little Pied Bat, Eastern Bent-wing Bat, Corben's Long-eared Bat, Yellow-bellied Sheath-tailed Bat, Eastern Cave Bat, Eastern False Pipistrelle, Greater Broad-nosed Bat and Hoary Wattled Bat), one endangered microchiropteran bat (Bristle-faced Free-tailed bat), three vulnerable birds (Grey-crowned Babbler, Brown Treecreeper and Speckled Warbler) and one migratory bird (Cicadabird). Out of the eight Vulnerable microchiropteran bats, six were recorded as "definite" calls and two (Greater Broad-nosed Bat and Hoary Wattled Bat) were recorded as "possible" calls by a microbat call identification expert. Brown Treecreeper and Speckled Warbler were recorded within the Project Boundary but not directly on the Subject Land.

An assessment of the impacts of the project on Matters of National Environmental Significance (MNES) within the Development Site was undertaken. A likelihood of occurrence and risk assessment was undertaken for each entity identified as likely to occur within the Project's locality. It was concluded that all TECs and threatened species as identified in the PMST would have low residual risk as a result of the Project. Therefore, assessments of significance and referral for assessment of the project to the Commonwealth Minister for the Environment is not required.

GAIA has undertaken significant steps to avoid, minimise and mitigate impacts to biodiversity. As part of the project refinement process, ERM provided advice to GAIA on areas which were of the highest priority for avoidance. This led to third and second order creeks and other conservation significant areas to be avoided.

The Project will result in the following direct impacts to flora and fauna:

- Removal of 40.97 ha of disturbed native vegetation pertaining to four PCTs and representing potential habitat for a range of threatened fauna species;
- Impacts to 116.83 ha of heavily disturbed grassland with vegetation integrity score <17;</p>
- Removal of 34 hollow-bearing trees representing potential roosting and/or breeding habitat for threatened microchiropteran bats and other hollow-dependent species including threatened forest owls:
- Impacts to suitable foraging habitat for the Eastern Cave Bat in less disturbed areas of PCTs 516, 544, 594 and 516;
- Impacts to suitable breeding and foraging habitat components for the Bristle-faced Free-tailed Bat; and
- Removal of nests, nest trees and foraging habitat for the Grey-crowned Babbler.

Potential indirect impacts associated with the proposed development have been identified, including edge effects, fragmentation, and sedimentation and pollutant run-off. With consideration of the proposed management and mitigation measures, it is expected that potential indirect impacts will be reduced to acceptable levels and any residual impacts will be negligible.

Direct impacts to native vegetation and fauna habitat requiring offsets include:

- Impacts on PCT 516 Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion, requiring 65 ecosystem credits;
- Impacts on PCT 544 Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest/woodland of the Nandewar Bioregion and New England Tableland Bioregion, requiring 15 ecosystem credits;

- Impacts on PCT 594 Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion ad Nandewar Bioregion, requiring 249 ecosystems credits;
- Impacts on PCT 596 Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion, requiring 269 ecosystem credits;
- Impacts on Bristle-faced Free-tailed Bat habitat, requiring 1,180 species credits;
- Impacts on Eastern Cave Bat foraging habitat, requiring 1,030 species credits;
- Impacts on potential (assumed) breeding habitat for the Masked Owl, requiring 680 species credits; and
- Impacts on potential (assumed) breeding habitat for the Barking Owl, requiring 680 species credits.

This assessment has been completed in accordance with the BAM (OEH 2017a, 2018a) on behalf of GAIA. Overall the Subject Land is considered to be of moderate biodiversity value with impacts related to direct removal of native vegetation and fauna habitat and displacement of resident fauna. The proposed Bonshaw Solar Farm development in its current footprint requires a total offset of 598 ecosystem credits and 3,570 species credits. Retirement of these credits will be carried out in accordance with the *Biodiversity Conservation Regulation 2017* and the NSW Biodiversity Offsets Scheme.

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page iii

STAGE 1: BIODIVERSITY ASSESSMENT

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page 1

1. INTRODUCTION

GAIA Australia (GAIA) is seeking to develop a large scale solar photovoltaic (PV) generation facility and associated infrastructure with the capacity of 200 megawatts (MW) situated near Bonshaw in the Inverell Shire Council (ISC) of New South Wales (NSW) (the Project).

The Project is classified as State Significant Development (SSD) in accordance with Clause 20 Schedule 1 of the State Environmental Planning Policy (State and Regional Development) 2011 (SEPP) and will be assessed under Part 4 of the Environmental Planning and Assessment Act 1979 (EP&A Act). This Biodiversity Development Assessment Report (BDAR) assesses the impacts of the Project according to the NSW Biodiversity Assessment Methodology (BAM) as required by the Secretary's Environmental Assessment Requirements (SEARs) for the proposal.

The aim of this BDAR is to undertake biodiversity and impact assessment of ecological values of the Subject Land in accordance with the BC Act. This BDAR also addresses the assessment requirements of the Commonwealth Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act).

1.1 Assessment Requirements

On 16 August 2018, the Department of Planning and Environment (DPE) provided Secretary's Environmental Assessment Requirements (SEARs) for the Bonshaw Solar Farm (the Project).

A copy of the SEARs is attached to the EIS as Appendix A. The assessment requirements that specifically relate to biodiversity, are listed in Table 1.1.

Table 1.1 SEARs requirements and how they have been addressed.

Requirement	Section Addressed
An assessment of the biodiversity values and the likely biodiversity impacts of the project in accordance with Section 7.9 of the <i>Biodiversity Conservation Act 2016</i> (NSW) the Biodiversity Assessment Method (BAM) and documented in a biodiversity development assessment report (BDAR), unless OEH (now BCD) and DPE determine that the proposed development is not likely to have any significant impacts on biodiversity values.	This BDAR has been prepared in accordance with the <i>Biodiversity</i> Conservation Act 2016, Biodiversity Conservation Regulation 2017 and the Biodiversity Assessment Method (BAM).
The BDAR must document the application of the avoid, minimise and offset framework including assessing all direct, indirect and prescribed impacts in accordance with the BAM.	Stage 2 of this BDAR, Chapter 6 to 8.
An assessment of the likely impacts on listed aquatic threatened species, populations or ecological communities, scheduled under the <i>Fisheries Management Act 1994</i> , and a description of the measures to minimise and rehabilitate impacts.	Section 5.2.

To inform preparation of the SEARs, DPE invited other government agencies to recommend matters to be addressed in the EIS. These matters were taken into account by the Secretary for DPE when preparing the SEARs. A copy of the former NSW Office of Environment and Heritage (OEH) advice to DPE was attached to the SEARs and matters relevant to the BDAR are listed in Table 1.2.

Table 1.2 Government agency requirements

Requirement

Section Addressed

DPI Fisheries

Aquatic Ecological Assessment

The aquatic ecological environmental assessment should include the following information;

- A recent aerial photograph (preferably colour) of the locality (or reproduction of such a photograph) should be provided.
- Area which may be affected either by the development or activity should be identified and shown on an appropriately scaled map (and aerial photographs).
- Waterways within the area of development are to be identified.
- The extent of aquatic habitat removal and riparian vegetation removal or modification which may result from the proposed development,
- Details of the location and design of the waterway crossings or underground cabling through waterways.
- Details of the methodology (e.g. trenching, boring) for any underground cabling passing through waterways.

Aquatic habitat is described in Section 5.7.7. Waterways are shown on the aerial photo in Figure 3.2, noting that they were dry (or very low water levels) during all survey periods due to extended drought conditions.

The second and third order watercourses will be avoided through detailed design (including 20-30m riparian buffer zones measured from the high bank of the streams).

Any waterway crossings required as a result of the Project will be designed in accordance with the Policy and Guidelines for Fish Habitat Conservation and Management and the Policy and Guidelines for Fish Friendly Waterway Crossings.

Construction methodology is further detailed in the EIS.

Waterway Crossings

The construction of permanent or temporary access tracks or underground cables through Little Oaky Creek, Little Limestone Creek and unnamed creeks running into the Beardy River should be in accordance with DPI Fisheries Guideline document: Policy and Guidelines for Fish Habitat Conservation and Management (Update 2013).

The activities at the
Development Site will not
require any access track or
cables through Little Oaky
Creek, Little Limestone Creek or
tributaries to Beardy River.

Any waterway crossings required as a result of the Project will be designed in accordance with the Policy and Guidelines for Fish Habitat Conservation and Management and the Policy and Guidelines for Fish Friendly Waterway Crossings.

Section Addressed Requirement **Threatened Species, Populations And Ecological Communities** A description of the aquatic habitat including consideration The proposal should include a threatened aquatic species assessment (as of the Purple Spotted Gudgeon per part 7A Fisheries Management Act 1994) to address whether there are is provided in Section 5.2 and likely to be any significant impacts on listed threatened species, an Assessment of Significance populations or ecological communities listed under the Fisheries is provided in Appendix E. Management Act 1994. It should be specifically noted that the proposal is located within an area considered habitat of the threatened species Purple Spotted Gudgeon (Mogurnda adspersa). This species is known or expected to occur in Little Oaky Creek and a number of nearby creeks. Threatened fish species mapping distributions are available at: http://www.dpi.nsw.gov.au/fishing/species-protection/threatenedspeciesdistributions-in-nsw **Biodiversity Conservation Division (BCD)** Biodiversity impacts related to the proposed development are to be This BDAR has been prepared assessed in accordance with Section 7.9 of the Biodiversity in accordance with the Conservation Act 2017, the Biodiversity Assessment Method and Biodiversity Conservation Act documented in a Biodiversity Development Assessment Report 2016, Biodiversity Conservation (BDAR). The BDAR must include information in the form detailed in Regulation 2017 and the the Biodiversity Conservation Act 2016 (s.6.12), Biodiversity **Biodiversity Assessment Method** Conservation Regulation 2017 (s6.8) and Biodiversity Assessment (BAM). Method, unless OEH (now BCD) and DPE determine that the proposed development is not likely to have any significant impacts on biodiversity values. The BDAR must document the application of the avoid, minimise and Chapter 7 offset framework including assessing all direct, indirect and prescribed impacts in accordance with the Biodiversity Assessment Method. The BDAR must include details of the measures proposed to address Chapter 9 the offset obligations as follows: The total number and classes of biodiversity credits required to be retired for the development/project; The number and classes of like-for0like biodiversity credits proposed to be retired; The number and classes of biodiversity credits proposed to be retired in accordance with the variation rules; Any proposal to fund a biodiversity conservation action; Any proposal to conduct ecological rehabilitation (if a mining project); Any proposal to make a payment to the Biodiversity Conservation Fund. If seeking approval to use the variation rules, the BDAR must contain details of the reasonable steps that have been taken to obtain requisite like-for-like biodiversity credits. The BDAR must be submitted with all spatial data associated with the All spatial data (ArcGIS format) survey and assessment as per Appendix A of the BAM. will be provided to BCD in electronic format.

Client: GAIA Australia

	Requirement	Section Addressed
5.	The BDAR must be prepared by a person accredited in accordance with the Accreditation Scheme for the Application of the Biodiversity Assessment Method Order 2017 under s.6.10 of the <i>Biodiversity Conservation Act 2016</i> .	Version 1 of this BDAR was co- prepared by Dr Adriana Corona Mothe (BAM Accreditation BAAS18113).
		Version 2 of this BDAR has prepared by Matt Jenkins (BAM Accreditation BAAS18029) by amending Version 1 where required to addressed comments from BCD.

Following public exhibition of Version 1 of this BDAR, dated 26 July 2019, additional comments were received from the Biodiversity Conservation Division (BCD) of the Environment, Energy and Science Group in the NSW Department of Planning, Industry and Environment (formerly part of the Office of Environment and Heritage). This amended BDAR addresses comments and recommendations provided by the BCD in correspondence dated 2 December 2019 (refer Section 1.4 for summary of consultation and response).

1.2 The Project

The Project (Proposed Development) will involve the construction, operation and maintenance of a solar PV generation facility and associated infrastructure with a capacity of up to 500 MW, supplying electricity to the national electricity grid. The Project is located approximately 16 kilometres (km) south of Bonshaw and 66 km north of Inverell and is wholly contained within the Inverell Local Government Area (LGA) (Figure 1.1). The Project would connect directly to the 330 kilovolt (kV) Dumaresq Substation located to the immediate west of the Project boundary.


The Project incorporates arrays of PV modules (commonly referred to as "solar panels"), transmission infrastructure and substations to enable connection into the existing electricity transmission network (Figure 1.2). The exact method and point of connection is being developed with TransGrid in parallel with this planning application and the detailed infrastructure layout developed during detailed design will confirm the generating capacity of the Bonshaw Solar Farm.


The key elements of the project include the construction and operation of:

- A network of PV modules in a fixed tilt or single axis tracking arrangement;
- Associated BESS(s) / battery storage;
- A switch yard to be connected to the 330 kV TransGrid Dumaresq Substation, on the boundary of the Project Area;
- Underground or overhead cabling for connection between arrays and inverters and transformers;
- Operations and maintenance (O&M) infrastructure, including O&M buildings including a control room, meeting facilities, a temperature controlled spare parts storage facility, supervisory control and data acquisition (SCADA) facilities, a workshop and associated infrastructure (e.g. kitchen, toilets and other facilities) car parking facilities;
- Access point to the site via the Bruxner Highway;
- A new internal road network to enable access from surrounding local roads to the array areas during construction and operations including internal access tracks, creek crossing & perimeter security fencing; and
- Temporary facilities during construction.

A full description of the Project is provided in the EIS and the final layout of the Project will be dependent on detailed design, availability and commercial considerations at the time of construction, however in terms of assessing biodiversity impacts, the following project features were considered:

- Four riparian crossings (indicative location on third and second order streams shown in Figure 1.2);
- Clearing of vegetation to be restricted to the identified 149.24 ha development footprint, noting
 that grazing beneath the solar panels will continue during operation of the solar farm (the height
 of the PV panels above natural ground is approximately 1.4 to 4.2 m based on tracker option to
 be used);
- Excavation of trenches and the laying of power and instrumentation cables;
- Main access point to the site via the Bruxner Highway;
- An internal private road network (up to a combined total length of approximately 13.7 km)
 connecting the arrays and other proposed infrastructure to the public road network; and
- A 10m-20m wide bushfire asset protection zone around the perimeter of the solar farm (located inside the identified development footprint).





1.3 Site Description

Key features of the Subject Land are summarised in Table 1.3 below.

For the purposes of this BDAR report:

- The Project Boundary refers to Lot 2 DP 1039185.
- The **Subject Land** is the area that would be directly impacted by the Project. In accordance with the BAM (OEH 2017a), the term Subject Land is used in the Stage 1 Biodiversity Assessment and the term **Development Site** is used in the Stage 2 Impact Assessment. The terms Subject Land and Development Site are used interchangeably within this BDAR and are considered to be synonymous for the purpose of this assessment.
- The term **Assessment Area** is used in Stage 1 of this BDAR to refer to the 1,500m buffer surrounding the Subject Land and other areas requiring consideration as part of the biodiversity values assessment in Stage 1 of the BAM (OEH 2017).

Table 1.3 Key Features of the Subject Land

Key Feature	Description	
Location Description	The proposed Bonshaw Solar Farm is located 16 km south of Bonshaw and 66 km north of Inverell.	
Lot Description (Project Boundary)	Lot 2 DP 1039185	
Subject Land	The Subject Land covers approximately 149.24 ha and corresponds to the western portion of Lot 2 DP1039185 and is partially bounded on its northern boundary by Bruxner Highway. To the south and east it is bounded by rural land and to the west, an unsealed road extends from Bruxner Highway to the existing 330kV TransGrid Dumaresq Substation located at the south-western corner.	
Local Government Area Inverell Local Government Area (LGA)		
Elevation	The Project site is dominated by a gently undulating landscape to the north, forming steep slopes to the south and east dissected by second and third order streams. Based on a review of topographic maps and aerial imagery, landforms present within the Subject Land include drainage depressions, gentle to steeply inclining slopes, and upper flat ridges. The elevation at the Bruxner Highway (north boundary) is approximately 335 m and rises up to approximately 420 m in the south-western portion of the site. The ridgelines to the south of the project rise up to approximately 660 m forming the dominant landscape feature.	
Previous Land Use	The Subject Land has undergone vegetation clearing associated with former land use for cropping and is currently used for sheep and cattle grazing.	

1.4 Consultation

Following completion of the Spring 2018 and Summer 2018 survey periods, ERM's principal ecologist Joanne Woodhouse met with BCD (Krister Wearn) to discuss the survey methodology and preliminary survey results, particularly the positive identification of the Bristle-faced Free-tailed Bat (*Mormopterus eleryi*). The results of this meeting and follow up consultation were used to guide the Autumn 2019 survey methodology (additional BAM plots within the areas of native grasslands) and supplementary bat call analysis to confirm the positive species identification. A summary of the consultation is provided in Table 1.4 below.

Table 1.4 Summary of Consultation with BCD during preparation of the BDAR

Date	Contact Name	Summary	Response/Action
12 March 2019	Meeting at BCD Grafton Office attended by: Krister Waern (BCD) Joanne Woodhouse (ERM) Luke Kim (GAIA, by phone)	ERM requested the meeting with BCD to discuss the survey methodology and preliminary survey results to date, particularly the positive identification of the Bristle-faced Free-tailed Bat (<i>Mormopterus eleryi</i>). Key points of discussion: Given the lack of information of the ecology of the Bristle-faced Free-tailed Bat (<i>Mormopterus eleryi</i>) ERM was seeking clarification on the species polygon determination and any additional survey requirements given that we already have a positive species identification. Given that acoustic detectors were the only survey method used, breeding habitat will be assumed to be present. Common Couch (<i>Cynodon dactylon</i>) is listed as native species in NSW however it is not listed in any of the PCT's identified within the Subject Land and has been introduced to the site as a pasture species. ERM was seeking clarification that we can exclude this species in the species richness calculations. ERM confirmed that the threatened flora surveys were undertaken as part of general observations and BAM plot areas, rather than parallel linear transect searches across the entire Subject Land.	BCD confirmed that information on the Bristle-faced Free-tailed Bat is not extensive and they would seek clarification from BCD's threatened species specialist. For other records of threatened microchiropteran bats e.g. Eastern Cave Bat (<i>Vespadelus troughtoni</i>) no additional targeted survey for breeding habitat is required although it is recommended that photographic evidence is provided as justification. Photos of rocky outcrops to be provided to confirm no viability as breeding habitat for this species. BCD will provide advice on the proposed exclusion of Common Couch as a native species within the Subject Land. Additional floristic plots required in derived native grasslands and cleared land. Derived native grassland determination to be updated based on vegetation integrity scores. Offset requirements to be determined based on integrity scores of: >15 (if PCT is a TEC), >17 (if PCT is associated with TS habitat) or >20 (if PCT is not a TEC or is associated with TS habitat). Vegetation mapping to be updated to merge PCT patches which are located within 100m of each other. BCD agreed that the threatened flora survey methodology was reasonable and should be supported by a figure to confirm that the Subject Land was effectively covered during the survey period.

Date	Contact Name	Summary	Response/Action
21 March 2019	Letter from OEH (now BCD) to ERM	 BCD provided formal advice following the meeting held on 12 March and confirmed that: Each vegetation zone (including areas of high weed occurrence and improved pasture areas) are required to detail the vegetation integrity survey plots as described in the BAM. All native plants in NSW (including Common Couch) are required to be considered as native plants for the purposes of applying the BAM, even if they do not naturally occur within the nominated PCT. The draft vegetation mapping appears to be focused on the canopy of the trees. The vegetation mapping should also consider the mid-storey and native ground cover to map the broader extent of the PCT. This is particularly important where canopy spacing within open forest and woodland communities can be up to 100m apart. BCD information suggests that Mormopterus eleryi cannot be easily distinguished from Scotorepens greyii by anabat call. Catching the bat is the most reliable way to determine the species presence. If Mormopterus eleryi is confirmed on site, and based on ERMs current records and the habitat requirements of the species, most of the subject property may be captured when determining the species polygon. 	 ERM undertook and additional four days in the field and an additional 28 floristic survey plots to ensure effective coverage of the all areas of grassland in accordance with advice from BCD and the requirements of the BAM. ERM have updated the assessment and Common Couch (<i>Cynodon dactylon</i>) is now included within the calculations. Vegetation mapping and calculations of areas have been updated. ERM have obtained a supplementary report to confirm the accuracy of the <i>Mormopterus eleryi</i> call identification. Greg Ford (Balance! Environmental) is a recognised expert on Australian bats, with specialist expertise in acoustic analysis of bat echolocation calls for species identification. As described in Appendix I, analysis by Balance! Environmental of numerous full-spectrum data-sets from several regions where <i>M. eleryi</i> and <i>S. greyii</i> are known to coexist has consistently found examples of two distinctive foraging sequence types that concur with the diagnostic descriptions of Corben (2010). An overview of these findings and promotion of the inclusion of feeding buzz analysis in call identification reporting was presented recently at the International Society of Ecoacoustics Congress (Ford 2018). Support for the use of this approach has also been received from several bat-call analysis experts based throughout eastern Australia. Based on this advice, ERM has prepared this assessment based on the results of the call identification and does not consider that harp trapping is required in this instance to confirm presence. Refer to Section 5.7.4.1 and ToS in Appendix E for detailed discussion and assessment. The species polygon has been prepared based on the advice from BCD and is provided in Section 5.7.4.1.

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development Assessment Report

Date	Contact Name	Summary	Response/Action
2 December 2019	Letter BCD to DPIE	BCD provided formal advice to DPIE following public exhibition of Version 1 of this BDAR, dated 26 July 2019. In summary, the following recommendations were provided in relation to biodiversity:	 ERM have engaged Matt Jenkins (Accredited Assessor - North Coast Ecology) to review, update and certify the BDAR and BAM Calculator assessment accordingly.
		 The BDAR should be reviewed, updated and certified to comply with the BC Act and resubmitted as part of the proposal. The BDAR should ensure adequate consideration of the NVR mapping for the site. Ensure the correct application of the streamlined assessment module – clearing paddock trees. The BDAR should be updated to include reference to existing threatened species habitat mapping. The BDAR should provide further justification of the Masked Owl to ensure it has been adequately considered. The BDAR should be updated to remove the recommendation to change the offset requirement for vegetation zone 11. The future vegetation integrity scores for all vegetation zones is to be reduced to zero. The BAM Calculator is to be updated and finalised for review. The mitigation measures as outlined in Table 7.1 of the BDAR should form part of any proposed development consent conditions. 	 NVR mapping has been addressed in Section 4.1.2 of this BDAR. The Streamlined assessment module – clearing of paddock trees has been removed from this BDAR due to the occurrence of threatened species habitat and application of the full BAM on the Subject Land in its entirety. Existing important habitat mapping has been addressed in Section 5.1.1. Masked Owl foraging and breeding habitat is assumed to be present and a species polygon has been provided as shown in Figure 5.3. The vegetation integrity score for vegetation zone 11 was incorrectly calculated in BAM-C. The revised calculations are provided in Section 4.3.2. Future vegetation integrity scores have been reduced to zero for all vegetation zones as shown in Section 4.3.2. The BAM Calculator has been updated and will be finalised upon finalisation and certification of this BDAR. The mitigation measures provided in Table 7.1 shall be used to formulate conditions of development consent.

1.5 **Key Sources of Information used in the Assessment**

The following key information sources were used in preparation of this BDAR: Proposed development layers and project footprint as provided by GAIA.

Key Sources of Information used in the Assessment

Online Resources

Australian Government's Species Profiles and Threats (SPRAT) database. Accessed online at http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl

NSW Department of Planning, Industry and Environment (DPIE) Threatened Biodiversity Profile Search. Accessed online at http://www.environment.nsw.gov.au/threatenedspeciesapp/

NSW Department of Primary Industry (DPI) profiles of threatened species, populations and ecological communities. Accessed online at https://www.dpi.nsw.gov.au/fishing/threatened-species

Commonwealth Department of Environment, Climate Change & Water (DECCW) National Recovery Plan. White Box - Yellow Box - Blakely's Red Gum Grassy Woodland and Derived Native Grassland. A critically endangered ecological community. May 2011.

Commonwealth Department of Environment and Energy (DEE) Protected Matters Search Tool Accessed online at http://www.environment.gov.au/epbc/protected-matters-search-tool

Australia's IBRA Bioregions and sub-bioregions. Accessed online at http://www.environment.gov.au/land/nrs/science/ibra/australias-bioregions-maps

Native Vegetation Regulatory (NVR) Map. Accessed online via https://www.lmbc.nsw.gov.au/Maps/index.html?viewer=NVRMap

NSW Department of Environment and Climate Change (DECC) (2007). Identification Guidelines for Endangered Ecological Communities. White Box - Yellow Box - Blakely's Red Gum Woodland.

NSW Department of Environment and Climate Change (DECC) (2002). Descriptions for NSW (Mitchell) Landscapes, Version 2.

NSW OEH Mitchell Landscapes database v3 2011. Accessed via https://data.gov.au/dataset/e64597db-453c-46be-a352-360b775d2852

NSW OEH's Biodiversity Assessment Method (BAM) calculator. Accessed online via https://www.lmbc.nsw.gov.au/bamcalc

NSW Department of Planning, Industry and Environment (DPIE) BioNet Atlas of threatened biodiversity data collection (TBDC). Accessed online at

http://www.environment.nsw.gov.au/atlaspublicapp/UI_Modules/ATLAS_/AtlasSearch.aspx

NSW Department of Planning, Industry and Environment (DPIE) Vegetation Classification Database. Accessed online via login at http://www.environment.nsw.gov.au/NSWVCA20PRapp/LoginPR.aspx

NSW Department of Planning, Industry and Environment (DPIE) VIS mapping. Accessed online via http://www.environment.nsw.gov.au/research/VISmap.htm

OEH (2017) Biodiversity Assessment Method.

OEH (2018a) Biodiversity Assessment Method Operational Manual - Stage 1.

Key Sources of Information used in the Assessment

NSW Government SEED Mapping. Accessed online via https://geo.seed.nsw.gov.au/Public Viewer/index.html?viewer=Public Viewer&locale=en-AU

NSW Biodiversity Values Map. Accessed online via https://www.lmbc.nsw.gov.au/Maps/index.html?viewer=BVMap

Threatened Species Scientific Committee (TSSC) Advice to the Minister for the Environment and Heritage from the TSSC on Amendments to the List of Ecological Communities under the *Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act) – Box Gum Grassy Woodland.

Literature Review

OEH (2015) BRG-Namoi Regional Native Vegetation Mapping. Technical Notes. NSW Office of Environment and Heritage, Sydney.

Peacock R., Rolhauser A., Thönell J. and Law E. (2009) Extant and potential natural vegetation of Yallaroi, Ashford, Bingara and Inverell 1:100,000 scale map sheet, NSW. NSW Department of Environment, Climate Change and Water.

Other literature consulted during preparation of this BDAR is referenced within the text.

The aerial imagery used in this BDAR is sourced from ESRI World Imagery (DigitalGlobe) dated 27 November 2016. Mapping has been produced using Geographic Information System (GIS). The following maps and data are provided:

- Digital mapping with aerial photography showing 1:1000 or fine
- Site Map (Figure 1.1) as described in subsection 4.2.1.1 and 4.2.1.3 of the BAM
- Location Map (Figure 3.2) as described in subsection 4.2.1.2 and 4.2.1.3 of the BAM

2. LEGISLATIVE CONTEXT

The project has been assessed against key biodiversity legislation and government policy, including:

- Commonwealth Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act);
- NSW Environmental Planning and Assessment Act 1979 (EP&A Act);
- NSW Biodiversity Conservation Act 2016 (BC Act);
- NSW Local Land Services Act 2013 (LLS Act);
- NSW Fisheries Management Act 1994 (FM Act); and
- NSW Biosecurity Act 2015 (BS Act).

Table 2.1 Applicable Legislation, Plan and Guidelines

Commonwealth Legislation

Commonwealth Environmental Protection and Biodiversity Conservation Act 1999 (EPBC Act)

The Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) requires approval of the Commonwealth Minister for Environment (formerly the Minister of Sustainability, Environment, Water, Population and Communities) for actions that may have a significant impact on Matters of National Environmental Significance (MNES). The EPBC Act is administered by the Commonwealth Department of Environment and Energy (DoEE) and lists threatened species, ecological communities and other MNES. Any proposed action that is expected to have an impact on MNES must be referred to the Minister for assessment under the EPBC Act, or assessed under the accredited process between the Commonwealth and the State of NSW.

Matters of National Environmental Significance	Application to the Subject Land	Addressed
World heritage properties	Not identified within the Subject Land	Not applicable
National heritage places	Not identified within the Subject Land	Not applicable
Ramsar wetlands of international importance	Not identified within the Subject Land	Not applicable
Listed threatened species and communities	Threatened species have been recorded within the locality and have potential habitat available within the Subject Land (Development Site). No Threatened Ecological Communities (TEC) are present.	Chapter 5 and Appendix C & D
Internationally protected migratory species	Migratory species are identified as potentially occurring within the Subject Land.	Appendix C & D
Commonwealth marine areas	Not identified within the Subject Land	Not applicable
The Great Barrier Reef Marine Park	Not identified within the Subject Land	Not applicable
Nuclear actions	Not applicable	Not applicable
A water resource, in relation to coal seam gas development and large coal mining development	Not applicable	Not applicable

Statutory Legislation and Guidelines

Biodiversity Conservation Act 2016 (BC Act)

The NSW *Biodiversity Conservation Act 2016* came into effect on 25 August 2017. The BC Act replaced the NSW *Threatened Species Conservation Act 1995*, the NSW *Nature Conservation Trust Act 2001* and parts of the NSW *National Parks and Wildlife Act 1974*. The BC Act establishes mechanisms for:

- The management and protection of listed threatened species of native flora and fauna (excluding fish and marine vegetation) and threatened ecological communities (TECs).
- The listing of threatened species, TECs and key threatening processes.
- The development and implementation of recovery and threat abatement plans.
- The declaration of critical habitat.
- The consideration and assessment of threatened species impacts in development assessment process.
- Biodiversity Offsets Scheme, including the Biodiversity Values Map and method to identify serious and irreversible impacts (SAII).

The BC Act establishes a new regulatory framework for assessing and offsetting biodiversity impacts on proposed developments. Where development consent is granted, the authority may impose as a condition of consent an obligation to retire a number and type of biodiversity credits determined under the Biodiversity Assessment Method (BAM). A Biodiversity Values Map and Biodiversity Offsets Scheme Entry Threshold (BOSET) tool are available to identify the presence of mapped biodiversity values within land proposed for development as well as the clearing thresholds that would trigger application of the BAM.

The Biodiversity Offsets Scheme applies to all local developments, major projects or the clearing of native vegetation where the State Environmental Planning Policy (Vegetation in Non - Rural Areas) 2017 applies. Any of these will also require entry into the Biodiversity Offsets Scheme if they occur on land mapped on the Biodiversity Values Map. ERM has reviewed and can confirm that part Lot 202 DP874273, is not currently mapped on the Biodiversity Values Map (see BOSET report in *Appendix A*) although as a major project, it does trigger the Biodiversity Offsets Scheme.

In terms of the proposed solar farm, the proposed development must take into account species likely to occur within available habitat based on existing records of threatened species and ecological communities, as well as those species likely to occur based on geographic distribution and presence of potential habitat (refer to *Appendix D*)

Water Management Act 2000

A controlled activity approval under the *Water Management Act 2000* (WM Act) is required for certain types of developments and activities that are carried out in or within 40 m of a river, lake or estuary.

The WM Act provides a number of mechanisms for protection of water sources via the water management planning process. If a 'controlled activity' is proposed on 'waterfront land', an approval is required under Section 91(2) of the WM Act. 'Controlled activities' include; the construction of buildings or carrying out of works; the removal of material or vegetation from land by excavation or any other means; the deposition of material on land by landfill or otherwise. 'Waterfront land' is defined as 'the bed of any river or lake, and any land lying between the river or lake and a line drawn parallel to and 40 metres inland from either the highest bank or shore'.

Major projects are exempt from requiring approvals under the Water Management Act 2000.

Biosecurity Act 2015

The NSW *Biosecurity Act 2015* came into effect on 1 July 2017, effectively replacing the *Noxious Weeds Act 1993*, and 13 other Acts, with a single Act. Under the Noxious Weeds Act all landowners have a responsibility to control noxious weeds on their property. Under the Biosecurity Act the same responsibility will apply and will be known as a General Biosecurity Duty.

The General Biosecurity Duty states "Any person who deals with biosecurity matter or a carrier and who knows, or ought reasonably to know, the biosecurity risk posed or likely to be posed by the biosecurity matter, carrier or dealing has a biosecurity duty to ensure that, so far as is reasonably practicable, the biosecurity risk is prevented, eliminated or minimised." The general biosecurity duty applies to all weeds listed in Schedule 3 of the Biosecurity Act (also included as Weeds of National Significance (WoNS).

As detailed in Section 5.7.1, a total of 32 exotic species, including five high threat exotic (HTE) were recorded within the Subject Land. Two of those exotic species, Tiger Pear (*Opuntia aurantiaca*) and Velvet Tree Pear (*Opuntia tomentosa*) are listed as WoNS. A weed species of genus *Senecio* was also recorded, and the potential presence of Fireweed (*Senecio madagascariensis*) cannot be precluded.

A strategic plan for each WoNS has been developed to define responsibilities and identify strategies and actions to control the weed species. These can be downloaded from:

http://www.environment.gov.au/biodiversity/invasive/weeds/weeds/lists/wons.html

Fisheries Management Act 1994

The Fisheries Management Act 1994 provides for the conservation, protection and management of fisheries, aquatic systems and habitats in NSW. Similar to the BC Act, the Fisheries Management Act 1994 lists threatened species, populations and ecological communities of fish and marine vegetation. Consideration of likely occurrence of threatened fish in the waterways in the Subject Land is provided in Section 5.2.

Any waterway crossings along the internal access roads will need to consider an appropriately designed structure that does not obstruct fish passage and will be designed in accordance with the Policy and Guidelines for Fish Habitat Conservation and Management and the Policy and Guidelines for Fish Friendly Waterway Crossings. Notwithstanding this, it is noted that a permit under section 219 would not be required for waterway crossings as section 89J(e) of the EP&A Act excludes projects approved under Part 4 of the EP&A Act from requiring "a permit under section 201, 205 or 219 of the Fisheries Management Act 1994".

SEPP No. 44 - Koala Habitat Protection

SEPP 44 aims to encourage the proper conservation and management of areas of natural vegetation that provide habitat for koalas to ensure a permanent free-living population over their present range and reverse the current trend of koala population decline. A review of SEPP 44 is currently under consideration. The key changes proposed in the amended SEPP 44 relate to the definitions of koala habitat; list of tree species; list of councils; and development assessment process.

SEPP 44 currently applies to land in relation to which a development application has been made within the LGAs as listed in Schedule 1, which includes Inverell LGA. An assessment of Koala habitat values has been provided in Section 5.1.

Local Land Services Act 2013

The Native Vegetation Regulatory (NVR) Map designates areas of NSW to which Part 5A of the *Local Land Services Act 2013* applies. NVR mapping guides the application of the land management code and allowable activities as part of the land management framework for NSW.

An assessment of NVR mapping and relevant land categories is provided in Section 4.1.2.

3. LANDSCAPE FEATURES


The identification of landscape features within the Subject Land was determined in accordance with Section 4 of the BAM (OEH 2017a), as summarised in Table 3.1 below.


Table 3.1 Summary of Landscape Features

Landscape Feature	Description
IBRA Bioregion	Bioregions are large, geographically distinct areas of land with common characteristics such as geology, landform patterns, climate, ecological features, and flora and fauna communities. The Subject Land is located within the NSW Nandewar Bioregion. The Nandewar region consists of hills on Palaeozoic sediments; lithosols and earths, including <i>Eucalyptus albens</i> woodlands. This region is characterized by summer rainfall (Environment Australia 2000).
IBRA Subregion	The Subject Land is located within the NSW Nandewar Northern Complex subregion. This subregion is described as low hills and ranges with abundant rock (granite) outcrop and tors. Short, steep gorges of major rivers. Karst landscapes on limestone. This IBRA Subregion is mapped as a Priority 4 investment region (OEH 2017b).
NSW Landscape Regions and Area	The Subject Land is mapped in the Ashford Mole Valleys Mitchell Landscape.
Percent Native Vegetation	A combination of existing mapping and ground truthed vegetation was used to estimate the Percent Native Vegetation Cover as follows:
	Existing vegetation mapping was used to estimate the percent native vegetation within the 1500 m buffer area surrounding the Subject Land (see Figure 3.2). In the buffer area, PCTs were allocated based on existing vegetation mapping of the Inverell Shire as mapped by Peacock et. al. (2009) (VIS ID 3794) and aerial imagery. GIS was used to estimate the area.
	 Ground-truthed vegetation was used to estimate the portion of native vegetation within the Subject Land. PCT allocation was based on existing mapping and current vegetation condition (see Chapter 4).
	The Percent Native Vegetation Cover was calculated by estimating the percent cover of native vegetation in the buffer area and Subject Land relevant to the benchmark for the PCT.
	Ten PCTs are mapped in the 1500 m buffer area (Table 4.1), covering approximately 1,273.70 ha. This represents 67.63% native vegetation cover within the assessment area (1,883.30 ha).
Cleared Areas	Approximately 107.79 ha (72 %) of land within the Subject Land represents cleared grazing land. Cleared areas are primarily grazed land (cattle and sheep), which provides limited foraging habitat for native species. Sheep can graze very close to the ground and like other livestock can lead to loss of vegetation and soil erosion particularly during the dry conditions encountered during the survey period.

Client: GAIA Australia

Landscape Feature	Description
Rivers, Streams and Estuaries	Three major un-named creeks (tributaries of the Dumaresq River) are present within the Project Boundary (Lot 2 in DP 1039185). The largest of these creeks traverses the development site in a general north-southerly direction and is mapped as a third order stream (Strahler, 1952). The other two creeks are mapped as First Order Streams, and are located to the east and west of the third order stream. These streams were all noted to be dry (or very low water levels) at the time of the survey due to extended drought conditions. A review of the NSW Department of Infrastructure (DPI) threatened freshwater fish records (accessed via SEED), confirms that these creeks are not mapped as habitat of threatened aquatic species. The nearest creek with threatened aquatic species habitat is Little Oak Creek, located approximately 1.7 km west from the Subject Land. Aquatic habitat is described further in Section 5.2 and a preliminary aquatic impact assessment is provided in Section 5.7.7.
Wetlands	No wetlands occur in or adjacent to the Development Site. Aerial photographs show six farm dams are present within the Subject Land. These dams were all noted to be dry (or very low water levels) at the time of the survey due to extended drought conditions.
Connectivity Features	A wildlife corridor is a link of wildlife habitat, generally native vegetation, which joins two or more larger areas of similar wildlife habitat. Corridors are critical for the maintenance of ecological processes including allowing for the movement of animals and the continuation of viable populations. The Subject Land is located immediately north of a regional vegetated corridor that connects Crooked Creek National Park and Torrington State Conservation Area. Isolated paddock trees scattered across the development site also represent limited connectivity features for highly mobile species to travel across the landscape. The creeks present within the Subject Land have the potential to provide aquatic connectivity to Dumaresq River although it is noted that these streams were all dry (or very low water levels) at the time of the survey due to extended drought conditions and any connectivity would be seasonal at best. The Subject Land is not located within (or close to) any identified migratory bird flyways including the East Asian – Australasian Flyway and the West Pacific Flyway.
Areas of Geological Significance	No karsts, caves, crevices or cliffs or other areas of geological significance occur within the Development Site or are likely to occur within the broader Subject Land.
Areas of Outstanding Biodiversity Value (AOBV)	No Areas of Outstanding Biodiversity Value occur within the Development Site of the broader Subject Land.
High Biodiversity Values Map	In accordance with the NSW Biodiversity Values Map (See Figure 3.1 and BOSET report in Appendix A), the Subject Land does not contain high biodiversity values. The nearest area mapped with high biodiversity values are Little Oak Creek and Beardy River, located at approximately 1.7 km to the west and 2.3 km to the east, respectively.
Native Vegetation Regulatory (NVR) Mapping	The Local Land Services Act 2013 (LLS Act) sets out several land categories that appear on the Native Vegetation Regulatory Map (NVR Map). The main purpose of this map is to govern clearing of vegetation associated with agricultural activities in rural areas. NVR mapping is shown in Figure 3.1 and addressed in detail in Section 4.1.2.





4. NATIVE VEGETATION

The extent of native vegetation within the development site was determined in accordance with Section 5 of the BAM (OEH 2017a).

The development site is currently primarily used for sheep and cattle grazing. Native vegetation is highly modified by both historical and ongoing management practices including clearance of the original vegetation type, livestock grazing and weed invasion. A large portion of the development site is cleared and disturbed land, the majority of which no longer reflects the species composition of the community from which it was likely derived. Forb diversity and coverage is very low, dominated by those species tolerant of heavy grazing.

Woodland areas show evidence of varying degrees of impact, the most notable being on-going clearing and presence of monospecific stands (e.g. White Cypress Pine) due to a long history of timber removal. Vegetation within the development site comprises:

- 29.17 ha Native Vegetation;
- 11.80 ha Derived Native Grasslands; and
- 107.79 ha Disturbed Grasslands.

These vegetation areas total 148.76 ha. The remaining 0.48 ha of the development footprint include cleared lands and six farm dams.

4.1 Native Vegetation Assessment Methodology

4.1.1 Review of Existing Information

A search was undertaken of the NSW Department of Planning, Industry and Environment (DPIE) Vegetation Information System (VIS) database and NSW SEED (Sharing and Enabling Environmental Data) mapping to access existing vegetation mapping information within the Subject Land. The State Vegetation Type Map – Borders Rivers Gwydir / Namoi (VIS ID 4681) provides vegetation mapping for the Inverell Shire. This state vegetation type map (SVTM) is based on regional mapping in Peacock *et. al.* (2009) (VIS ID 3794) and identifies seven Plant Community Types (PCTs) within the Subject Land and eleven PCTs within the 1,500m buffer area as listed in Table 4.1 below and depicted in Figure 4.1.

Table 4.1 Mapped Vegetation Communities (SVTM VIS 4681) within the Buffer Area and Subject Land (not ground-truthed)

PCT ID	PCT Name	Extent in Buffer Area (ha)	Extent in Development Site (ha)
NA	Cleared Land	609.60	108.03
1	Candidate Native Grasslands	389.86	0
78	River Red Gum riparian tall woodland / open forest wetland in the Nandewar Bioregion and Brigalow Belt South Bioregion	27.38	0
84	River Oak – Rough-barked Apple – Red Gum – Box riparian tall woodland (wetland) of the Brigalow Belt South and Nandewar Bioregions	14.73	0
505	Black Cypress Pine - Tumbledown Red Gum - Narrow-leaved Ironbark - Stringybark She Oak open forest on acid volcanics of the western New England Tableland Bioregion	1.07	0
516	Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	42.04	3.00

PCT ID	PCT Name	Extent in Buffer Area (ha)	Extent in Development Site (ha)
544 *	Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / Woodland of the Nandewar Bioregion and New England Tableland Bioregion	8.37	0.83
549	Silver-leaved Ironbark – Black Cypress Pine +/- White Box shrubby open forest mainly in the northern Nandewar Bioregion	194.97	0
578	Tumbledown Red Gum - Black Cypress Pine - Caley's Ironbark shrubby open forest of the Nandewar Bioregion and western New England Tableland Bioregion	2.96	0
594	Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	225.39	13.65
596	Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	366.21	11.70
599	Blakely's Red Gum - Yellow Box grassy tall woodland on flats and hills in the Brigalow Belt South Bioregion and Nandewar Bioregion	0.71	0

^{*} PCT 544 corresponds to the BC Act listed endangered ecological community (EEC) "White Box Yellow Box Blakely's Red Gum Woodland", which also corresponds to part of the EPBC listed critically endangered ecological community (CEEC) "White Box Yellow Box Blakely's Red Gum Woodland".

4.1.2 Native Vegetation Regulatory Mapping

The Local Land Services Act 2013 (LLS Act) sets out several land categories that appear on the Native Vegetation Regulatory Map (NVR Map). The main purpose of this map is to govern clearing of vegetation associated with agricultural activities in rural areas. However, there are important implications that affect DA assessment for one of the land categories: Category 1 – Exempt Land. This category is broadly intended to identify land that has been previously legally cleared or that comprises low conservation value grasslands. The Category 1 layer has not yet been released, but will appear as a blue layer on the completed map. While the map is being finalised, rural landholders are responsible for determining the categorisation of their land based on the criteria listed in the LLS Act.

Clearing of native vegetation for development on land that meets the definition of Category 1 land does not require assessment or offsetting under the BAM. In practice, this means that native vegetation on Category 1 land is not included in any area clearing calculations when determining whether a BDAR should be prepared. A BDAR may still be required if the development will result in:

- A prescribed impact (these are listed in <u>clause 6.1 of the Biodiversity Conservation</u> <u>Regulation</u>), not including native vegetation clearing associated with the prescribed impact
- A likely significant impact on threatened species or ecological community based on the threatened species test of significance

As identified in Section 60F of the LLS Act (Transitional arrangement until preparation of maps), an area is taken, during the transitional period, to be low conservation value grasslands if it comprises only groundcover whose clearing was permitted by section 20 of the Native Vegetation Act 2003, as in force immediately before the repeal of that Act. Note. Generally that section permitted clearing if the vegetation comprised less than 50% of indigenous species of vegetation.

Disturbed grassland areas of the Subject Land (i.e. previously cleared grazing land) are not considered to represent Category 1 – exempt land as they do not meet the definition provided for low conservation value grasslands (i.e. remaining groundcover vegetation is generally not comprised of <50% of indigenous species). Furthermore, given the presence of some threatened species habitat components (e.g. scattered hollow-bearing trees and stags) the application of the BAM to the entirety of the Subject Land is considered to be appropriate in this instance.

The Subject Land is therefore considered to represent Category 2 – Regulated Land. Some areas of steep land in the south-eastern portion of Lot 2 DP1039185 are currently mapped as Vulnerable Regulated Land, as shown in Figure 3.1.

4.1.3 Targeted Floristic Survey

Based on the results of the desktop assessment presented in Table 4.1 above, flora surveys were undertaken in September 2018, December 2018, March 2019 and January 2020 as follows:

- The first survey was undertaken between the 10th and 14th September 2018 by ERM ecologists, Joanne Woodhouse and Dr Adriana Corona Mothe. The aim of the initial survey was to determine the PCTs on the development site and their condition on site. Random meander transects were conducted in areas of native vegetation and rapid vegetation assessments were undertaken to record the presence of native vegetation, threatened flora and to ground-truth existing mapping. PCT's were identified from the native species present, landforms and physiography and location within the IBRA subregion with reference to the BioNet Vegetation Classification Database. The Subject Land was then stratified into areas of similar condition class to determine vegetation zones for each PCT and detailed floristic surveys were undertaken on 11 12 September 2018.
- The second survey was undertaken between the 10th and 14th December 2018 by ERM ecologists Tom Cotter and Dr Adriana Corona Mothe. Vegetation integrity plots (20m by 50m) were established in each vegetation zone over three days on the 11 13 December 2018. Data was collected on the composition, structure and function of the vegetation in accordance with the methodology presented in the BAM 2017 and under the directions of persons accredited under the BAM. The number of plots for each zone was based on the area of each zone and the minimum number of plots and transects required per zone area as specified by the BAM.
- The third survey was undertaken between the 25 and 29 March by Dr Toivo Zoete and Dr Adriana Corona Mothe and included additional vegetation integrity plots over four days from 25 to 28 March 2019.
- The fourth survey was undertaken on 21 and 22 January 2020 by accredited assessor Matt Jenkins (North Coast Ecology) and ecologist Chloe Hanrahan (ERM). This survey was completed in order to review existing PCT and vegetation zone mapping prepared during previous surveys.

A total of four PCTs comprising 12 vegetation zones were identified on the Subject Land as listed in Table 4.4 and mapped in Figure 4.2. Based on the assessment of each PCT, it was decided to stratify each of the four PCTs based on the following:

- Absence of upper stratum, condition of the stratum and composition. Due to the high level of disturbance at the Development Site, treed areas were commonly represented by a single species of trees or the presence of trees were significantly reduced when compared with tree composition and structure as per PCTs descriptions.
- Structure and function of the vegetation in other stratum, e.g. ground cover.
- Presence of exotic species and their abundance/cover as well as bare ground.

Each vegetation zones was considered homogeneous and well represented by the plot data. As outlined in Table 4.2, a total of 42 flora plots were collected in accordance with section 5 of the BAM (OEH 2017) by ecologists trained in the BAM (Joanne Woodhouse and Adriana Corona Mothe) and under the direction of Dr Adriana Corona Mothe who is accredited under the BAM. An additional two plots were completed during the January 2020 survey by Matt Jenkins (accredited under the BAM) and Chloe Hanrahan (ERM ecologist). At each plot location the following was undertaken:

- one 20 x 20 m plot for assessment of composition and structure; and
- one 20 x 50 m plot for assessment of function, including a series of five 1 x 1 m plots to assess average leaf litter cover.

The assessment of composition and structure recorded species name, stratum, growth form, cover and abundance rating for each species present within the plot. Cover (foliage cover) was estimated for all species rooted in or overhanging the plot, and recorded using decimals (if less than 1%, rounded to whole number (1 - 5%) or estimated to the nearest 5% (5 - 100%). Abundance was counted (up to 20) and estimated above 20.

The assessment of function recorded the number of large trees, the presence of tree stem size class, tree regeneration, number of trees with hollows and length of fallen logs, as well as leaf litter cover within the 20 x 50 m plot and five 1 x 1 m subplots. The minimum number of plots and transects per vegetation zone was determined using Table 4 of the BAM (OEH 2017a). Portions of the land within the study area are either cleared land, cropped or consist of exotic grassland. Under the BAM (OEH 2017a), land not containing native vegetation is not subject to assessment beyond Section 5.4 (determination of a vegetation integrity score). A large proportion of grassland occurs within the development site, ranging from exotic with few native grass species, to areas of low diversity native pasture. Cleared land and disturbed grasslands were assigned to the most likely PCT and mapped as "Disturbed Grasslands".

4.1.4 PCT Identification

The method used for PCT identification included the following:

- Use of the BioNet Vegetation Classification's Community Identification keys. Search Criteria used included Vegetation Formation, Vegetation Class, IBRA Region and selected community species in the upper / mid / low stratum. This search produced a shortlist of potential PCTs.
- Selection of the PCT was undertaken by comparing the descriptions of each potential PCT with characteristics of the vegetation such as landform location, species composition and other landscape features relevant to the vegetation community. When the vegetation community under assessment was likely to conform to more than one PCT, the decision on a given PCT over other options was based on presence of characteristic species, species richness and other aspects of the PCT description.
- The condition of the PCT was defined based on the absence of upper stratum, dominance of exotics over natives and percentage bare land present.
- Where derived grasslands were identified, the most likely PCT was allocated based on number of native species shared with a given PCT, its location, landscape features and the neighbouring PCTs.

4.2 PCTs Identified in the Subject Land

Determination of PCTs within the Subject Land identified the presence of four Plant Community Types (PCT) as shown in Table 4.2 below.

Table 4.2 **Plant Community Types Occurring within the Subject Land**

	Plant Community Type	Vegetation	Vegetation Class	
ID No	Name	Formation		
516	Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	Grassy Woodlands	Western Slopes Grassy Woodlands	
544	Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion	Grassy Woodlands	Western Slopes Grassy Woodlands	
594	Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	Dry Sclerophyll Forests (Shrub/grass sub-formation)	North-west Slopes Dry Sclerophyll Woodlands	
596	Tumbledown Red Gum – White Cypress Pine – Silver- leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	Dry Sclerophyll Forests (Shrub/grass sub-formation)	North-west Slopes Dry Sclerophyll Woodlands	

A detailed description of each of these PCTs follows in Table 4.3 to Table 4.6.

PCT 594: Silver-leaved Ironbark – White Cypress Pine shrubby Table 4.3 open forest of Brigalow Belt South Bioregion and Nandewar Bioregion

PCT 594: Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South **Bioregion and Nandewar Bioregion**

Vegetation Formation	Dry Sclerophyll Forest (Shrub/grass sub-formation)		
Vegetation Class	North-west Slopes Dry Sclerophyll Woodlands		
Vegetation Type	PCT ID:	594	
	PCT's Common Name:	Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	
	Condition	Moderate Low	
Description	Tall woodland to open forest dominated by Silver-leaved Ironbark (<i>Eucalyptus melanophloia</i>) and Black Cypress Pine (<i>Callitris endlicheri</i>). Other trees may include White Box (<i>Eucalyptus albens</i>), White Cypress Pine (<i>Callitris glaucophylla</i>), Tumbledown Red Gum (<i>Eucalyptus dealbata</i>), Narrow-leaved Ironbark (<i>Eucalyptus crebra</i>) and less often		
Smooth-barked Apple (Angophora leiocarpa) The shrub la includes Olearia elliptica, Notelaea microcarpa var. microcurceolatus, Xanthorrhoea johnsonii, Acacia elongata, Brey		a microcarpa var. microcarpa, Hibbertia obtusifolia, Melichrus	

Client: GAIA Australia

18 May 2020

PCT 594: Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South **Bioregion and Nandewar Bioregion**

Acacia leiocalyx subsp. leiocalyx, Acacia penninervis var. penninervis, Beyeria viscosa and Pimelea neo-anglica. The ground layer is sparse with frequent grass species including Eragrostis leptostachya, Enneapogon gracilis, Aristida ramosa, Cymbopogon refractus, Austrostipa scabra and Aristida vagans. Forb species include Dichondra repens, Vittadinia cuneata, Brunoniella australis, Dichondra sp. A, Rostellularia adscendens subsp. adscendens, Scleria mackaviensis and Phyllanthus virgatus. The rock ferns Cheilanthes sieberi subsp. sieberi and Cheilanthes distans are common. Climbers include Parsonsia eucalyptophylla, Desmodium brachypodum and Desmodium varians. Occurs on shallow loamy sand soils mostly derived from sandstone on hills with rocky outcrops mainly in the northern Nandewar Bioregion from north of Bingara to Arakoola with outliers in the Mole River district. The PCT is found on conglomerate and sandstone on hills and low hills.

At the Subject Land the PCT is present in both moderate and low condition. Dominant tree species recorded were Silver-leaved Ironbark, White Box and White Cypress Pine. Some areas of the community lack Eucalypt trees completely and identification of likely PCT was based on landform and characterisic ground cover species. Species characteristic of the community include Oxalis perennans, Poison Pimelea (Pimelea neo-anglica), Mielichrus urceolatus, Purple Wiregrass (Aristida ramosa), Speargrass (Austrostipa scabra subsp. scabra), Yellow Burr-daisy (Calotis lappulacea), Slender Flat-sedge (Cyperus gracilis), Barbed Wrie Grass (Cymbopogon refractus), Cheilates sieberi, Glycine clandestina and Corrugated Sida (Sida corrugata).

Approximate extent within Subject Land Moderate: 4.10 ha Low: 9.54 ha

BAM Plots in PCT

Moderate: Four plots (P2, P7, P15 and P16) Low: five plots (P1, P9, P17, P21 and P40)

Justification of evidence used to identify the **PCT**

PCTs 594 and 596 had similar distributional patterns and a total of nine species characteristic of each PCT was recorded in some of these plots, including White Cypress Pine, which is listed as a dominant species in both PCTs.

The decision was made to select PCT 594 based on the abundance of Austrostipa scabra subsp. scabra, Sporobolus creber and/or Poison Pimelea which is are characteristic species of PCT 594 and are absent in PCT 596.

This PCT was identified relying on the presence of species typical of the PCT as follows:

Upper stratum: dominated by Silver-leaved Ironbark

Mid-stratum: presence of White Cypress Pine and/or Poison Pimelea

Ground stratum: presence of Oxalis perennans, Austrostipa scabra subsp. scabra, Cymbopogon refractus and Aristida ramosa

The condition of this PCT was selected based on the presence of upper stratum, cover of exotic species (e.g. Tiger Pear) and intensity of sheep and cattle grazing.

TEC Status

PCT 594 is not associated with a TEC

Estimate of percent cleared

53%

Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page 27

PCT 594: Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South **Bioregion and Nandewar Bioregion**

Images of PCT within Subject Land



Photograph 4-1 View of PCT 594 in Plot 2



Photograph 4-2 View of PCT 594 in Plot 21

PCT - Plant Community Type; TEC - Threatened Ecological Community

Page 29

PCT 596: Tumbledown Red Gum - White Cypress Pine - Silver-Table 4.4 leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion

PCT 596: Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion

mainly in the nort	hern Nandewar Bioregion						
Vegetation Formation	Dry Sclerophyll Forest (Shrub/grass sub-formation)						
Vegetation Class	North-west Slopes Dry Sclerophyll Woodlands						
Vegetation Type	PCT ID:	596					
	PCT's Common Name:	Tumbledown Red Gum – White Cypress Pine – Silver- leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion					
	Condition	Moderate Low Very Low Derived Native Grassland					
Description	Mid-high to low woodland or open forest dominated by Tumbledown Red Gum (Eucalyptus dealbata), White Cypress Pine (Callitris glaucophylla) and Silver-leav Ironbark (Eucalyptus melanophloia). Other tree species may include Caley's Ironbark (Eucalyptus caleyi subsp. caleyi), Rough-barked Apple (Angophora floribunda), Lifruited Bloodwood (Corymbia dolichocarpa), Black Cypress Pine (Callitris endliche Rusty Fig (Ficus rubiginosa). There is usually a dense shrub/small tree layer of sp. such as Leptospermum brevipes, Acacia cheelii, Notelaea microcarpa var. microcal Melichrus urceolatus, Xanthorrhoea johnsonii and Leucopogon muticus. The vine Pandorea pandorana is often abundant. The ground layer is sparse with Cheilanti distans, Cheilanthes sieberi subsp. sieberi, Aristida vagans, Cymbopogon refracti Entolasia stricta and Aristida ramosa most frequent. Forb species include Plectral parviflorus, Commelina cyanea, Scleria mackaviensis, Dichondra species A, Cale lappulacea, Phyllanthus virgatus, Vittadinia sulcata and Galium guadichaudii. Occishallow loamy sand soils in hilly areas at low altitudes.						
	is present in a disturbed state due to the long history of emoval. Upper stratum is represented by Tumbledown Red Silver-leaved Ironbark (<i>Eucalyptus melanophloia</i>), Black theri) and White Cypress Pine (<i>Callitris glaucophylla</i>). Other ble Wiregrass (<i>Aristida ramosa</i>), Bristly Cloak Fern Wire Grass (<i>Cymbopogon refractus</i>), Slender Flat-sedge ck-trefoil (<i>Desmodium varians</i>), Urn-heath (<i>Melichrus</i> ightshade (<i>Solanum sturtianum</i>), Yellow Burr-daisy (<i>Calotis</i> is Tree (<i>Xanthorrhoea johnsonii</i>), <i>Glycine clandestina</i> and						
Approximate extent within the Subject Land	Moderate: 11.15 ha Low: 0.25 ha Very Low: 0.30 ha Derived Grassland in Low cor	ndition: 9.04					
BAM Plots in PCT	Moderate: four plots (P8, P10 Low: two plots (P3 and P5) Very Low: one plot (P41) Derived Grassland in Low cor	ndition: four plots (P14, P23, P26 and P29)					

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

PCT 596: Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion

Justification of evidence used to identify the PCT

This PCT was identified relying on the presence of dominant tree species and other typical species of the PCT as follows:

- Upper stratum: dominated by Tumbledown Red Gum and Silver-leaved Ironbark
- Mid-stratum: dominated by White Cypress Pine, presence of other species such as Black Cypress Pine
- Ground stratum: presence of Cheilantes distans, a frequent species in the PCT; along with other species such as Thargomindah Nighshade.

The condition of this PCT was selected based on the presence of upper stratum, cover of exotic species (e.g. Tiger Pear) and intensity of sheep and cattle grazing.

Species relied upon for PCT identification

Tumbledown Red Gum, Silver-leaved Ironbark, Black Cypress Pine, White Cypress Pine, Barbed Wire Grass and Bristly Cloak Fern (*Cheilantes distans*).

TEC Status

PCT 596 is not associated with a TEC

Estimate of percent cleared

38%

Images of PCT within the Development Site



Photograph 4-3 View of PCT 596 in Plot 10

PCT 596: Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion



Photograph 4-4 View of PCT 594 in Plot 5

Notes: PCT - Plant Community Type; TEC - Threatened Ecological Community

Table 4.5 PCT 516: Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion

PCT 516: Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion

Vegetation Formation	Grassy Woodla	Grassy Woodlands							
Vegetation Class	Western Slope	Western Slopes Grassy Woodlands							
Vegetation Type	PCT ID:	516							
	PCT's Common Name:	Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion							
	Condition	Moderate (occurring adjacent to Subject Land) Very Low (occurring within Subject Land) Derived Native Grasslands in Moderate condition (occurring within Subject Land)							
Description	This PCT is described as widespread, mid-high to very tall woodland or of forest occurring on flat to undulating sites at low to mid elevation, mainly or grained soils on sedimentary and metasedimentary substrates. It is domin by Grey Box								

Client: GAIA Australia

18 May 2020

PCT 516: Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion

(Eucalyptus moluccana) (crossing with Eucalyptus albens at sites with basalt influence) which is associated with Blakely's Red Gum (Eucalyptus blakelyi), Yellow Box (Eucalyptus melliodora) and/or White Cypress Pine (Callitris glaucophylla), with occasional Rough-barked Apple (Angophora floribunda). A sparse shrub layer of Notelaea microcarpa var. microcarpa and/or Dodonaea viscosa subsp. spatulata is sometimes present. The ground layer is dense and diverse, containing a mix of grasses and forbs including Cymbopogon refractus, Austrostipa scabra subsp. scabra, Aristida ramosa, Asperula conferta, Bothriochloa decipiens, Daucus glochidiatus, Dichanthium sericeum subsp. sericeum, Carex inversa, Chrysocephalum apiculatum, Cymbonotus lawsonianus, Dichondra sp. A, Glycine tabacina, Poa sieberiana, Desmodium varians, Eremophila debilis, Austrodanthonia racemosa var. obtusata, Austrostipa verticillata, Dichondra repens, Hydrocotyle laxiflora, Plantago debilis, Rostellularia adscendens subsp. adscendens, Chloris ventricosa, Geranium solanderi var. solanderi, Cyperus gracilis, Hypericum gramineum, and (rarely) Panicum paludosum. This community is found in

the undulating floors of the major river valleys of the Nandewar and far western New England bioregions.

At the Subject Land, the community is highly disturbed with few characteristic species present, including: White Cypress Pine (*Callitris glaucophylla*), Silverleaved Ironbark (*Eucalyptus melanophoia*), Small-leaf Bluebush (*Maireana microphylla*), Purple Wiregrass (*Aristida ramosa*), *Dichondra* sp. A, Red Grass (*Botriochloa decipiens*) and Speargrass (*Austrostipa scabra subsp. scabra*),

Approximate extent within the Subject Land

Moderate: 0 ha Very Low: 3.00 ha

Derived Grassland in Moderate condition: 2.76 ha

BAM Plots in PCT

Moderate: one plot (P38)

Very Low: three plots (P31, P39 and P42)

Derived Grassland in Moderate condition: two plots (P28 and P32)

Justification of evidence used to identify the PCT

PCT 516 was identified based on the presence of characteristic species of this vegetation community and its distribution at landform elements such as drainage depression.

This PCT was identified relying on the presence of species typical of the PCT as follows:

- Upper stratum: Grey Box
- Mid-stratum: presence of Small-leaf Bluebush
- Ground stratum: presence of Dichondra sp. A, Purple Wiregrass, Speargrass and Red GrassDistribtuion near drainage depression

The condition of this PCT was selected based on the presence of upper stratum, cover of exotic species (e.g. Tiger Pear) and intensity of sheep and cattle grazing.

Grey Box, Small-leaf Bluebush, Purple Wiregrass, Dichondra sp. A, Red Grass

Species relied upon for PCT identification

PCT 516 is not associated with a TFC

TEC Status

85%

and Speargrass

Estimate of percent cleared

Images of PCT within the Subject Land

Tableland Bioregion



Photograph 4-5 View of Moderate Condition PCT 516 in Plot 38



Photograph 4-6 View of Very Low Condition PCT 516 in Plot 42

Notes: PCT - Plant Community Type; TEC - Threatened Ecological Community

18 May 2020

Page 34

Table 4.6 PCT 544: Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion

PCT 544: Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion

Vegetation Formation	Grassy Woodlands					
Vegetation Class	Western Slopes Grassy Woodlands					
Vegetation Type	PCT ID: 544					
	PCT's Common Name:	Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion				
	Condition	Moderate (occurring adjacent to Subject Land) Low (occurring within Subject Land)				
Description	barked Apple (Ar (Eucalyptus blak (Eucalyptus melli (Eucalyptus albe Notelaea microca transmontanum, Cassinia quinque anglica, Mairean dense and is congrasses including Austrostipa vertici blady grass (Imp Dichondra repen adscendens subsicynoglossum au and Carex inversat some sites. Oc loamy-clay soil a and west New Er Riparian Woodla barked Apple woweed infested. At the Subject Larecorded were R (Eucalyptus melli	ribed as tall open forest to woodland dominated by Rough- ngophora floribunda) in association with Blakely's Red Gum elyi), White Cypress Pine (Callitris glaucophylla) or Yellow Box iodora) or Black Cypress Pine (Callitris endlicheri) or White Box ns). The shrub layer is sparse comprising species such as arpa var. microcarpa, Leptospermum polygalifolium subsp. Olearia ellitptica, Leptospermum brevipes, Beyeria viscosa, efaria, Dodonaea viscosa subsp. angustifolia, Pimelea neo- a microphylla and Acacia neriifolia. The ground layer is often nprised of the mat-rush (Lomandra longifolia) and a range of g Microlaena stipoides var. stipoides, Bothriochloa macra, cillata, Echinopogon caespitosus, Eragrostis leptostachya and erata cylindrica). Forb species include Dichondra sp. A, s, Plantago debilis, Oxalis perennans, Rostellularia sp. adscendens, Swainsonia galegifolia, Urtica incisa, strale and Vittadinia dissecta. Sedges include Cyperus gracilis sa. The invasive weed Coolatai grass (Hyparrhenia hirta) occurs cours on a variety of alluvial or colluvial soils from sand to long stream banks and on valley flats throughout the Nandewar ngland Tableland Bioregions. Grades into ID84 River Oak nd or ID78 River Red Gum woodland. Similar to the Rough- odland ID281 in the BBS Bioregion. Mainly cleared and often threatened community. and the PCT is present in low condition. Dominant tree species ough-barked Apple (Angophora floribunda), Yellow Box iodora) and White Cypress Pine (Callitris glaucophylla). Other				
	characteristic species included Notelaea microcarpa var. microcarpa, Beyeria viscosa, Maireana microphylla, Pimelea neo-anglica, Microlaena stipoides var. stipoides and Austrostipa verticillata.					
Approximate extent within the Subject Land	Moderate: 0 ha Low: 0.83 ha					
BAM Plots in PCT	Moderate: one pl Low: one plot (P					
Justification of evidence used to identify the PCT	vegetation comm	entified based on the presence of characteristic species of this nunity and its distribution at landform elements such as bank ream bed, stream channel.				

PCT 544: Rough-barked Apple - White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion

This PCT was identified relying on the presence of species typical of the PCT as follows:

- Upper stratum: Rough-barked apple and Yellow box
- Mid-stratum: presence of Notelaea microcarpa var. microcarpa, Beyeria viscosa, Maireana microphylla, Pimelea neo-anglica
- Ground stratum: presence of Microlaena stipoides var. stipoides and Austrostipa verticillata

The condition of this PCT was selected based on the cover/integrity of upper stratum.

Species relied upon for PCT identification

As above

TEC Status

PCT 544 corresponds to the NSW listed endangered ecological community (EEC) "White Box Yellow Box Blakely's Red Gum Woodland".

The occurrence of this TEC on the Subject Land is not considered to satisfy the minimum condition criteria for the Commonwealth listing of this TEC. This determination has been made on the basis of very small patch sizes occurring on the Subject Land, low diversity of native understorey species and absence of "important species" as described in the Threatened Species Scientific Committee listing advice and National Recovery Plan for the TEC.

Estimate of percent cleared

65%

Images of PCT within the Subject Land



Photograph 4-7 View of Moderate Condition PCT 544

Project No.: 0470861

Client: GAIA Australia

PCT 544: Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion



Photograph 4-8 View of Low Condition PCT 544

Notes: PCT - Plant Community Type; TEC - Threatened Ecological Community

4.2.1 Cleared and Disturbed Land

Cleared and disturbed land within the Subject Land has a long history of heavy grazing and does not correspond to any recognisable PCT. However to facilitate this assessment and in accordance with the BAM (and advice from BCD confirming that Common Couch is to be included in the BAM calculator as a native species), these areas have been assigned a likely PCT based on surrounding vegetation and location within the landscape. Three vegetation zones for cleared and disturbed land were assessed:

- 516_Disturbed Grassland: a total of 7.39 ha of cleared and/or highly disturbed land. A total of three BAM plots were assessed in this vegetation zone (P27, P33 and P35).
- 594_Disturbed Grassland: a total of 48.81 ha of cleared and/or highly disturbed land. A total of eight BAM plots were assessed in this vegetation zone (P6, P18, P19, P22, P24, P25, P36 and P37).
- 596_Disturbed Grassland: a total of 50.59 ha of cleared and/or highly disturbed land. A total of five BAM plots were assessed in this vegetation zone (P4, P13, P20, P30 and P34).

Cleared and disturbed land was widespread across the Subject Land which is consistent with its historical and current land uses which include clearing, cropping and grazing. Cattle, sheep and goats were present across the Subject Land and a very intense level of grazing was observed. Flora species included a mix of native and exotic species, including high threat exotics (see flora list in Appendix H). Upper and mid-stratum were generally absent, with the exception of some isolated scattered trees as shown in Photographs 4-7 to 4-10.

4.3 Vegetation Integrity Assessment

4.3.1 Vegetation Zones and Survey Effort

The random meander surveys and detailed floristic plots have been used to assist the delineation of zones. Each of the identified PCTs were stratified into zones with a similar broad condition state. These zones were defined based on the overstorey condition, understorey condition and observed land management practices.

A total of 9 native vegetation zones plus cleared land were identified in the development site as listed in Table 4.6 and mapped in Figure 4.2.

A total of 44 vegetation integrity plots were collected in accordance with section 5 of the BAM (OEH 2017) to determine their condition.



Photograph 4-7: View of cleared land in plot P4



Photograph 4-8: View of cleared land in plot P13



Photograph 4-9: View of cleared and disturbed land in plot P35



Photograph 4-10: View of cleared and disturbed land in plot P22

Table 4.7 **Vegetation Zones on the Subject Land**

Zone ID	PCT ID	Stratification Unit / Condition Class	Extent within development site	BAM plots required	Survey effort (number of Plots)	Patch Size (ha) (Number of patches)	Reference
Native	Vegetation						
1	594_ Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	Moderate	4.10 ha	2	4	>1000 ha (1)	Table 4.3
2	594_ Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	Low	9.54 ha	3	5	<5 ha (3); >1000 ha (1)	
3	596_ Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	Moderate	11.15 ha	3	4	>1000 ha (1)	Table 4.4
4	596_ Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	Low	0.25 ha	1	2	>1000 ha (1)	
5	596_ Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	Very Low	0.30 ha	1	1	>1000 ha (1)	
6	596_Derived Grassland	Low	9.04 ha	3	4	<5 ha (5)	
7	516_Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	Very Low	3.00 ha	2	3	>1000 ha (1)	Table 4.5

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

Zone ID	PCT ID	Stratification Unit / Condition Class	Extent within development site	BAM plots required	Survey effort (number of Plots)	Patch Size (ha) (Number of patches)	Reference	
8	516_Derived Grassland	Moderate	2.76 ha	2	2	<5 ha (7)		
9	544_Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion	Low	0.83 ha	1	1	>1000 ha (1)	Table 4.6	
Predom	ninantly Cleared and Disturbed land			1				
NA	516_Disturbed Grassland	Other	7.39 ha	3	3	N/A		
10	594_Disturbed Grassland	Other	49.81 ha	4	8	N/A	Refer Section 4.2.1	
11	596_Disturbed Grassland	Other	50.59 ha	5	5	N/A		
12	Cleared Land / Dams	Other	0.48 ha	-	-	N/A		

Project No.: 0470861 18 May 2020 Page 39 www.erm.com Version: 2.0 Client: GAIA Australia

4.3.2 Vegetation Integrity Assessment Results


A total of 134 plant species were recorded within the 44 vegetation integrity survey plots (see Table H.6 in Appendix H). The field data sheets with results of the vegetation plots and photos of each plot are shown in Appendix G.


The plot data from the vegetation integrity survey plots were entered into the BAM calculator. The results of the vegetation integrity assessment are provided in


Table 4.6.

Table 4.6 Summary of Vegetation Integrity Scores

No	Vegetation Zone ID	Composition Score	Structure Score	Function Score	Vegetation Integrity Score
1	PCT 594_Moderate	52.9	43.6	99.8	61.3
2	PCT 594_Low	39.3	18.4	50.9	33.3
3	PCT 596_Moderate	67.4	52.3	69.4	62.6
4	PCT 596_Low	18.9	40.9	71.8	38.2
5	PCT 596_Very Low	18.4	22.7	31.6	23.6
6	PCT 596_Derived_Low	48.7	0.4	29.3	8
7	PCT 516_Very Low	31.6	11.2	24.9	20.7
8	PCT 516_Derived_Moderate	20	50.3	15	24.7
9	PCT 544_Low	26.7	24.3	65.9	35
10	516_Disturbed Grasslands	19.3	13.5	15	15.8
11	594_Disturbed Grasslands	25.7	2.4	26.2	11.8
12	596_Disturbed Grasslands	17.7	0.9	11	5.6







5. THREATENED SPECIES

5.1 Fauna Habitat Assessment

5.1.1 Important Habitat Mapping

For a small number of threatened species, a habitat constraint may refer to a mapped location that identifies areas that are considered important for those species. Maps currently available include important areas for the Swift Parrot (*Lathamus discolor*) and Regent Honeyeater (*Anthochaera phrygia*). These maps have been reviewed and the Subject Land does not form part of the important habitat mapping for either of these species.

5.1.2 Onsite Habitat Assessment

A habitat assessment was undertaken onsite seeking to identify the following fauna habitat features within the Subject Land:

- Habitat trees including hollow-bearing trees and nest trees;
- Foraging resources including fruiting and flowering plants and other feed tree species;
- Waterbodies including condition assessment;
- Rocks, ground litter and logs;
- Burrows, dens, nests and dreys; and
- Koala food trees.

Searches were also completed for indirect evidence of fauna habitat including scats, bones, tracks and trails.

The onsite habitat assessment identified that the majority of the Subject Land is highly disturbed, only supporting fauna species which are able to persist in highly modified agricultural landscapes and within woodland remnants. Additional habitat features assessed are provided in Table 5.1.

Table 5.1 Additional Habitat Features Considered

Habitat Features	Available with the Development Site								
Occurrences of Karst, Caves, Crevices and Cliffs	Not present within the Development Site								
Occurrences of Rock	A total of eleven rocky areas were recorded within the Subject Land (see Figure 5.b). Targeted reptile searches were undertaken in six of those rocky areas. Rocky areas were present on crests and hillsides as shown in Photograph 5-1 to Photograph 5-4 below.								
	Threatened species that could occur in these rocky outcrops are the:								
	Pink - tailed Legless Lizard (Aprasia parapulchella)								
	Rodd's Star Hair (Astrotricha roddii)								
	■ Border Thick-tailed Gecko (Uvidicolus sphyrurus)								
	As these species were not detected, the rocky outcrops were considered unlikely to provide habitat for threatened species.								

Habitat Features	Available with the Development Site								
Occurrences of Human made structures and Non- native Vegetation	The only human made structures present within the Subject Land are fences, gates and stockyards. No built structures, such as sheds or dwellings, are present.								
Termite/ ant mounds	A total of 39 ant/termite mounds were observed across the Subject Land as shown in Figure 5.b. An example of mounds found within the Subject Land are shown in Photograph 5-5 and Photograph 5-6 below.								
Hollows and Hollow Bearing Trees	A total of 60 hollow bearing trees were recorded across the Subject Land (see Table H.7 in Appendix H and Figure 5.b). Trees with hollows were eucalypt trees and stags (see Photograph 5-7 to Photograph 5-10). As shown in Graph 5-1 below, most hollows were medium size of less than or equal to 20cm diameter, followed by large hollows (greater than 20cm diameter) and small hollows (less or equal to 5cm diameter).								
	Hollow frequency Hollow frequency Hollow frequency The state of the								
	Graph 5-1 Tree hollow sizes and their frequency								
Bird nests	Numerous bird nest were observed across the Subject Land (see Photograph 5-11 to Photograph 5-14). Grey-crowned Babbler were observed actively adding sticks to one nest which indicates that the Subject Land is occupied by a breeding population of this species. Twenty-one nests were considered likely to be used by the Grey-crowned Babbler. The distribution of bird nests is shown in Figure 5.b and Figure 5.2.								
Koala Habitat Values	Only nine records of koala exists within the 10km locality as per BioNet atlas. The nearest records were recorded in 1999 at approximately 1.5km south from the Subje Land. Threatened species surveys indicated that no koalas, koala scats or evidence koala scratches were observed within the Subject Land. This is not unexpected due to the high level of clearance and disturbance at the Subject Land. Only one species listed as Koala Feed Tree (SEPP44), White Box (<i>Eucalyptus albert</i> was recorded in low numbers within the Subject Land.								



Photograph 5-1 View of rocky are on a hillside



Photograph 5-2 View of rocky area on a hillside



Photograph 5-3 View of rocky area on a hillside and crest



Photograph 5-4 View of rocky area on a hill crest



Photograph 5-5 Example of termite/ant mound



Photograph 5-6 Example of termite/ant mounds



Photograph 5-7 View of tree with hollow



Photograph 5-9 View of tree with hollows



Photograph 5-8 View of stag with hollow



Photograph 5-10 View of hollowed stag



Photograph 5-11 View of Grey-crowned Babbler nest



Photograph 5-12 Bird's nests on tree



Photograph 5-13 Tree with bird nest



Photograph 5-14 View of bird's nest on Eucalypt tree

5.2 Rivers, Streams and Wetlands

Riparian areas and farm dams were inspected along random meander transects in the assessment area surrounding the Subject Land.

5.2.1 Creek Lines

Third, second and first order un-named creek lines are present within Lot 2 in DP1039185 and are mapped tributaries of the Dumaresq River (see Figure 3.2). The creeks within the Project Boundary can be grouped as follows:

- Western Creek: This is a first order creek located within the Subject Land. This creek was dry during all the three survey periods (September 2018, December 2018 and March 2019) and provides very limited, if any aquatic habitat value.
- Central Creek and tributaries: This is a third order creek within the northern portion of the Project boundary and then diverges into a third order (eastern arm) and second order (western arm) to the south. These creeks were dry during all the three survey periods (September 2018, December 2018 and March 2019) and provides limited, seasonal aquatic habitat value only. Approximately six first order streams also converge into this central creek and provide very limited, if any aquatic habitat value.
- Eastern Creek: The southernmost portion of a first order stream is located on the eastern portion of the Project boundary (Lot 2 DP1039185) and outside of the Subject Land. This creek was not surveyed and will not be affected as part of the proposed development.

The overall condition of creek lines within the development footprint and the Project boundary is poor. Creek lines were dry during all three survey periods and erosion was noted along their banks (see **Photograph 5-15** to **Photograph 5-18**). Clearly defined river banks and channels are evident within the western, central and eastern creek lines indicating that at some point, these areas do contain large amounts of water and would provide some limited aquatic habitat for those species not dependent on a permanent water resource. The remaining 1st order creek lines within the Subject Land do not have conspicuous banks and bed and are not considered an aquatic habitat resource.

Riparian vegetation is mostly absent with scattered trees or patches of trees present. However, remnants of the Threatened Ecological Community – White Box Yellow Box Blakely's Red Gum Woodland were observed in several locations in association with stream banks and channels. Several vehicle crossing are present on creek lines within the Project Boundary.

It is worth noting that in natural conditions, when an aquatic species is present in a major river, that species is expected to also be distributed in the tributaries of that river because the river system is connected and water flows freely across the tributaries of the system. Where natural connectivity in riparian systems occur, naturally occurring restrictions to aquatic biodiversity are related to species-specific habitat and ecological requirements. Natural connectivity between the Dumaresq River and its tributaries at the Project Boundary no longer exists based on the presence of Bruxner Way, which runs in a general east-west direction along the northern boundary of the Project Boundary and beyond. It is worth noting that bridges were built on the highway at some of the major creeks and rivers intersections, such as Little Oaky Creek and Beardy River. This was not the case for tributaries crossing the Project Boundary.

Given that no connectivity exists between other tributaries of the Dumaresq River/Beardy River system and the creeks present at the Project Boundary, it is unlikely for those creeks to currently represent suitable habitat for aquatic biodiversity from the Dumaresq River.



Photograph 5-15 View of riparian area on the Project Boundary



Photograph 5-16 View of riparian area with vegetated bed.



Photograph 5-17 View of dry and eroded creek line with sparse trees on river bank



Photograph 5-18 View of creek near vehicle crossing

5.2.2 Farm Dams

A total of six farm dams are located within the Subject Land. Most of the farm dams were dry during surveys although they would provide a catchment for overland flow. As shown in photographs below, no fringing or submerged aquatic vegetation was present in these farm dams. An unidentified turtle was observed.



Photograph 5-19 View of a farm dam with water within the assessment area



Photograph 5-20 View of a small farm dam within the assessment area



Photograph 5-21 View of dry farm dam within the assessment area



Photograph 5-22 View of farm dam within the assessment area

5.3 Ecosystem Credit Species

A list of ecosystem credit species predicted to occur within the Subject Land, based on the PCTs present and generated by the calculator associated within the BAM (OEH 2017a) is provided in Table 5.2. The potential for these species to occur within the Subject Land was assessed in accordance with Section 6.2 of the BAM (OEH 2017a).

Two ecosystem credit species, the Glossy Black Cockatoo and White-bellied sea-eagle, were excluded from the assessment based on lack of foraging habitat across the Subject Land, all other species have been assumed to occur and contribute to ecosystem credits.

Table 5.2 Ecosystem Credit Species (BAM Calculator)

Ecosystem Credit Species	BC Act Listing	EPBC Act Listing	Justification for inclusion/exclusion
Anomalopus mackayi Five-clawed Work skink	E	V	Included. Potential habitat for the species is present within fallen logs and open paddocks with scattered trees.
Anthochaera phrygia Regent Honeyeater (Foraging)	CE	CE	Included. Potential foraging habitat is present. This woodland bird species forages in woodlands with significantly large numbers of mature trees, high canopy cover and abundance of mistletoes. No habitat is available within the disturbed grasslands.
Artamus cyanopterus cyanopterus Dusky Woodswallow	V	-	Included. Potential foraging habitat is present in most woodland vegetation zones, excluding grasslands, cleared and disturbed land.
Calyptorhynchus lathami Glossy Black-Cockatoo (Foraging)	V	-	Excluded as the Subject Land does not contain preferred foraging resources (<i>Allocasuarina</i> or Casuraina sp) for the species
Chalinolobus picatus Little Pied Bat	V		Included. The species' calls were recorded in the assessment area and hollow bearing trees are present across the Subject Land.
Chthonicola sagittata Speckled Warbler	V	-	Included. This species was recorded adjacent to the Subject Land during potential offset site investigations. Suitable habitat for the species is considered to occur in woodland PCTs. Grasslands and cleared land (disturbed grasslands) do not represent suitable habitat for the species.
Circus assimilis Spotted Harrier	V	-	Included. Potential habitat for the species occurs across all vegetation zones in the Subject Land.
Climacteris picumnus victoriae Brown Treecreeper (eastern subspecies)	V	-	Included. This species was recorded adjacent to the Subject Land during potential offset site investigations. Suitable habitat for the species is considered to occur in woodland PCTs. Grasslands and cleared land (disturbed grasslands) do not represent suitable habitat for the species.
Daphoenositta chrysoptera Varied Sittella	V	-	Included. Potential habitat for the species occurs in woodland vegetation within the Subject Land.
Glossopsitta pusilla Little Lorikeet	V	-	Included. Potential habitat for the species occurs in Eucalypt trees across all vegetation zones within the Subject Land.
Grantiella picta Painted Honeyeater	V	V	Included. Potential habitat for the species occurs in Box and Ironbark trees across the Subject Land.

Ecosystem Credit Species	BC Act Listing	EPBC Act Listing	Justification for inclusion/exclusion
Haliaeetus leucogaster White-bellied Sea-eagle (Foraging)	V	-	Excluded. Habitats are characterised by the presence of large areas of open water including larger rivers, swamps, lakes, and the sea. No foraging habitat available within the Subject Land.
Hieraaetus morphnoides Little Eagle (Foraging)	V	-	Included. Potential foraging habitat for the species occurs in woodland areas within the Subject Land.
Lathamus discolour Swift Parrot (Foraging)	E	CE	Included. Potential foraging habitat for the species occurs in Eucalypt trees in treed vegetation zones within the Subject Land. Derived grasslands and cleared (disturbed grasslands) excluded.
Lophoictinia isura Square-tailed Kite (Foraging)	V	-	Included. Potential foraging habitat for the species occurs over the canopy of woodlands along riparian zones within the Subject Land.
Macropus dorsalis (Black-striped Wallaby)	E		Included. Potential foraging habitat for the species occurs across the Subject Land.
Melanodryas cucullata cucullata Hooded Robin (south-eastern form)	V	-	Included. Potential foraging habitat is present in woodland vegetation zones within the Subject Land.
Melithreptus gularis gularis Black-chinned Honeyeater (eastern subspecies)	V	-	Included. Potential foraging habitat is present in woodland vegetation zones within the Subject Land.
Miniopterus orinae oceanensis Large Bent-winged bat (Foraging)	V	-	Included. The species' calls were recorded within the Subject Land. The species forages over the canopy of trees, so suitable habitat is present in woodland vegetation zones within the Subject Land.
Neophema pulchella Turquioise Parrot	V	-	Included. Suitable habitat for the species includes woodlands and native grasslands. Roosting and breeding habitat for this woodland bird are hollow-bearing trees in woodlands. It forages in woodlands and native grasslands.
Ninox connivens Barking Owl (Foraging)	V	-	Included. Potential foraging habitat is present in woodland vegetation zones within the Subject Land.
Nyctophilus corbeni Corben's Long-eared Bat	V	-	Included. The species' calls were recorded in the assessment area and hollow bearing trees are present across the Subject Land.
Petroica boodang Scarlet Robin	V	-	Included. Limited suitable habitat for the species occurs in woodlands with fallen logs and timber wtihtin the Subject Land.

Ecosystem Credit Species	BC Act Listing	EPBC Act Listing	Justification for inclusion/exclusion
Petroica phoenicea Flame Robin	V	-	Included. Limited potential habitat is present in woodland vegetation zones within the Subject Land.
Phascolarctos cinereus Koala (Foraging)	V	-	Included. A small number of suitable feeding trees (i.e. <i>E. albens</i>) were recorded within the Subject Land. Koalas, or evidence of Koala activity, were not recorded during surveys.
Pomatostomus temporalis temporalis Grey-crowned Babbler (eastern subspecies)	V	-	Included. Recorded on site. Foraging and breeding habitat for the species is present in woodland vegetation zones. Cleared land (i.e. disturbed grasslands) excluded.
Pteropus poliocephalus Grey-headed Flying-fox (Foraging)	V	V	Included. Potential foraging habitat is present in canopy of Eucalypt trees. Grasslands and cleared land excluded as potential foraging habitat.
Saccolaimus flaviventris Yellow-bellied Sheathtail-bat	V	-	Included. The species' calls were recorded in the assessment area and hollow bearing trees are present across the Subject Land.
Stagonopleura guttata Diamond Firetail	V	-	Included. Potential roosting and breeding habitat for the species occurs in woodlands, foraging habitat for the species occurs across the entire Subject Land.
Tyto novaehollandiae Masked Owl (Foraging)	V	-	Included. Potential foraging habitat is present in eucalypt forests/woodlands and along their edges within the Subject Land.

5.4 Species Credit Species

An assessment of habitat constraints for threatened species was undertaken in accordance with Step 2 of Section 6.4 of the BAM (OEH 2017a). For those threatened species predicted to occur and for which habitat constraints are listed, an assessment was undertaken of the presence of the habitat features within the development site.

The species generated by the calculator with habitat constraints, as well as the results of the habitat constraints assessment, are shown in Table 5.3.

Under Section 6.4.1.17 of the BAM, a species credit species can be considered unlikely to occur on a development site (or within specific vegetation zones) if following field assessment, it is determined that the habitat is substantially degraded such that the species is unlikely to utilise the development site (or specific vegetation zones).

A total of eight candidate species credit species were excluded as no suitable habitat is present within the Subject Land (see Table 5.3). The species excluded are: Regent Honeyeater (Breeding), Whitebellied Sea Eagle (Breeding), Swift Parrot (Breeding), Square-tailed Kite (Breeding), Grey-headed Flying Fox (Breeding), Eastern Bentwing Bat (Breeding), Koala (Breeding) and Ovenden's Ironbark.

Table 5.3 **Species Credit Species**

Species Credit Species	BC Act Listing	EPBC Act Listing	Habitat components and geographic restrictions	Sensitivity to gain class	Included or Excluded	Reason for inclusion or exclusion
Adelotus brevis Tusked Frog population in the Nandewar and New England Tableland Bioregion	E	-	-	Very High	Included	Six first order streams and several farm dams are present within the Subject Land although they do not provide any preferred habitat for this species. This species are usually found near creeks, ditches and ponds, and call while hidden amongst vegetation or debris. Limited riparian or and no fringing vegetation was noted in creeks and near the dams, respectively. Therefore, aquatic habitat is marginal at best
<i>Amalosia rhombifer</i> Zigzag Velvet Gecko	E	-	-	High	Included	Suitable habitat for the species occurs within the woodland habitats. This species is largely arboreal, living and foraging in trees.
Anthochaera phrygia Regent Honeyeater (Breeding)	CE	CE	-	High	Excluded	The Subject Land is not considered to be used as breeding or foraging habitat for the species. There are four known key breeding areas for this species, three of them in NSW - Capertee Valley, Bundarra-Barraba and Hunter Valley districts. For the Bundarra-Barraba area, three subsidiary foraging and breeding areas are known, Inverell-Ashford-Emmaville, Pilliga and Warrumbungles.
Astrotricha roddii Rodd's Star Hair	E	E	Rocky areas, Granite or acid volcanic outcrops	High	Included	A total of eleven rocky areas were recorded within the Subject Land and provide potential habitat for the species.
Burhinus grallarius Bush Stone-curlew	Е	-	Fallen/standing dead timber including logs	High	Included	This species inhabits open forests and woodlands with a sparse grassy groundlayer and fallen timber. Potential habitat is available within the grassy woodlands only. The heavily grazed areas do not constitute preferred habitat.

Species Credit Species BC Ac Listin		EPBC Act Listing	Habitat components and geographic restrictions	Sensitivity to gain class	Included or Excluded	Reason for inclusion or exclusion
Calyptorhynchus lathami Glossy Black-Cockatoo (Breeding)	V	-	-	High	Included	Suitably sized hollows have been recorded within the Subject Land although no preferred foraging habitat has been recorded which would limit the likelihood of this species nesting on site.
Dichanthium setosum Bluegrass	V	V	-	High	Included	The species has potential to occur in the Subject Land.
Digitaria porrecta Finger panic Grass	Е	-	-	Moderate	Included	The species has potential to occur in the Subject Land.
Eucalyptus caleyi subsp. ovendenii Ovenden's Ironbark	V	V	-	High	Excluded	The preferred altitudes for the species are between 610 and 820 m, on granitic substrates. The Subject Land in located at much lower elevations between 335 and 420 m.
Geophaps scripta scripta Squatter Pigeon (southern)	CE	V	-	High	Included	The species has potential to occur within the woodland habitats although the species prefers sandy areas and usually close to water (within 3 km of water bodies or courses).
Haliaeetus leucogaster White-bellied Sea-Eagle (Breeding)	V	-	-	High	Excluded	Breeding habitat for this species consists of mature tall open forest, open forest, tall woodland, and swamp sclerophyll forest close to foraging habitat (not available within the Subject Land). Nest trees are typically large emergent eucalypts and often have emergent dead branches or large dead trees nearby which are used as 'guard roosts'.
Hieraaetus morphnoides Little Eagle (Breeding)	V	-	-	Moderate	Included	Potential breeding habitat is available within the woodland habitats. This species nests in tall living trees within remnant vegetation where pairs build a large stick nest in winter.

Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page 56 www.erm.com Version: 2.0

Species Credit Species	BC Act Listing	EPBC Act Listing	Habitat components and geographic restrictions	Sensitivity to gain class	Included or Excluded	Reason for inclusion or exclusion		
Homopholis belsonii Belson's Panic	Е	V	-	NA	Included	Grows in dry woodland (e.g. Belah) often on poor soils, although sometimes found in basalt-enriched sites north of Warialda and in alluvial clay soils. The species has potential to occur in the Subject Land		
Hoplocephalus bitorquatus Pale-headed Snake				High	Included	Suitable habitat for the species occurs within the Subject Land. A small number of historical records are known for the New England Tableland from Glenr Innes and Tenterfield; however, the majority of records appear to be from sites of lower elevation. The species is found mainly in dry eucalypt forests and woodlands, cypress forest and occasionally in rainforest or moist eucalypt forest. It has limited potential to occur in the Subject Land		
<i>Indigofera baileyi</i> Bailey's Indigo	E	-	-	High	Included	The species is known from around Inverell and Ashford in the western inland slopes. It prefers open woodlands on loam and clay loam soils, typically from granite or basalt, and has potential to occur in the Subject Land		
Lathamus discolour Swift Parrot (Breeding)	E	CE	-	Moderate	Excluded	The species breeds in Tasmania during spring and summer, migrating in the autumn and winter months to south-eastern Australia.		

Species Credit Species	BC Act Listing	EPBC Act Listing	Habitat components and geographic restrictions	Sensitivity to gain class	Included or Excluded	Reason for inclusion or exclusion
Lophoictinia isura Square-tailed Kite (Breeding)	V	-	-	Moderate	Excluded	No suitable breeding habitat for the species is present within the Subject Land, as the species prefers coastal and subcostal environments and inland wooded watercourses. Second and third order streams are excluded from the Subject Land. Scattered records of the species throughout NSW indicate that the species is a regular resident in the north, north-east and along the major west-flowing river systems. Breeding is from July to February, with nest sites generally located along or near watercourses, in a fork or on large horizontal limbs.
Miniopterus orinae oceanensis Large Bent-winged bat (Breeding)	V	-	-	Very High	Excluded	Caves are the breeding habitat for the species. Caves, suitable for maternity areas, are not present within the Subject Land. Being a dual credit species, foraging habitat resources are already accounted for in the ecosystem credits.
Ninox connivens Barking Owl (Breeding)	V	-	Hollow bearing trees. Living or dead trees with hollows greater than 20cm diameter and greater than 4m above the ground	High	Included	Hollow bearing trees occur within woodland habitats and also as isolated trees within highly disturbed vegetation zones on the Subject Land. However, only a small number of these trees are likely to represent suitable breeding habitat for this species due to limiting factors associated with hollow size and height above ground. Potential breeding habitat is considered to comprise Vegetation Zones 594_Moderate, 594_Low, 596_Moderate, 544_Low and 516_Very low given that hollow-bearing trees (with suitable hollows) were observed to be concentrated within these zones.

Species Credit Species	BC Act Listing	EPBC Act Listing	Habitat components and geographic restrictions	Sensitivity to gain class	Included or Excluded	Reason for inclusion or exclusion
Petaurus norfolcensis Squirrel Glider	V	-	-	High	Included	Substandard suitable habitat is present in Eucalypt trees in the Subject Land.
Phascolarctos cinereus Koala (Breeding)	V	V	-	High	Excluded	Breeding habitat is not considered to be present on the Subject Land. This determination has been made on the basis that no koalas, or evidence of koala activity, has been recorded on the Subject Land and given that recognised feed trees occur only in very low numbers onsite.
Polygala linariifolia Native Milkwort	Е	-	-	High	Included	The species has potential to occur in the Subject Land
Pomaderris queenslandica Scant Pomaderris	Е	-	-	High	Included	The species has potential to occur in the Subject Land
Pteropus poliocephalus Grey-headed Flying-fox (Breeding)	V	V	-	High	Excluded	No known breeding camp is present within the Subject Land.
Swainsona sericea Silky Swainson-pea	V	-	-	High	Included	The species has the potential to occur within the Subject Land.
Thesium austral Austral Toadflax	V	V	-	Moderate	Included	The species has potential to occur in the Subject Land

Species Credit Species	BC Act Listing	EPBC Act Listing	Habitat components and geographic restrictions	Sensitivity to gain class	Included or Excluded	Reason for inclusion or exclusion
Tyto novaehollandiae Masked Owl (Breeding)	V	-	-	High	Included	The Masked Owl is a large forest owl and prefers uncleared or lightly cleared areas with high densities of old hollow bearing trees. Given that habitat is present on site that meets the breeding habitat constraint for this species (living or dead trees with hollows greater than 20cm diameter), breeding habitat has been assumed to be present for this species. Potential breeding habitat is assumed to occur within Vegetation Zones 594_Moderate, 594_Low, 596_Moderate, 544_Low and 516_Very low given that hollow-bearing trees (with suitable hollows) were observed to be concentrated within these zones. However, it is considered unlikely that the Masked Owl would use these habitat features given the highly disturbed and highly fragmented context in which theses habitat features occur.
Uvidicolus sphyrurus Border Thick-tailed Gecko	V	V	-	High	Included	The species has potential to occur in the Subject Land

5.5 Additional Species Considered

Based on results of the BioNet Atlas search (see Appendix B), the PMST (see Appendix C) and the likelihood of occurrence assessment undertaken during desktop review (see Appendix D), the following species were also considered for assessment:

- Eastern False Pipistrelle (Falsistrellus tasmaniensis);
- Bristle-faced Free-tailed Bat (Setirostris eleryi previously Mormopterus eleryi);
- Eastern Cave Bat (Vespadeuls vulturnus);
- Greater Broad-nosed Bat (Scoteanax rueppellii);
- Narrow-leaved Black Peppermint (Eucalyptus nicholii); and
- Heath Wrinklewort (Rutidosis heterogama).

5.6 Candidate Species Surveyed

The species listed in Table 5.4 are those that are considered to have habitats present at the Subject Land. Surveys have been conducted for these species. The results are summarised in Table 5.5.

Details of the survey methodologies for each surveyed species are provided in Appendix F and Appendix H. Targeted survey locations are mapped on Figure 5.1a.

Species polygons have been defined for relevant species credit species that have been confirmed, or are assumed, to be present on the Subject Land as mapped on Figure 5.3.

Table 5.4 **Candidate Species Surveyed**

Species Credit Species	BC Act Listing	EPBC Act Listing	Biodiversity Risk	Biodiversity Risk Weighting	Survey Time	Assumed to occur/survey/ expert report	Recorded during field surveys	Species polygon area or count
Adelotus brevis Tusked Frog population in the Nandewar and New England Tableland Bioregion	E	-	Very High	3	December	No	No	No
Amalosia rhombifer Zigzag Velvet Gecko	Е	-	High	2	December	No	No	No
Anthochaera phrygia Regent Honeyeater	CE	CE	Very High	3	December	No	No	No
Astrotricha roddii Rodd's Star Hair	E	Е	High	2	September December	No	No	No
Burhinus grallarius Bush Stone-curlew	Е	-	High	2	December	No	No	No
Calyptorhynchus lathami Glossy Black-Cockatoo	V	-	High	2	No	No	No	No
Dichantium setosum Bluegrass	V	V	High	2	December March	No	No	No
Digitaria porrecta Finger Panic Grass	Е	-	High	2	December March	No	No	No
Geophaps scripta scripta Squatter Pigeon (southern)	CE	V	Very High	3	December	No	No	No
Hieraaetus morphnoides Little Eagle	V	-	Moderate	1.5	December	No	No	No
Homopholis belsonii Belson's Panic	Е	V	N/A	1	December March	No	No	No

Species Credit Species	Species Credit Species BC Act Listing Listin		Biodiversity Risk	Biodiversity Risk Weighting	Survey Time	Assumed to occur/survey/ expert report	Recorded during field surveys	Species polygon area or count	
Hoplocephalus bitorquatus Pale-headed Snake	V	-	High	2	December	No	No	No	
<i>Indigofera baileyi</i> Bailey's Indigo	E	-	High	2	September December	No	No	No	
Lophoictinia isura Square-tailed Kite	V	-	Moderate	1.5	December	No	No	No	
Mormopterus eleryi (syn. Setirostris eleryi) Bristle-faced Free-tailed Bat	Е	-	High	2.0	December	Survey	Yes	Yes	
Miniopterus orianae oceanensis Large Bent-winged bat	V	-	Very High	3.0	December	Survey	Yes	No	
Ninox connivens Barking Owl	V	-	High	2	None	Assumed Present	N/A	Yes	
Petaurus norfolcensis Squirrel Glider	V	-	High	2	December	No	No	No	
Phascolarctos cinereus Koala	V	V	High	2	December	No	No	No	
Polygala linariifolia Native Milkwort	Е	-	High	2	December January	No	No	No	
Pomaderris queenslandica Scant Pomaderris	Е	-	High	2	September December March	No	No	No	
Pteropus poliocephalus Grey-headed Flying-fox	V	V	High	2	December	No	No	No	
Swainsona sericea Silky Swainson-pea	V	-	High	2	September December	No	No	No	

Project No.: 0470861 18 May 2020 www.erm.com Version: 2.0 Client: GAIA Australia Page 63

Species Credit Species	BC Act Listing	EPBC Act Listing	Biodiversity Risk	Biodiversity Risk Weighting	Survey Time	Assumed to occur/survey/ expert report	Recorded during field surveys	Species polygon area or count
Thesium austral Austral Toadflax	V	V	Moderate	1.5	September December	No	No	No
<i>Tyto novaehollandiae</i> Masked Owl	V	-	High	2	None	Assumed Present	N/A	Yes
Uvidicolus sphyrurus Border Thick-tailed Gecko	V	V	High	2	December	No	No	No
Vespadeuls vulturnus Eastern Cave Bat	V	-	Very High	3.0	December	Survey	Yes	Yes

5.7 Targeted Survey Results

Targeted surveys for flora and fauna species were completed across the Subject Land. A description of survey methodology and effort is provided in Appendix F. Figure 5.1a shows the location of survey tracks across the Subject Land.

5.7.1 Flora Species Recorded

No threatened flora species were recorded during site surveys. A complete list of flora species recorded on the Subject Land is provided in Table H.6 (Appendix H).

A total of 143 flora species in 47 families were recorded within the assessment area. This included a total of 111 native (78 %) and 32 exotic (22 %) species. The most numerous families were Poaceae (30 species), Asteraceae (21 species) and Fabaceae (Faboideae) (10 species).

Exotic species included two HTE and WoNS, Tiger Pear (*Opuntia aurantiaca*) and Velvet Tree Pear (*Opuntia tomentosa*). Due to the high level of grazing and lack of reproductive material, identification of some flora specimens was only viable to genus level, including *Senecio* sp., *Hypericum* sp. and *Solanum* sp. Therefore, presence of an additional WoNS, Fireweed (*Senecio madagascariensis*), cannot be precluded. Fireweed is also listed as a Priority Weed in the Northern Tablelands. Similarly, exotic species within genera *Hypericum* and *Solanum* were recorded. Therefore, potential presence of priority weeds St John's Wort (*Hypericum perforatum*) and Silverleaf Nightshade (*Solanum elaeagnifolium*), cannot be precluded. As per the Biosecurity Act, all exotic species require management and preparation of a Weed Management Plant for the proposed solar farm will be required.

5.7.2 Fauna Species Recorded

A total of 78 fauna species were recorded on and adjacent to the Subject Land. A summary of faunal groups recorded is presented in Table 5.5 and a complete fauna species list is provided in Appendix H.

A total of eleven threatened species were recorded within the Subject Land (see Table 5.5 below). The species included eight vulnerable microchiropteran bats (Little Pied Bat, Eastern Bentwing-bat, Corben's Long-eared Bat, Yellow-bellied Sheath-tailed Bat, Eastern Cave Bat, Eastern False Pipistrelle, Greater Broad-nosed Bat and Hoary Wattled Bat), one endangered microchiropteran bat (Bristle-faced Free-tailed bat), one vulnerable bird (Grey-crowned Babbler) and one migratory bird (Cicadabird). Out of the eight Vulnerable microchiropteran bats, six were definite call identifications and two (Greater Broad-nosed Bat and Hoary Wattled Bat) were potential calls. Two additional Vulnerable bird species (Brown Treecreeper and Speckled Warbler) were also recorded adjacent to the Subject Land during potential offset site investigations.

Table 5.5 Summary of Fauna Species Recorded

Fauna Type	Number of Species	BC Act	EPBC Act		
Woodland Birds	46 on the Subject Land and an additional 7 adjacent to the Subject Land	One species, the Grey-crowned Babbler is listed as Vulnerable and has been further assessed in Section 5.7.3 Two additional Vulnerable species were recorded adjacent to the Subject Land during potential offset site investigations – Brown Treecreeper (Climacteris picmnus victoriae) and Speckled Warbler (Chthonicola sagittata)	One species, the Cicadabird is listed as Marine and has been further assessed in Section 5.7.6		
Raptors	2 flying over Subject Land	No	No		
Forest Owls	0	No	No		
Reptiles	9	No	No		
Amphibians	3	No	No		
Terrestrial Mammals	11 including 5 introduced species	No	No		
Arboreal Mammals	1	No	No		
Microchiropteran Bats	12 definite + 6 potential	Six Vulnerable species: Little Pied Bat, Eastern Bentwing-bat, Corben's Long-eared Bat, Yellow-bellied Sheath-tailed Bat, Eastern Cave Bat (this species has been further assessed in Section 5.7.4) and Eastern False Pipistrelle Two potential Vulnerable species: Greater Broad-nosed Bat and Hoary Wattled Bat. One Endangered species: Bristle-faced Free-tailed Bat. This species has been further assessed in Section 5.7.4.	One vulnerable species: Corben's Long-eared Bat		

Results of each threatened species recorded on the Subject Land are summarised in sub-sections below.

5.7.3 Grey-crowned Babbler

The Grey-crowned Babbler (*Pomatostomus temporalis*) is listed as an ecosystem credit species.

It inhabits open Box-Gum Woodlands on the slopes, and Box-Cypress-pine and open Box Woodlands on alluvial plains. Woodlands on fertile soils in coastal regions. Its flight is laborious so birds prefer to hop to the top of a tree and glide down to the next one. Birds are generally unable to cross large open areas. It lives in family groups that consist of a breeding pair and young from previous breeding seasons. A group may consist of up to fifteen birds. All members of the family group remain close to each other when foraging. It is insectivorous and it forages on the trunks and branches of eucalypts and other woodland trees or on the ground, digging and probing amongst litter and tussock grasses. It builds nests that are used as dormitory and roosting and uses them all year round. It breeds between July and February. Territory ranges from one to 50 hectares (usually ten hectares) and are defended all year (OEH 2019).

Grey-crowned Babblers are obligate communal breeders that form a family group, in which offspring from the previous season and other unrelated birds help to raise the current's year's brood. Young birds stay with the family group for at least one year after fledging and may remain for two or more years acting as non-breeding helpers. As breeding spaces become available in the population, some helpers may disperse to establish their own breeding group. In some populations, breeding success is related to the number of helpers. Population viability studies in Victoria suggests that a viable population is likely to contain more than ten family groups, while populations with less than ten family groups are likely to have high rate of extinction. In NSW, the species breeds between July and February (OEH 2019).

It has been suggested that cooperative breeder species, such as the Grey-crowned Babbler, are more sensitive to habitat fragmentation and loss (including loss due to fire) as availability of resources for breeding decreases (Fischer 2011). Habitat fragmentation and predation by introduced species being the major threats to the species. The species is reported to survive in disturbed landscapes, such as urban areas, where proportion of ground cover and leaf litter provides sufficient food (Lambert and Ford 2016, Stevens *et. al.* 2015).

The Grey-crowned Babbler was observed during surveys in September and December 2018, heard during the March 2019 survey and observed more recently during surveys in November 2019 and January 2020. In December 2018, the Grey-crowned Babbler was recorded at 20 points within the Subject Land and a total of 21 bird nests likely to belong to the species were observed (see Figure 5.2). The records included up to 15 individuals observed, 21 calls heard with an additional two potential calls heard at distance. Also, a pair was observed adding twigs to a nest located on the lower branches of a Silver-leaved Ironbark (*Eucalyptus melanophloia*).

The Grey-crowned Babbler was observed in the following vegetation zones:

- 594_Moderate (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion). A total of 4.1 ha of this vegetation zone is present across the Subject Land;
- 596_Moderate (Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion). A total of 11.15 ha of this vegetation zone is present across the Subject Land;
- 544_ Low (Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion). A total 0.83 ha are present within the Subject Land;
- 516_Very Low (Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion), a total of 3.00 ha of this vegetation zone is present across the Subject Land; and
- 516_Disturbed Grasslands (Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion). A total 7.39 ha of this vegetation zone is present within the Subject Land.

A Test of Significance for the species was undertaken and is included in Appendix E.

5.7.4 Microchiropteran Bats

A total of nine threatened microchiropteran bat species were recorded within the assessment area, including eight vulnerable species and one endangered species. In accordance with the 'Species Credit' threatened bats and their habitats guide (OEH 2018b), the species fall into the following credit species types:

- Ecosystem Credit Species:
 - Little Pied Bat (Chalinolobus picatus) (V)
 - Large Bent-winged Bat (Miniopterus orianae oceanensis) (V) Foraging Habitat only
 - Corben's Long-eared Bat (Nyctophilus corbeni) (V)
 - Eastern False Pipistrelle (Falsistrellus tasmaniensis) (V)
 - Greater Broad-nosed Bat (Scoteanax rueppellii) (V)
 - Hoary Wattled Bat (Chalinolobus nigrogriseus) (V)
 - Yellow-bellied Sheath-tailed Bat (Saccolaimus flaviventris) (V)
- Species Credit Species:
 - Bristle-faced Fee-tailed Bat (Mormopterus eleryi) (E)
 - Eastern Cave Bat (Vespadeuls vulturnus) (V)
 - Large Bent-winged Bat (*Miniopterus orianae oceanensis*) (V) Breeding habitat only listed as SCS

Habitat polygons for Species Credit Species (SCS) have been assessed in accordance with Step 5 of Section 6.4 of the BAM (OEH 2017a and OEH 2018a). Based on the analysis of habitat availability for SCS within the Subject Land (see Table 5.6), a species polygon has been prepared for the Eastern Cave Bat and the Bristle-faced Free-tailed Bat.

Suitable breeding habitat is not considered to occur on the Subject Land for the Eastern Bentwing Bat or Eastern Cave Bat.

Table 5.6 Habitat Summary for Microchiropteran Bat Species Credit Species

Scientific Name BC Common Name Act Listin			Habitat as	Habitat Constraints		PCTs	Comments			
	Act Listing	Roosting	Breeding	Foraging	Are paddock trees important habitat?	Patch Size	Description	Present		
Mormopterus eleryi (syn. Setirostris eleryi) Bristle-faced Free- tailed Bat	E	Hollows and tree fissures. Habitat is within 500m of riparian areas (including dry river/creek beds) and water bodies within the PCTs associated with the species.	Tree hollows (Ellis 2006)	Insectivorous species that forage typically well-above the canopy and only coming low in relatively open areas (Ellis and Wilson 1992)	No	5-24 ha	Land within 500 m of watercourses or dams surrounded by eucalypts containing hollows.	Yes	516, 544, 594, 596	Hollow bearing trees within 500m of watercourses and dams are the relevant habitat component for the species.
Vespadelus troughtoni Eastern Cave Bat	V	Caves, Scarps, cliffs, disused mine workings.	Potential breeding habitat is the PCTs associated with the species (as per the TBDC) within 100m of rocky areas, caves, overhangs crevices, cliffs and escarpments, or old mines or tunnels, old buildings and sheds within potential habitat.	Found in dry open forests and woodland, near cliffs or rocky overhangs.	No	5-24 ha	Caves within two kilometres of rocky areas containing caves, overhangs, escarpments, outcrops, crevices or boulder piles, or within two kilometres of old mines, tunnels, old buildings or sheds.	No	516, 544, 594, 596	Trees and treed areas represent potential foraging habitat for the species within the Subject Land. No roosting or breeding habitat is present.
Miniopterus orianae oceanensis Large Bent-winged Bat	V	Caves are the primary roosting habitat, but also use derelict mines, storm-water tunnels, buildings and other man-made structures	Maternity caves. Potential breeding habitat is caves, tunnels, mines or other structures known or suspected to be used by the species	Being an insectivorous species it is assumed foraging habitat is treed areas. Within 300km range of maternity caves.	No	<5 ha	Breeding: Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding including species records with microhabitat code "IC - in cave;" observation type code "E nest-roost;" with numbers of individuals >500	No	516, 544, 594, 596	Trees and treed areas represent potential foraging habitat for the species within the Subject Land. No roosting or breeding habitat is present.

5.7.4.1 Bristle-faced Free-tailed Bat

As described in Table 5.6, the Bristle-faced Free-tailed Bat roosts in tree hollows and tree fissures, breeds in tree hollows and forages for insects over the canopy of trees (Ellis 2001, NSW SC 2004, OEH 2019). In defining the habitat polygon for the Bristle-faced Free-tailed Bat, ecological information from the TBDC and scientific literature was taken into account including the following from BioNet:

This species may move distances greater than 10km, dispersal distance of 100m - 10km was selected as this is the distance at which females regularly move to establish new colonies in available habitat. Additionally, selected <1 for average number of offspring because females do not give birth every (often miscarry etc).

Potential habitat is riparian areas (including dry river/creek beds) within the PCTs associated with the species. Survey should sample the available range of suitable vegetation along riparian areas on the Subject Land. Traps or nets should be set near water holes (especially if isolated), under/beside large trees, in/beside creek beds, or in 'flyway' spaces between vegetation. NB. Use of acoustic detection alone is not suitable for this species as the call is difficult to distinguish from other common species. Refer to Threatened Bat Survey Guide.

All habitat on the Subject Land where the Subject Land is within 500m of a river, creek or riparian area must be mapped. Use aerial imagery to map river, creek or riparian areas (including dry creek channels, former creek channels, billabongs etc.) on or within 500m of the Subject Land. Species polygon boundaries should align with PCTs on the Subject Land to which the species is associated that are within 500m of waterbodies mapped.

In accordance with Table 1 of the OEH (2018b) 'Species Credit' threatened bats and their habitats guide, the species polygon for the Bristle-faced Free-tailed Bat has been prepared and includes the following features:

 All habitat on the Subject Land where the Subject Land is within 500m of a river, creek or riparian area.

A review of information on the Bristle-faced Free-tailed Bat has been undertaken and is presented in Appendix E. It is acknowledged that little information is available regarding the ecology and biology of this species. It is assumed that the species roosts and breeds on tree hollows and tree fissures. Based on feeding behaviour of species within the *Mormopterus / Setirostris* genus, it is assumed that the species is aerial insectivorous and that is a vegetation dependent species which forage typically well above the canopy of any vegetation type when feeding. Therefore, suitable roosting, breeding and foraging habitat for the species within the Subject Land includes individual trees (particularly hollow-bearing trees, trees with fissures and tree canopies) and woodland areas within 500m of riparian areas. Cleared areas are not considered to represent suitable habitat for the species.

A species habitat polygon for the Bristle-faced Free-tailed Bat is provided in Figure 5.3.

5.7.4.2 Eastern Cave Bat

As described in Table 5.6, the Eastern Cave Bat roosts in Caves, Scarps, cliffs, disused mine workings. Potential breeding habitat is the PCTs associated with the species (as per the TBDC) within 100m of rocky areas, caves, overhangs crevices, cliffs and escarpments, or old mines or tunnels, old buildings and sheds within potential habitat.

The BioNet Atlas indicates that the Eastern Cave Bat:

"Any impacts on breeding habitat could be considered potentially serious and irreversible. This species is retained as dual credit because foraging habitat is broad ranging but breeding habitat is highly specific. At lower altitudes this species is usually more abundant during winter months, the lower numbers of individuals from October to February are due to females moving to maternity sites. Additionally, selected <1 for average number of offspring because females do not give birth every (often miscarry etc)."</p>

"All breeding habitat including the cave, or other features, used for breeding and the area immediately surrounding this feature must be mapped. Species polygon boundaries should have a 100m radius buffer around an accurate GPS point location centred on the cave/feature entrance."

In accordance with Table 1 of the OEH (2018b) 'Species Credit' threatened bats and their habitats guide, a species polygon for the Eastern Cave Bat would need to be prepared that includes the following features:

- "All habitat on the Subject Land where the Subject Land is within 2km of caves, scarps, cliffs, rock overhangs and disused mines."
- "Note: any breeding habitat identified for this species (see Table 2) is a potential serious and irreversible impact."

In accordance with Table 2 of the OEH (2018b), the features and approach required to develop the species polygon for the Eastern Cave Bat, a species that require identification of breeding habitat are:

- "All breeding habitat on or within 100m of the Subject Land and the area immediately surrounding the feature. Artificial structures should be inspected and included if the species is using these features for breeding (see Section 3.2)."
- "Note all habitat for this species should also be mapped if present (i.e. including that described in Table 1)."

Roosting and/or breeding habitat for the Eastern Cave Bat does not occur on the Subject Land. Aerial images covering an area of 2km surrounding the Project Boundary were assessed for the potential presence of caves, cliffs, rock overhangs and disused mines. No suitable breeding and/or roosting habitat features were identified. However, rocky areas containing boulder piles and crevices were observed within 2km of the Subject Land.

It is considered that only foraging resources for the Eastern Cave Bat are available on the Subject Land. In accordance with the TBDC, the four PCTs identified at the Subject Land (i.e. PCT 516, 544, 594 and 596) represent potential foraging habitat. The profile for the species (OEH 2019) indicates that "little is understood of its feeding or breeding requirements or behaviour". All vegetation zones, with the exception of disturbed and derived grasslands, have therefore been considered as potential foraging habitat for the Eastern Cave Bat. A species habitat polygon for the Eastern Cave Bat is provided in Figure 5.3.

A test of significance for the species is presented in Appendix E. Given the absence of roosting and/or breeding habitat for the species at the Subject Land, it is considered that the proposed development does not represent a significant impact on this species.

5.7.5 Owls

5.7.5.1 Barking Owl

The TBDC provides the following information regarding the preferred habitat of the Barking Owl:

- Inhabits woodland and open forest, including fragmented remnants and partly cleared farmland. It is flexible in its habitat use, and hunting can extend into closed forest and more open areas. Sometimes able to successfully breed along timbered watercourses in heavily cleared habitats (e.g. western NSW) due to the higher density of prey found on these fertile riparian soils.
- Roosts in shaded portions of tree canopies, including tall mid storey trees with dense foliage such as Acacia and Casuarina species. During nesting season, the male perches in a nearby tree overlooking the hollow entrance.
- Preferentially hunts small arboreal mammals such as Squirrel Gliders and Common Ringtail
 Possums, but when loss of tree hollows decreases these prey populations the owl becomes more

reliant on birds, invertebrates and terrestrial mammals such as rodents and rabbits. Can catch bats and moths on the wing, but typically hunts by sallying from a tall perch.

- Requires very large permanent territories in most habitats due to sparse prey densities.
 Monogamous pairs hunt over as much as 6000 hectares, with 2000 hectares being more typical in NSW habitats.
- Two or three eggs are laid in hollows of large, old trees. Living eucalypts are preferred though dead trees are also used. Nest sites are used repeatedly over years by a pair, but they may switch sites if disturbed by predators (e.g. goannas).
- Nesting occurs during mid-winter and spring, being variable between pairs and among years. As a rule of thumb, laying occurs during August and fledging in November. The female incubates for 5 weeks, roosts outside the hollow when chicks are 4 weeks old, then fledging occurs 2-3 weeks later. Young are dependent on their parents for several months.

The TBDC also provides that where a breeding site has been identified in accordance with the BAM the species polygon should be established by providing a circular buffer with a 100m radius around the nest tree. This includes suitable habitat for male roosts, feeding/grooming perches and fledgling requirements. It does not account for foraging habitat. The shape of the buffer can be modified where evidence indicates an alternative shape would better meet the species needs in the context of the assessment site.

As no targeted surveys (e.g. call playback and identification of nest trees) were completed for this dual credit species, breeding habitat has been assumed to be present by identifying suitable habitat. As described in Table 5.3, a species polygon has been mapped for this species by including entire vegetation zones in which hollow-bearing trees were concentrated and contained hollows of a suitable size (i.e. >20cm wide) and height above the ground (i.e. >4m) in accordance with breeding habitat constraints provided in BAM-C.

Potential breeding habitat is therefore assumed to occur within the following Vegetation Zones as shown in Figure 5.3:

- PCT 594_Moderate;
- PCT 594_Low;
- PCT 596_Moderate;
- PCT 544 Low; and
- PCT 516_Very low.

Given the approach described above, the species polygon for Barking Owl is considered to represent a conservative overestimate of actual/potential breeding habitat in that only a small proportion of hollow-bearing trees within the polygon are suitable for nesting.

5.7.5.2 Masked Owl

The TBDC provides the following information regarding the preferred habitat of the Masked Owl:

- Lives in dry eucalypt forests and woodlands from sea level to 1100 m.
- A forest owl, but often hunts along the edges of forests, including roadsides.
- The typical diet consists of tree-dwelling and ground mammals, especially rats.
- Pairs have a large home-range of 500 to 1000 hectares.
- Roosts and breeds in moist eucalypt forested gullies, using large tree hollows or sometimes caves for nesting. Although there is no evidence to suggest that caves are utilised for nesting in NSW.
- Dead stags are especially popular for roosting/breeding habitat.

The TBDC also provides that where a breeding site has been identified in accordance with the BAM the species polygon should be established by providing a circular buffer with a 100m radius around the nest tree. This includes suitable habitat for male roosts, feeding/grooming perches and fledgling requirements. It does not account for foraging habitat. The shape of the buffer can be modified where evidence indicates an alternative shape would better meet the species needs in the context of the assessment site.

As no targeted surveys (e.g. call playback and identification of nest trees) were completed for this dual credit species, breeding habitat has been assumed to be present by identifying suitable habitat. Although no breeding habitat constraints are explicitly identified in BAM-C for this species, the Approved Recovery Plan for Large Forest Owls (DEC 2006) states that the Masked Owl nests in hollows greater than 40cm wide and greater than 100cm deep, with hollow entrances at least 3m above ground, in trees of at least 90cm diameter at breast height. As described in Table 5.3, a species polygon has been mapped for this species by including entire vegetation zones in which hollow-bearing trees were concentrated and contained hollows of a suitable size and height above ground.

Potential breeding habitat is therefore assumed to occur within the following Vegetation Zones as shown in Figure 5.3:

- PCT 594_Moderate;
- PCT 594 Low;
- PCT 596 Moderate;
- PCT 544_Low; and
- PCT 516_Very low.

Given the approach described above, the species polygon for Masked Owl is considered to represent a conservative overestimate of actual/potential breeding habitat, particularly given that this species is generally known to roost and breed in more forested habitats in NSW. The Approved Recovery Plan for Large Forest Owls (DEC 2006) also provides that the Masked Owl prefers unlogged or lightly logged forest with high densities of old hollow trees and avoids young regrowth.

5.7.6 Cicadabird

One individual of the Cicadabird (*Coracina tenuirostris*) was observed within the Subject Land during surveys. A significant impact assessment for the species was undertaken (see Appendix E) and it was concluded that significant impacts on this migratory/marine species are unlikely to result due to the proposed solar farm development. The Cicadabird is not further assessed.

5.7.7 Threatened Aquatic Species

No threatened aquatic species are considered likely to occur within the Development Site. Threatened species maps prepared by the NSW Department of Primary Industries were reviewed (https://www.dpi.nsw.gov.au/fishing/species-protection/threatened-species-distributions-in-nsw/freshwater-threatened-species-distribution-maps) via the SEED Portal. The following threatened species listed under the NSW Fisheries Management Act have been considered:

- Southern Purple Spotted Gudgeon (Mogurnda adspersa): The Subject Land is located within the indicative distribution area of the Southern Purple Spotted Gudgeon, which is listed as Endangered under the FM Act. DPI's indicative distribution map indicates that none of the creeks within the Subject Land are mapped as habitat for the species (see Figure 5.4)The nearest creeks mapped as habitat for the Southern Purple Spotted Gudgeon are:
 - Dumaresq River: it is located at approximately 1.2 km north from the Development Site. This river runs in a general east-west direction.

- Little Oaky Creek: it is located at approximately 1.7 km west from the Development Site. Little Oaky Creek is a tributary of Dumaresq River.
- Crooked Creek: it is located at approximately 4.5 km west from the Development Site and
 2.5 km west from Little Oaky Creek. Crooked Creek is a tributary of Dumaresq River.
- Beardy River: is located to the east from the Development Site. The nearest point is located at approximately 3 km from the development footprint. Beardy River runs on a general north-south direction from its convergence with Dumaresq River. None of the first order tributaries of Beardy River are located within the Development Site.

All creeks within the Development Site were dry during the three survey periods in September 2018, December 2019 and March 2019 and have low habitat value for the Southern Purple Spotted Gudgeon.

- Eel Tailed Catfish (*Tandanus tandanus*): The Subject Land is located within the indicative distribution area of the Murray-Darling Basin population of Eel Tailed Catfish (Endangered Population). DPI's indicative distribution map indicates that none of the creeks within the Subject Land are mapped as habitat for the species (see Figure 5.4). The nearest creeks mapped as habitat for the Eel Tailed Catfish are:
 - Dumaresq River: it is located at approximately 1.2 km north from the Subject Land. This river runs in a general east-west direction.
 - Beardy River: is located to the east of the Subject Land. The nearest point is located at approximately 3 km from the development footprint. Beardy River runs on a general north-south direction from its convergence with Dumaresq River. None of the first order tributaries of Beardy River are located within the Subject Land.
- Olive Perchlet (Ambassis agassizii): The Development Site is located within the indicative distribution area of the Western Population of Olive Perchlet which is listed as an endangered population under the FM Act. The nearest creeks mapped as habitat for the Olive Perchlet are:
 - Dumaresq River: it is located at approximately 1.2 km north from the Subject Land. This river runs in a general east-west direction.
 - Beardy River: is located to the east from the Subject Land. The nearest point is located approximately 3 km from the development footprint. Beardy River runs on a general north-south direction from its convergence with Dumaresq River. None of the first order tributaries of Beardy River are located within the Subject Land.


5.8 Limitations To Data


It is possible that some species were not detected on the Subject Land due to surveys being completed outside of the appropriate/recommended survey period for those species. Where survey effort or timing is not consistent with the BAM or relevant guidelines, this is stated explicitly in the assessment and measures identified to address the limitation (e.g. assumed presence of Masked Owl and Barking Owl breeding habitats).


The calculation of hollow-bearings trees, in particular the size and number of hollows, was made from ground level. It is possible that some hollows are present that were not visible from ground level, which may result in underestimates.


The combined impacts of intensive livestock grazing and extended drought period may have altered the visible foliage cover of native grasses, forbs and high threat exotic species. Some native grasses and/or forbs may not have been present/visible at the time of survey. Likewise, some heavily grazed grasses were unidentifiable to species level.


It is also noted that at the time of surveys, all streams were dry and no detailed aquatic assessment has been undertaken on the Subject Land.











STAGE 2: IMPACT ASSESSMENT (BIODIVERSITY VALUES)

6. IMPACT ASSESSMENT (BIODIVERSITY VALUES)

This chapter identifies the potential impacts of the project on the biodiversity values of the Development Site. Measures taken to date to avoid and minimise impacts are summarised and recommendations are provided, which will assist GAIA to design a development that further avoids, minimises and mitigates impacts.

6.1 Avoiding and Minimising Impacts On Native Vegetation and Habitat

As discussed in the EIS, various options relating to location, technology and scale of the project were evaluated in developing the proposal. The site was selected as being a suitable site for a solar plant based on:

- A mostly cleared landscape with minimal vegetation removal required;
- Compatible land use zoning of the land; and
- Proximity to the transmission network.

The development footprint is of a scale that allows for flexibility in the design, allowing ecological and heritage constraints to be avoided (refer to EIS for greater detail on the site selection process).

The Project site has had extensive environmental investigation and assessment which has informed the current concept layout. The layout has been reassessed and reduced to minimise its impact on the environment. Careful consideration of the existing environmental constraints has seen the total development footprint reduced from 1,097 ha (including adjacent Lots to the west and east of the Project Boundary identified during the Preliminary Environmental Assessment) to 149.24 ha.

In terms of biodiversity values, GAIA has undertaken significant steps to avoid, minimise and mitigate impacts, as per the process outlined below:

- Identification of biodiversity values through comprehensive, rigorous and thorough biodiversity surveys;
- Communication of identified values to the project team;
- Consultation between the design team and project ecology leader to consider direct and indirect impacts and work through the design process to achieve a feasible project with the least biodiversity impact; and
- Consultation with BCD, to seek input and discuss measures proposed to avoid and minimise impacts (see Table 1.4).

Through continued detailed design the Project will avoid the following areas of high biodiversity value:

- Avoid the large areas of intact vegetation communities within the eastern and south eastern portion of the Project Boundary;
- Avoid the majority of areas identified as a TEC, with the exception of <1 ha of low condition vegetation representing a TEC. All areas of moderate condition vegetation representing a TEC have been avoided;
- The second and third order watercourses will be avoided through detailed design and survey, with riparian buffers to be applied either side of the streams, measured from the edge of the top of bank. The buffers applied to this BDAR are 20m either side of the second order stream and 30m to either side of the third order streams; and
- Locating ancillary facilities in areas where there are minimal biodiversity values.

Regardless of the above, the proposed development will result in unavoidable impacts to some areas containing native vegetation and habitat values to accommodate the necessary infrastructure associated with the proposal. These impacts are described in the following sections.

6.2 Avoiding and Minimising Prescribed Biodiversity Impacts

The BC Regulation (clause 6.1) identifies actions that are prescribed as impacts to be assessed under the biodiversity offsets scheme. Prescribed Biodiversity impacts relevant to the proposal are:

- Impacts of development on the habitat of threatened species associated with rocks.
- Impacts of development on the habitat of threatened species associated with human made structure.
- Impacts of development on the habitat of threatened species associated with non native vegetation.
- Impacts of development on the connectivity of different areas of habitat of threatened species that facilitates the movement of those species across their range.
- Impacts of development on movement of threatened species that maintains their life cycle.
- Impacts of development on water quality, water bodies and hydrological processes that sustain threatened species and threatened ecological communities.
- Impacts of vehicle strikes on threatened species or on animals that are part of a TEC.

Prescribed impacts requiring assessment are identified in Section 8.2.1.2 of the BAM (OEH 2017a). How these prescribed impacts have been avoided and minimised by the proposal is detailed in Table 6.1.

Table 6.1 Avoidance and Minimisation of Impacts

Impact	Avoidance/Minimisation Measures	Outcome
Prescribed Impact		
 (a) Impacts of development on the habitat of threatened species or ecological communities associated with: (i) karst, caves, crevices, cliffs and other geological features of significance, or (ii) rocks, or (iii) human made structures, or (iv) non-native vegetation 	A total of eleven rocky areas were recorded within the Subject Land. Five of these have been avoided by the proposal. Only two outcrops occur in the middle of the cleared paddocks and cannot be avoided by the development footprint. The remaining four may also be avoided during the detailed design and survey of the required riparian buffers and perimeter road. Vegetation within the Development Site comprises 108.03 ha of Cleared Land, Disturbed Grasslands and Dams. These areas were selected to be removed as they represent areas with reduced biodiversity value.	Avoidance of five rocky areas within the Project Boundary. Loss of only two rocky areas located in the centre of the Development Site. Four additional rocky areas likely to be avoided during detailed design.
(b) Impacts of development on the connectivity of different areas of habitat of threatened species that facilitates the movement of those species across their range.	The predominantly cleared landscape provides low quality connectivity across the Development Site itself. Isolated trees scattered across the Development Site represent limited connectivity features for highly mobile species to travel across the landscape. The second and third order creeks within the Development Site have some, albeit limited, potential to provide aquatic connectivity to Dumaresq River and have been avoided.	Retention of the second and third order streams which are the landscape features with highest potential as linking corridor.

Impact	Avoidance/Minimisation Measures	Outcome
(c) Impacts of development on movement of threatened species that maintains their life cycle	The predominantly cleared landscape provides low quality connectivity across the Development Site and the movement of the Grey-crowned Babbler largely confined to the woodland habitats and riparian corridors as this species has limited flying capacity. The riparian corridors (with a 20-30m buffer) have been avoided.	Retention of the second and third order creek lines along which movement of threatened species will continue to occur.
	Regarding the Cicadabird and microchiropteran bats, it is not considered the project will restrict the movement of these highly mobile aerial species and they will continue to utilise retained habitats surrounding the solar farm.	
(d) Impacts of development on water quality, water bodies and hydrological processes that sustain threatened	The second and third order watercourses will be avoided through detailed design (including 20-30m riparian buffer zones measured from the top of bank of the streams).	Avoidance of second and third order creek lines. Loss of six first order creeks and six farm dams.
species and threatened ecological communities (including from subsidence or upsidence resulting from underground mining)	Any waterway crossings required as a result of the Project will be designed in accordance with the Policy and Guidelines for Fish Habitat Conservation and Management and the Policy and Guidelines for Fish Friendly Waterway Crossings.	
	Six farm dams are present within the development site. These farm dams could not be avoided due to the size constraints of the solar panels. These farm dams would be filled in during construction of the solar farm. The impacts proposed to these dams are not anticipated to have any broader impacts for environments that sustain and interact with rivers, streams or wetlands either on or offsite.	
	Appropriate drainage features would be constructed along internal access roads to minimise the risk of polluted water leaving the site or entering water bodies. Ground cover would be maintained beneath the solar arrays and there would be a low risk of contamination in the event of a chemical spill (fuels, lubricants, herbicides etc.) as storage and emergency handling protocols will be implemented.	
(e) Impacts of wind turbine strikes on protected animals	Not applicable	Not applicable

Impact	Avoidance/Minimisation Measures	Outcome
(f) Impacts of vehicle strikes on threatened species or on animals that are part of a TEC.	The proposal would not directly increase impacts of vehicle strikes on threatened species. Threatened species would not be funnelled into transport corridors. An increase in vehicle traffic may indirectly increase vehicle strikes on native fauna. Site design would be unlikely to reduce impacts associated with vehicle strikes. Site management to enforce and reduce site speed limits would minimise the likelihood of vehicle strikes.	Increased traffic may indirectly increase vehicle strikes with fauna, including threatened species such as Greycrowned Babbler.
Impacts to Other Biodiversity	Values	
Loss of Species Credit Species Habitat or Individuals	Habitat for the Bristle-faced Free-tailed Bat will be retained within riparian buffers of 2 nd and 3 rd creek lines. The bat was also detected in one of the first order creeks to be retained. To further avoid impact, vegetation clearance shall only be undertaken outside breeding period for the species to prevent miscarriage in gravid females. Hollow bearing trees within the development footprint will be inspected prior to removal. If this species is confirmed utilising any of the hollow bearing trees, the trees will be left undisturbed until further advice is sought from BCD and a suitably recognised bat expert.	The species polygon for the Bristle-faced Free-tailed Bat occupies the entire Development Site. Offsets will be required to compensate for unavoidable impacts.
Loss of habitat for the Grey- crowned Babbler, Brown Treecreeper and Speckled Warbler	A total of 21 Grey-crowned Babbler nests were recorded across the assessment area, with 13 of these to be avoided. The predominantly cleared landscape provides low quality connectivity across the development site and the movement of the Grey-crowned Babbler is largely confined to woodland areas and riparian corridors. Riparian corridors (including 20-30m buffer) have been avoided. However, there will be some loss of suitable (albeit degraded) woodland habitat for this species and potentially for the Brown Treecreeper and Speckled Warbler (recorded adjacent to Subject Land during investigation of potential offset areas). To further minimise impacts, enhancement of linking corridors along 2 nd and 3 rd order riparian zones will be provided. This is expected to improve the current fragmented condition of riparian corridors and assist movement and dispersal of these and other fauna species.	Offsets will be required to compensate for unavoidable impacts.

Impact	Avoidance/Minimisation Measures	Outcome
Loss of Hollow Bearing Trees (HBT)	A total of 60 hollow bearing trees were recorded across the assessment area, including 34 HBT within the development footprint. Trees will be planted at a rate of 2:1 with two new trees planted for each HBT removed. A minimum of 68 trees shall be planted within the riparian corridor with commitment to 100% recruitment rate. As naturally formed tree hollows take many decades to develop, nest-boxes suitable for relevant hollow dependent species will be installed prior to HBT removal. Nest boxes shall be installed within suitable retained habitat (e.g. riparian corridors) and will be monitored and maintained during the construction and operational phases of the development.	The impacts of the loss of HBT will be minimised via revegetation and the provision of nest boxes .
Impacts of Development on the Habitat of Threatened Species or Ecological Communities	One TEC was recorded on and adjacent to the Subject Land – White Box Yellow Box Blakely's Red Gum Woodland. Impacts on this TEC have been largely avoided by retaining riparian and buffer areas. Threatened species recorded within the Development Site included the Grey-crowned Babbler and a number of microbats. In addition, Brown Treecreeper and Speckled Warbler were recorded adjacent to the Subject Land during investigation of potential offset areas. Large areas of intact vegetation representing better quality habitat for the above species have been avoided during the design phase of the project. Avoided areas include forest, woodland and riparian habitats. However, there will be some loss of suitable (albeit degraded) habitat for these species.	Implementation of mitigation measures will result in negligible impact on retained habitat of threatened species on and adjacent to the Development Site.
Impacts of Development on the Habitat of EPBC Listed Migratory Species	The project will not result in impacts on habitat of EPBC listed migratory species. One individual of the Cicadabird was recorded within the Development Site. The species occupies several types of vegetation across its range and forages on fruits and insects. No species-specific avoidance measures have been proposed for the Cicadabird. However, measures proposed for other threatened species will enhance roosting habitat and foraging resource availability for this migratory bird.	Implementation of mitigation measures will result in negligible impacts on migratory species.

6.3 Impact that Cannot be Avoided

6.3.1 Direct Impacts

Direct impacts associated with the construction and operational phases of the proposed development are outlined in Table 6.2.

Table 6.2 Direct Impacts to Biodiversity during the Construction and Operational Phases

Nature of Impact	Area of impact	Frequency/Timing	Description of the Impact	Threatened species and habitats likely to be affected
Loss of habitat	Up to approx. 150ha	Construction	Direct removal of native vegetation representing flora and fauna habitat. Direct removal of fallen timber and bush rock representing fauna habitat.	Grey-crowned Babbler and other threatened woodland birds. Microchiropteran bats, such as Bristle-faced Free- tailed Bat.
Displacement of resident fauna	Unknown	Construction and Operation	Direct loss of native fauna from the Subject Land.	Grey-crowned Babbler and other threatened woodland birds. Microchiropteran bats, such as Bristle-faced Free- tailed Bat.
Potential injury or death of fauna	Unknown	Construction and Operation	Injury and/or mortality of fauna during removal of vegetation and other fauna habitat components. Potential decline in local fauna populations.	Resident and dispersing fauna
Removal of important habitat features e.g. Hollow bearing trees	34 trees with hollows to be removed	Construction	Direct loss of native fauna habitat. Injury and mortality of fauna during clearing of fauna habitat and habitat trees.	Hollow dependent fauna, including arboreal mammals (e.g. Possum), microbats and birds
Bush Rock removal and disturbance	Up to six outcrops may be disturbed	Construction	Direct loss of native fauna habitat. Injury and mortality of fauna during clearing of fauna habitat	Reptiles Ground-dwelling mammals
Shading by solar infrastructure	Up to approx. 150ha	Operation	Modification of ecosystem grassland composition.	Flora species

The likely changes in vegetation integrity scores as a result of clearing for the solar array, laydown areas and access roads are documented for each vegetation zone in Table 6.3.

Table 6.3 Current and Future Vegetation Integrity Scores for each Vegetation Zone

Vegetation Zone		getation Zone PCT Area of Impact A	Area of Impact ^A	Vegetation integrity scores	
No	Name	-		Current	Future
1	594_Moderate	594	4.10	61.3	0
2	594_Low	594	9.54	33.3	0
3	596_Moderate	596	11.15	62.6	0
4	596_Low	596	0.25	38.2	0
5	596_Very Low	596	0.30	23.6	0
6	596_Derived_Low	596	9.04	8	0
7	516_Very Low	516	3.00	20.7	0
8	516_Derived_Moderate	516	2.76	24.7	0
9	544_Low	544	0.83	35	0
10	516_Disturbed Grassland	NA	7.39	15.8	0
11	594_Distrubed Grassland	NA	49.81	11.8	0
12	596_Disturbed Grassland	NA	50.59	5.6	0
13	Cleared Land / Dams	NA	0.48	-	-

6.3.2 Indirect Impacts

Identified indirect impacts that could occur as a result of the project include:

- Increased noise, vibration and dust levels;
- Increased sedimentation and erosion;
- Increased mortality and/or injury of fauna;
- Artificial lighting impacting nocturnal species behaviour; and
- Increase in invasive species, weeds and pathogens.

 Table 6.4
 Potential Indirect Impacts Associated with the Proposed Development

Nature of Impact	Area of impact	Frequency/ Timing	Description of the Impact and Consequence for Bioregional Persistence	Threatened species and habitats likely to be affected
Increased noise, vibration and dust levels	Development Site	Working hours during the Construction and Operational phases	Construction activities may result in increased levels of noise and vibration. No significant impacts are anticipated as the fauna abundance is low across the development site and largely limited to highly mobile species. A number of Grey-crowned Babblers and one confirmed nest were observed within the Development Site, family groups of the species stay together within their home range (generally 10 ha and up to 50 ha). It is expected that individuals of the local family group would move to roosting nests outside of the Development Footprint. No other threatened species are anticipated to rely on any of the habitats currently present and no sensitive receptors have been identified.	All fauna species, including the Greycrowned Babblers
Increase sedimentation and erosion	Development Site & Project Boundary	During and following the Construction and Operational Phases	Mobilisation of soils/sediments may occur during inclement weather over disturbed soils and sediments in areas where vegetation has been cleared and/or areas where soil and construction material has been stockpiled. Reduction in watercourse bank stability following any nearby construction and any clearing of riparian vegetation could also result in bank erosion and input of sediments into watercourses.	Aquatic fauna and habitat within the Dumaresq riparian system
Artificial lighting impacting	Development Site & Project Boundary	Night time during the Construction and Operational Phase	The project will require limited permanent night lighting, most likely for the operations and maintenance buildings and substations. Temporary, localised night lighting may be required during general maintenance activities conducted during the operational stage of the project. Lighting has the potential to impact species behaviour. Any impacts are anticipated to be highly localised and are not anticipated to be significant given the low diversity and abundance of fauna recorded within the Development Site.	All fauna, including the Grey-crowned Babbler and microbats
Increase in invasive species	Development Site & Project Boundary	During and following the Construction and Operational Phases	Increased movement of vehicles has the potential to transport weeds and pathogens into the development site and adjacent vegetation. Given the high levels of disturbance within the development site, there is also the risk that weeds may be transported off-site. Infection of native plants by <i>Phytophthora cinnamomi</i> is listed as a key threatening process under the BC Act and EPBC Act. <i>P. cinnamomi</i> can lead to death of trees and shrubs, resulting in devastation of native ecosystems (DECC 2008). As described by DoE (2014), infection of susceptible communities with <i>P. cinnamomi</i> can lead to:	Predation of fauna species, including threatened birds (e.g. Grey-crowned Babbler) and microbats (e.g. Bristle-faced Freetailed Bat)

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

Nature of Impact	Area of impact	Frequency/ Timing	Description of the Impact and Consequence for Bioregional Persistence	Threatened species and habitats likely to be affected
			 changes in the structure and composition of native plant communities; a significant reduction in primary productivity and functionality; and habitat loss and degradation for dependent flora and fauna. Clearing of vegetation and construction of roads provides increased potential for the introduction/spread of invasive pest species. Invasive species can also be unintentionally transported in vehicles and machinery. Monitoring and management of invasive species, such as cats, foxes and cane toads should be completed. 	
Impacts to Riparian Habitats	Development Site & Project Boundary	During and following the Construction and Operational Phases	Second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of these streams, measured from the edge of the high bank. This includes a 20m buffer either side of second order streams and 30m either side of third order streams. Furthermore, watercourses within the Development Site are highly disturbed being heavily grazed and eroded. The project is unlikely to result in any increased impact to the aquatic habitat, and the change in land use may improve habitats by reducing stocking rate and revegetating disturbed riparian areas.	Movement of fauna species, including the Grey-crowned Babbler and microbats.
Increased Fragmentation	Development Site	Permanent after clearing	The removal of native vegetation has the potential to result in fragmentation of fauna habitat, with resultant effects on fauna species movement, reproduction and gene flow. The impact of this vegetation clearance in an already highly modified landscape is anticipated to be negligible given that no significant fauna movement corridors currently exist within the development site (excluding riparian corridors).	All fauna species, including the Grey- crowned Babbler and microbats

6.3.3 Prescribed Impacts

Prescribed impacts requiring assessment are identified in Section 8.2.1.2 of the BAM (OEH 2017a). Avoidance of prescribed impacts is discussed in Section 6.2. These are discussed in Table 6.5 below.

Table 6.5 Identification of Prescribed Impacts associated with the Proposed Development

Development				
Prescribed Impact	Likely to occur as result of the proposed Solar Farm			
 (a) Impacts of development on the habitat of threatened species or ecological communities associated with: karst, caves, crevices, cliffs and other geological features of significance, or rocks, or human made structures, or non-native vegetation 	A total of eleven rocky areas were recorded within the Development Site. Up to six of these may be impacted by the proposed development. No threatened species were observed, or likely to be dependent on these six outcrops. No threatened species are likely to be dependent on any human made structures or areas of non-native vegetation.			
(b) Impacts of development on the connectivity of different areas of habitat of threatened species that facilitates the movement of those species across their range.	Isolated paddock trees scattered across the Development Site represent limited connectivity features for highly mobile species to travel across the landscape. A number of these trees would be removed as a result of the proposed development. The Subject Land does not represent a known connectivity link for threatened species and the proposal is therefore unlikely to significantly disrupt the movement of a threatened species across its range. However, the removal of some degraded woodland habitat is likely to represent a minor reduction in the availability of dispersal habitat for the Grey-crowned Babbler and potentially Brown Treecreeper and Speckled Warbler (recorded adjacent to Subject Land during investigation of potential offset areas).			
(c) Impacts of development on movement of threatened species that maintains their life cycle	A breeding population of Grey-crowned Babbler have been recorded on the Subject Land. The movement of this species is largely confined to the woodland habitats and riparian corridors as this species has limited flying capacity. Although riparian corridors (with a 20-30m buffer) have been avoided, there will be some impacts to degraded woodland habitat that may facilitate the movement of this species across the Subject Land. Regarding the Cicadabird and microchiropteran bats, it is not considered the project will restrict the movement of these highly mobile aerial species and they will continue to utilise retained habitats adjacent to the solar farm. The Subject Land does not represent a known migratory path for any threatened species. Due to the highly cleared and fragmented landscape the proposal is not likely to disrupt the movement of a threatened species that maintains their lifecycle.			

Prescribed Impact	Likely to occur as result of the proposed Solar Farm
(d) Impacts of development on water quality, water bodies and hydrological processes that sustain threatened species and threatened ecological communities (including from subsidence or upsidence resulting from underground mining)	Six farm dams are present within the development site. These farm dams would be filled in during construction of the solar farm. The filling of these dams is not anticipated to have any broader impacts for environments that sustain and interact with the rivers, streams and wetlands either onsite or offsite. These dams are not considered to represent important habitat for any threatened species or ecological community.
	The proposed development will affect six first order streams which are located within the proposed Development Site. These six first order creeks will be lost permanently.
	The proposal has the potential to affect surface water quality and quantity due to sediment runoff and/or contaminant runoff into adjacent watercourses. Pollutants to water courses may be associated with vehicle movement, spill of chemicals used for solar panel maintenance/cleaning and maintenance of associated infrastructure.
(e) Impacts of wind turbine strikes on protected animals	Not applicable
(f) Impacts of vehicle strikes on threatened species or on animals that are part of a TEC.	Vehicle strikes may occur during the construction and operational phases of the proposed development given the likelihood of increased vehicle movements.

6.4 Impacts to Matters of National Environmental Significance

An assessment of the impacts of the project on MNES within the Development Site was prepared to determine whether referral of the project to the Commonwealth Minister for the Environment is required. Matters of MNES relevant to the development site are summarised in Table 6.6 below.

A likelihood of occurrence assessment considering each entity individually is provided in Appendix D. Based on the results of the likelihood of occurrence and risk assessment, all TECs and threatened species as identified in the PMST were concluded to have a low residual risk. Given that one individual of the Cicadabird was recorded on the Subject Land, an assessment of significance for the species was undertaken as a precautionary measure (see Appendix E). No additional assessments of significance are considered to be required for other MNES (Appendix D).

Referral of the project to the Commonwealth Minister for the Environment for assessment is not required.

Table 6.6 Assessment of the Project in relation to EPBC Act

MNES	Predicted MNES	Result
Wetlands of International Importance	Three wetlands of national importance were identified as being between 1,100 and 1,300 km form the Development Site: Banrock Station Wetland Complex Riverland The Coorong, and lakes Alexandrina and Albert wetland.	No direct or indirect effects on these wetlands of international importance will result from the proposal.
Threatened Ecological Communities (TECs)	Four TECs were identified as likely to occur within the project area: Natural Grasslands on basalt and finetextured alluvial plains of northern New South Wales and Southern Queensland (CEEC) New England Peppermint (Eucalyptus nova-anglica) Weeping Myall Woodland (CEEC) Weeping Myall Woodlands (EEC) White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland (CEEC)	A likelihood of occurrence analysis was undertaken for these TECs (see Table D.3 in Appendix D). This analysis concluded that the proposed development is unlikely to affect EPBC listed TECs. Although the NSW listing of White Box – Yellow Box – Blakely's Red Gum Grassy Woodland is considered to be represented on the Subject Land by PCT 544, this vegetation does not satisfy the minimum condition criteria for the Commonwealth listing of this TEC. This determination has been made on the basis of very small patch sizes occurring on the Subject Land, low diversity of native understorey species and absence of "important species" as described in the Threatened Species Scientific Committee listing advice and National Recovery Plan (DECCW 2011).
Threatened Species	A total of 36 threatened species were predicted to occur within the project area, including: Eight mammals Seven birds Four reptiles One fish 16 flora	Seven species were identified as likely to occur within the Development Site and were taken into account during survey design. No EPBC listed threatened species have been recorded within the Development Site.
Migratory Species	A total of 11 migratory species were predicted to occur within the project area.	One migratory bird, Cicadabird (<i>Coracina tenuirostris</i>), was recorded during the surveys. A Significant Impact Assessment for this migratory species was undertaken (see Appendix E) and it was concluded that the species is unlikely to be significantly impacted as result of the proposed development. No further assessment for the Cicadabird is required.

7. MANAGEMENT AND MITIGATION MEASURES

The key measures required to mitigate the impacts of the proposal is provided below in Table 7.1.

Table 7.1 Recommended Mitigation Measures for Residual Impacts

Impact	Mitigation Measure	Responsibility	Timing	Reporting
Loss of Species Credit Species Habitat or Individuals	 Vegetation clearance: Preparation and implementation of a vegetation clearing protocol. Clearing to be supervised by an experienced fauna catcher / ecologist. Time works to avoid critical life cycle events such as breeding. Monitoring of tree hollows prior to removal to avoid impacting any breeding females or juveniles. If Bristle-cased Freetail Bat is confirmed utilising any of the hollow bearing trees, the trees will be left undisturbed and managed in accordance with the Biodiversity Management Plan. Replacement of trees at a rate of 2:1 (i.e. two trees will be planted to replace each hollow bearing tree removed). A minimum of 68 trees will be planted within the riparian corridor with commitment to a 100% recruitment rate. Nest-boxes suitable for hollow dependent microbats will be installed prior to HBT removal. Nest boxes shall be installed within suitable retained habitat (e.g. riparian corridors) and will be monitored and maintained during the construction and operational phases of the development. 	 Principal contractor to ensure implementation of vegetation clearing protocol. Experienced fauna catcher / Ecologist to supervise clearing and relocate native fauna. 	 Vegetation clearing protocol to be prepared by an Ecologist prior to vegetation clearing. Clearing supervision to occur during the entire clearing process. Letter with results of clearing to be prepared by the fauna catcher/Ecologist supervising the clearance. Letter to be available for review by delegated authority (if requested). 	Vegetation Clearing Protocol as part of the Biodiversity Management Plan

Impact	Mitigation Measure	Responsibility	Timing	Reporting
	 No stockpiling or storage within dripline of any mature trees. Approved clearing limits to be clearly delineated with temporary fencing or similar prior to construction commencing. 			
Grey-crowned Babbler habitat	 Nest removal: Approved clearing limits to be clearly delineated with temporary fencing or similar prior to construction commencing. Removal of trees with nests will be included in the vegetation clearing protocol including any seasonal constraints to avoid impacting any juveniles or unfledged chicks. Removal of trees with nests will be supervised by an experienced fauna catcher or ecologist. A portion of felled trees will be salvaged as habitat for fauna and translocated in suitable areas in the remainder of the Project Boundary. 	 Principal contractor to ensure implementation of vegetation clearing protocol. Experienced fauna catcher / Ecologist to supervise clearing and relocate native fauna 	 Clearing supervision to occur during the entire clearing process. Letter with results of clearing to be prepared by the fauna catcher/Ecologist supervising the clearance. Letter to be available for review by delegated authority (if requested). 	Vegetation Clearing Protocol as par of the Biodiversity Management Plan

Impact	Mitigation Measure	Responsibility	Timing	Reporting
Loss of Hollow Bearing Trees	 Replacement of trees at a rate of 2:1 (i.e. two trees will be planted to replace each hollow bearing tree removed). A minimum of 68 trees will be planted within the riparian corridor with commitment to a 100% recruitment rate. Nest-boxes suitable for hollow dependent microbats will be installed prior to HBT removal. Nest boxes shall be installed within suitable retained habitat (e.g. riparian corridors) and will be monitored and maintained during the construction and operational phases of the development. 	 GAIA to appoint a qualified ecologist for preparation of the monitoring plan. GAIA to liaise with relevant authority regarding outcomes. Appointed Ecologist to undertake monitoring as required. 	 Monitoring Plan to be prepared and approved prior to commencement of clearing. Monitoring to be implemented as required. 	Tree Replacement and Nest Box Monitoring Plan as part of the Biodiversity Management Plan
Impacts of Development on the Habitat of Threatened Species or Ecological Communities	 A tree replacement and nest box monitoring plan will be prepared for the Project. The plan will provide details of monitoring and Key Performance Indicators (KPIs) to ensure objectives of tree replacement and nest box monitoring is achieved. Monitoring and reporting to be undertaken by a qualified ecologist. Avoidance of use of chemicals, such as pesticides and herbicides, within the solar farm during the construction and operational phases to prevent contributing to the global decline in insect population and diversity. Facilitation of natural regeneration of native ground cover within viable solar farm footprint (e.g. under solar panel arrays) and in retained areas. This will include management of weeds. 	 GAIA to appoint a qualified ecologist for preparation of the monitoring plan. GAIA to liaise with relevant authority regarding outcomes. Appointed Ecologist to undertake monitoring as required. 	 Monitoring Plan to be prepared and approved prior to commencement of clearing. Monitoring to be implemented as required. 	Tree Replacement and Nest Box Monitoring Plan as part of the Biodiversity Management Plan

Impact	Mitigation Measure	Responsibility	Timing	Reporting
Impacts of Development on the Habitat of EPBC Listed Migratory Species	None required	NA	NA	NA
Impacts of Development on Water Quality, Water Bodies and Hydrological Processes that sustain Threatened Species and TECs	 An erosion and sediment control plan (ESCP) would be prepared in conjunction with the final design and implemented. Design of creek crossings to meet best practice industry standards. ESCP to include requirements for water quality monitoring, chemical use and control. 	 GAIA has overall responsibility to ensure meeting that they are meeting environmental commitments. 	 ESCP to be prepared and approved prior to commencement of works. Reporting, evaluation and auditing as per the ESCP. 	CEMP ESCP
Impacts of Vehicle Strikes on Threatened Species of animals or on animals that are part of a TEC	Actions to minimise mortality of wildlife involved in vehicle strikes: Appropriate vehicle speeds and signage to be installed within the solar farm. Protocol detailing actions to be undertaken in the event of a vehicle strike. Identification of a wildlife veterinary and/or wildlife carer group and agreement for injured wildlife to be taken care of or being humanely euthanised.	GAIA has overall responsibility to ensure meeting that they are meeting environmental commitments.	 CEMP to be prepared and approved prior to commencement of works. Reporting, evaluation and auditing as per the CEMP. 	CEMP
	Avoid night works.Direct lights away from retained native vegetation.	 GAIA has overall responsibility to ensure meeting that they are meeting environmental commitments. 	 CEMP to be prepared and approved prior to commencement of works. Reporting, evaluation and auditing as per the CEMP. 	СЕМР

Impact	Mitigation Measure	Responsibility	Timing	Reporting
Invasive Species	 CEMP will include a Management protocol for declared priority weeds under the <i>Biosecurity Act 2015</i> during and after construction. Hygiene protocols to prevent the spread of weeds or pathogens between infected areas and uninfected areas. Monitoring and management protocol for invasive feral/pest species, including cats, foxes and cane tods. 	GAIA has overall responsibility to ensure meeting that they are meeting environmental commitments.	 CEMP to be prepared and approved prior to commencement of works. Reporting, evaluation and auditing as per the CEMP. 	СЕМР

8. SERIOUS AND IRREVERSIBLE IMPACTS (SAII)

A Serious and Irreversible Impact (SAII) is listed under the BC Act as an impact that is likely to contribute significantly to the risk of extinction of a threatened entity.

In accordance with the BAM, species and ecological communities with a 'very high' biodiversity risk weighting will be a potential serious and irreversible impact (SAII). Whenever potential SAII are identified for a Development Site, those SAII need to be address as per Section 10.2 of the BAM (OEH 2017a).

The following guidelines were consulted to identify potential SAII:

- OEH (2017b) Guidance to assist a decision-maker to determine a serious and irreversible impact.
- OEH (2018b) 'Species Credit' threatened bats and their habitats guide.

8.1 Potential Serious and Irreversible Impact Entities

Based on candidate ecosystem credit species, species credit species and results of field surveys, the potential SAII for the Development Site are listed in Table 8.1 below.

Table 8.1 Potential SAII within the Development Site

Scientific Name Common Name	Justification as potential SAII	Corresponding habitat constraint	Is SAII present?
Miniopterus orianae oceanensis Large Bent-winged Bat	The species is dependent on non- responding attribute (breeding habitat only) (OEH 2017b)	Caves are the primary breeding habitat for the Eastern Bentwing Bat. Breeding habitat does not occur on the Subject Land.	No
Vespadelus troughtoni Eastern Cave Bat	Breeding habitat is classified as SAII (OEH 2018b)	Breeding habitat for the species is caves, overhangs, crevices, cliffs and escarpments, or old mines or tunnels, old buildings and sheds within the potential habitat. Breeding habitat is not considered to occur on the Subject Land.	No
White Box Yellow Box Blakely's Red Gum Woodland	Threatened Ecological Community with very high sensitivity to loss due to small population size (extent)	Fertile lower footslopes and flats. This TEC is represented onsite by PCT 544 which is associated with stream banks, beds and channels. Riparian areas associated with this TEC have largely been avoided.	No

Client: GAIA Australia

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

8.1.1 Threatened Ecological Communities

One threatened ecological community, that is a candidate for serious and irreversible impacts, was identified on the Subject Land – White Box Yellow Box Blakely's Red Gum Woodland. This TEC is represented on the Subject Land by disturbed areas of PCT 544 – Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion of New England Tableland Bioregion. The majority of this TEC has been avoided by retaining riparian corridors and associated buffers. The proposed development is not considered to represent a SAII on this TEC as impacts on 0.83 ha of highly disturbed vegetation representing PCT 544 are unlikely to contribute to the risk of extinction of the TEC. This TEC will be allowed to naturally regenerate within retained riparian areas and buffers. Infill plantings will also be provided that will expand the extent of this TEC on the Subject Land and improve its current condition.

8.1.2 Threatened Species

Two potential SAII entities were identified, the Eastern Bentwing Bat and the Eastern Cave Bat. The echolocation calls of these species were recorded within the assessment area. However, breeding habitat for these microbat species, i.e. caves, overhangs crevices, cliffs or escarpments, are not present on the Subject Land. It is concluded that due to the lack of suitable breeding habitat, the Subject Land represents foraging habitat only. Therefore, the proposed development does not represent a SAII on these species.

8.1.3 Additional Potential Entities

No other threatened species or communities, were identified as potential candidates for serious and irreversible impacts.

8.2 Assessment of SAII

The proposed development is not considered to represent a serious and irreversible impact on any identified SAII entities.

www.erm.com Version: 2.0 Project No.: 0470861 0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

9. REQUIREMENT TO OFFSET

This chapter provides an assessment of the impacts requiring offsetting in accordance with Section 10 of the BAM (OEH 2017a).

9.1 Impacts Requiring an Offset

9.1.1 Impacts on Native Vegetation – Ecosystem Credits

Impacts to native vegetation requiring offsets include:

- Direct impacts on PCT 516 Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion;
- Direct impacts on PCT 544 Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion;
- Direct impacts on PCT 594 Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion; and
- Direct impacts on PCT 596 Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion

A total of 598 ecosystem credits are required to offset impacts of the proposed development. The impacts on the vegetation zones within each of the four PCTs is shown in Table 9.1.

Table 9.1	Summary	y of Ecos	ystem Credits
-----------	---------	-----------	---------------

Vegetation Zone Number	Vegetation Zone Name	Area (ha)	Vegetation Integrity Score	Future Vegetation Integrity Score	Change in Vegetation Integrity Score	Credits Required
PCT 594 – Silver-leaved Ironbark – White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion					Bioregion	
1	594_Moderate	4.1	61.3	0	-61.3	110
2	594_Low	9.54	33.3	0	-33.3	139
	Total Ecosystem Credits required to offset impacts on PCT 594					249
	umbledown Red Gum – W Nandewar Bioregion	hite Cyp	ress Pine – Sil	ver-leaved Iror	nbark shrubby woodlan	d mainly in
3	596_Moderate	11.15	62.6	0	-62.6	262
4	596_Low	0.25	38.2	0	-38.2	4
5	596_Very Low	0.30	23.6	0	-23.6	3
Total Ecosystem Credits required to offset impacts on PCT 596					269	

Vegetation Zone Number	Vegetation Zone Name	Area (ha)	Vegetation Integrity Score	Future Vegetation Integrity Score	Change in Vegetation Integrity Score	Credits Required
PCT 516 – Grey Box grassy woodland or open forest of the Nandewar Bioregion of New England Tableland Bioregion						
7	516_Very Low	3.00	20.7	0	-20.7	31
8	516_Derived_Moderate	2.76	24.7	0	-24.7	34
Total Ecosystem Credits required to offset impacts on PCT 516					65	
PCT 544 – Rough-barked Apple – White Cypress Pine – Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion of New England Tableland Bioregion					voodland of	
9	544_Low	0.83	35	0	-35	15
Total Ecosystem Credits required to offset impacts on PCT 544					15	

9.1.2 Impacts on Threatened Species – Species Credits

Impacts to habitat of Threatened Species requiring offsets include:

- Habitat for the Bristle-faced Free-tailed Bat in PCTs 516, 544, 594 and 596, requiring 1,180 species credits;
- Foraging habitat for Eastern Cave Bat in PCTs 516, 544, 594 and 596, requiring 1,030 species credits; and
- Potential (assumed) breeding habitat for Masked Owl and Barking Owl in PCTs 516, 544, 594 and 596, where suitably sized hollows occur, requiring 680 species credits for each of these species.

A total of 3,570 species credits are required to offset loss of fauna habitat within the Development Site.

9.1.3 Offsets Required Under the EPBC Act

No offsets requirements under the EPBC Act have been identified for the proposed development.

9.2 Impacts Not Requiring an Offset

In accordance with the BAM, an offset is not required for impacts on native vegetation where the vegetation integrity score is below those set out in Paragraph 10.3.1.1. These thresholds are as follows:

- a) A vegetation zone that has a vegetation integrity score <15 where the PCT is representative of an endangered or critically endangered ecological community, or
- A vegetation zone that has a vegetation integrity score <17 where the PCT is associated with threatened species habitat (as represented by ecosystem credits), or is representative of a vulnerable ecological community, or

c) A vegetation zone that has a vegetation integrity score <20 where the PCT is not representative of a TEC or associated with threatened species habitat.

Areas of the Subject Land that are not required to be offset in accordance with the thresholds described above are provided in Table 9.2. These vegetation zones represent highly disturbed areas, including cleared areas, heavily grazed grasslands and low integrity derived grasslands.

Table 9.2 Summary of Impacts on Vegetation Not Requiring Offsets

Vegetation Zone Number	Vegetation Zone Name	Area (ha)	Vegetation Integrity Score	Future Vegetation Integrity Score	Change in Vegetation Integrity Score	Credits Required
PCT 596 – Tumbledown Red Gum – White Cypress Pine – Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion						
6	596_Derived_Low	9.04	8	0	-8	0
12	596_Disturbed Grasslands	50.59	5.6	0	-5.6	0
PCT 594 - Silve and Nandewar E	er-leaved Ironbark – Wh Bioregion	nite Cypre	ss Pine shrubby	open forest of B	rigalow Belt Sou	ıth Bioregion
11	594_Disturbed Grasslands	49.81	11.8	0	-11.8	0
PCT 516 – Grey Box grassy woodland or open forest of the Nandewar Bioregion of New England Tableland Bioregion						
10	516_Disturbed Grassland	7.39	15.8	0	-15.8	0

9.3 Summary of Offset Credits Required

The proposed development will incur loss of native vegetation and threatened fauna habitat requiring offsets for a total of 598 ecosystem credits and 3,570 species credits.

9.4 Biodiversity Offset Framework

This section outlines several options that are available to proponents to offset project related impacts on biodiversity in accordance with the Biodiversity Offset Scheme (BOS). The final offset strategy for the project is yet to be developed. However, GAIA is committed to satisfying all offset requirements prior to the commencement of works (including bulk earthworks and vegetation removal).

In accordance with the BOS, proponents may use one or more of the following methods to satisfy an offset requirement:

- Purchasing credits: Identify and purchase the required 'like for like' credits in the market and then
 retire those credits via Biodiversity Offsets and Agreement Management System (BOAMS).
- Generating credits: Identify potential offset land containing the biodiversity values required to be
 offset. Generate credits by applying the BAM and placing a Biodiversity Stewardship Agreement
 (BSA) on proposed offset land.

- Offsets Payment Calculator: Determine the cost of the credit obligation and transfer this amount to the Biodiversity Conservation Fund via the BOAMS. The Biodiversity Conservation Trust is then responsible for identifying and securing the credit obligation.
- Submit approval to the consent authority to apply variation rules. This excludes impacts on threatened species and TECs.

Given that impacts on threatened species habitat are associated with the proposed development, variation rules are not applicable to the Bonshaw Solar Farm development. GAIA will use one, or a combination, of the first three options above to meet the offset obligation.

The BAM Biodiversity Credit Report (like-for-like) detailing requirements for offsetting impacts on PCTs (i.e. 516, 544, 594 and 596) and threatened fauna habitat (i.e. Bristle-faced Free-tailed Bat, Eastern Cave Bat, Barking Owl and Masked Owl) is provided in Appendix K.

10. CONCLUSIONS

GAIA is proposing to develop a solar farm in disturbed land located in Lot 2 DP 1039185 (Project Boundary), Bonshaw within the Inverell LGA. The Development Site is located in the western portion of the Project Boundary and will occupy approximately 149.24 ha. The purpose of this BDAR was to address the requirements of the BAM and to address the biodiversity matters raised in the SEARs.

Based on the results of extensive field surveys, a total of 12 vegetation zones were identified on the Subject Land pertaining to the following four plant community types (PCTs):

- PCT 516 Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion;
- PCT 544 Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest/woodland of the Nandewar Bioregion and New England Tableland Bioregion;
- PCT 594 Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion; and
- PCT 596 Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion.

PCT 544 is considered to represent the following Threatened Ecological Community, which is listed as Endangered under the NSW *Biodiversity Conservation Act 2016:*

White Box Yellow Box Blakely's Red Gum Woodland

The occurrence of this TEC on the Subject Land is not considered to satisfy the minimum condition criteria for the Commonwealth listing of this TEC. This determination has been made on the basis of very small patch sizes occurring on the Subject Land, low diversity of native understorey species and absence of "important species" as described in the Threatened Species Scientific Committee listing advice and National Recovery Plan for the TEC.

An assessment of the potential impacts on the biodiversity values of the Subject Land has identified the following direct impacts to native vegetation requiring offset:

- Impacts on PCT 516 Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion, requiring 65 ecosystem credits;
- Impacts on PCT 544 Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest/woodland of the Nandewar Bioregion and New England Tableland Bioregion, requiring 15 ecosystem credits;
- Impacts on PCT 594 Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion ad Nandewar Bioregion, requiring 249 ecosystems credits; and
- Impacts on PCT 596 Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion, requiring 269 ecosystem credits

A total of 598 ecosystem credits are required to offset the impacts of the project.

A total of thirteen threatened fauna species were recorded within the Project Boundary. Threatened species included eight vulnerable microchiropteran bats (Little Pied Bat, Eastern Bent-wing Bat, Corben's Long-eared Bat, Yellow-bellied Sheath-tailed Bat, Eastern Cave Bat, Eastern False Pipistrelle, Greater Broad-nosed Bat and Hoary Wattled Bat), one endangered microchiropteran bat (Bristle-faced Free-tailed bat), three vulnerable birds (Grey-crowned Babbler, Brown treecreeper and Speckled Warbler) and one migratory bird (Cicadabird). Out of the eight Vulnerable microchiropteran bats, six were recorded as "definite" calls and two (Greater Broad-nosed Bat and Hoary Wattled Bat) were recorded as "possible" calls by a microbat call identification expert. Brown Treecreeper and Speckled Warbler were recorded within the Project Boundary but not directly on the Subject Land.

Impacts on threatened fauna habitat generated the following species credits:

Revised Biodiversity Development Assessment Report

- Bristle-faced Free-tailed Bat, requiring a total of 1,180 species credits;
- Eastern Cave Bat, requiring a total of 1,030 species credits;
- Masked Owl, requiring a total of 680 species credits; and
- Barking Owl, requiring a total of 680 species credits.

The proposed Bonshaw Solar Farm development in its current footprint requires a total offset of 598 ecosystem credits and 3,570 species credits. Retirement of these credits will be carried out in accordance with the *Biodiversity Conservation Regulation 2017* and the NSW Biodiversity Offsets Scheme.

11. REFERENCES

- DECC (2002) **Descriptions for NSW (Mitchell) Landscapes Version 2.** NSW Department of Environment and Climate Change.
- DECCW (2011) National Recovery Plan. White Box Yellow Box Blakely's Red Gum Grassy Woodland and Derived Native Grassland. A critically endangered ecological community. May 2011.
- DoE (2013) **Matters of National Environmental Significance**. Significant Impact Guidelines 1.1 under the Environment Protection and Biodiversity Conservation Act 1999. Australian Government. Department of the Environment.
- DoEE (2012) Interim Biogeographic Regionalization for Australia, version 7.
- DPI (2008) **Threatened Species Assessment Guidelines**. The assessment of significance. NSW DPI Threatened Species Management.
- DPI (2012) Controlled Activities on Waterfront Land Guidelines for watercourse crossings on waterfront land. NSW Department of Industry (July 2012).
- DPI (2018) Fact Sheet: Controlled Activity approval exemptions. NSW Department of Industry (August 2018).
- Environment Australia (2000) **Revision of the Interim Biogeographic Regionalisation for Australia (IBRA) and development version 5.1. Summary Report.**
- NPWS (1999) Natural tree hollows Essential for Wildlife. Voluntary Conservation on Private and Public Land Note 5. NSW National Parks and Wildlife Service.
- OEH (2016) 'Species credit' threatened bats and their habitats. Office of Environment and Heritage for the NSW Government, Sydney, NSW.
- OEH (2017a) **Biodiversity Assessment Methodology (BAM).** Office of Environment and Heritage for the NSW Government, Sydney, NSW.
- OEH (2017b) Guidance to assist a decision-maker to determine a serious and irreversible impact. Office of Environment and Heritage for the NSW Government.
- OEH (2018a) **Biodiversity Assessment Method Operational Manual Stage 1**. Office of Environment and Heritage for the NSW Government.
- OEH (2018b) 'Species credit' threatened bats and their habitats. NSW survey guide for the Biodiversity Assessment Method. Office of Environment and Heritage for the NSW Government, Sydney, NSW.
- OEH (2019) Threatened Biodiversity Profiles. NSW Office of Environment and Heritage. On-line profiles accessed via: https://www.environment.nsw.gov.au/threatenedSpeciesApp/
- Peacock, R., Rolhauser, A., Thönell, J. & Law, E. (2009). Extant and potential natural vegetation of Yallaroi, Ashford, Bingara and Inverell 1:100,000 scale map sheets, NSW. Appendix 1 Local Vegetation Community profiles. Draft Version 1.1 Revision date 31 July 2009.

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development Assessment Report APPENDIX A BOSET REPORT

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx



Biodiversity Offset Scheme (BOS) Entry Threshold Map 1: 59,978 3.0 Kilometers 1.52 This map is a user generated static output from an Internet mapping site and is for reference only. Data layers that appear on this map may or may not be accurate, current, or otherwise reliable. GCS_GDA_1994 THIS MAP IS NOT TO BE USED FOR NAVIGATION

Legend

Biodiversity Values that have been mapped for more than 90 days

Biodiversity Values added within last 90 days

Notes

© Office of Environment and Heritage | NSW Environment & Heritage



Biodiversity Values Map and Threshold Report

Results Summary

Date of Calculation	08/01/2019 10:12 AM	BDAR Required*
Total Digitised Area	356.35 ha	
Minimum Lot Size Method	Lot size	
Minimum Lot Size	199.02 ha	
Area Clearing Threshold	1 ha	
Area clearing trigger Area of native vegetation cleared	Unknown [#]	Unknown [#]
Biodiversity values map trigger Impact on biodiversity values map(not including values added within the last 90 days)?	no	no

*If BDAR required has:

- at least one 'Yes': you have exceeded the BOS threshold. You are now required to submit a Biodiversity Development Assessment Report with your development application. Go to https://customer.lmbc.nsw.gov.au/assessment/AccreditedAssessor to access a list of assessors who are accredited to apply the Biodiversity Assessment Method and write a Biodiversity Development Assessment Report
- 'No': you have not exceeded the BOS threshold. You may still require a permit from local council. Review the development control plan and consult with council. You may still be required to assess whether the development is "likely to significantly affect threatened species' as determined under the test in s. 7.3 of the Biodiversity Conservation Act 2016. You may still be required to review the area where no vegetation mapping is available.
- # Where the area of impact occurs on land with no vegetation mapping available, the tool cannot determine the area of native vegetation cleared and if this exceeds the Area Threshold. You will need to work out the area of native vegetation cleared refer to the BOSET user guide for how to do this.

Disclaimer

This results summary and map can be used as guidance material only. This results summary and map is not guaranteed to be free from error or omission. The State of NSW and Office of Environment and Heritage and its employees disclaim liability for any act done on the information in the results summary or map and any consequences of such acts or omissions. It remains the responsibility of the proponent to ensure that their development application complies will all aspects of the *Biodiversity Conservation Act 2016*.

The mapping provided in this tool has been done with the best available mapping and knowledge of species habitat requirements. This map is valid for a period of 30 days from the date of calculation (above).

Acknowledgement

I as the applicant for this development, submit that I have correctly	depicted the area that will be impacted or	likely to be impacted as a
result of the proposed development.		

Signature	Data:	08/01/2019 10:12 AM

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development Assessment Report					
Revised blodiversity Development	nosessillett report				
APPENDIX B	NSW BIONET ATLAS RECORDS				

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

Results of search of existing records of threatened species and Threatened Ecological Communities (TECs) in the NSW BioNet Atlas is summarised in Table B.1 below.

Table B.1 BioNet Atlas results

Family	Scientific Name	Common Name	NSW status	Comm. status	Records within 10km radius
Reptilia					
Gekkonidae	Uvidicolus sphyrurus	Border Thick-tailed Gecko	V	V	1
Aves	'				
Cacatuidae	Calyptorhynchus lathami	Glossy Black-Cockatoo	V		1
Psittacidae	Glossopsitta pusilla	Little Lorikeet	V		2
Psittacidae	Neophema pulchella	Turquoise Parrot	V		6
Climacteridae	Climacteris picumnus victoriae	Brown Treecreeper (eastern subspecies)	V		2
Acanthizidae	Chthonicola sagittata	Speckled Warbler	V		2
Pomatostomidae	Pomatostomus temporalis temporalis	Grey-crowned Babbler (eastern subspecies)	V		3
Neosittidae	Daphoenositta chrysoptera	Varied Sittella	V		1
Artamidae	Artamus cyanopterus cyanopterus	Dusky Woodswallow	V		1
Petroicidae	Petroica boodang	Scarlet Robin	V		1
Estrildidae	Stagonopleura guttata	Diamond Firetail	V		2
Mammalia	·				
Phascolarctidae	Phascolarctos cinereus	Koala	V	V	9
Petauridae	Petaurus norfolcensis	Squirrel Glider	V		1
Macropodidae	Petrogale penicillata	Brush-tailed Rock-wallaby	E	V	1
Vespertilionidae	Falsistrellus tasmaniensis	Eastern False Pipistrelle	V		1

GAIA BONSHAW SOLAR EIS


Revised Biodiversity Development Assessment Report

Family	Scientific Name	Common Name	NSW status	Comm. status	Records within 10km radius
Vespertilionidae	Miniopterus orianae oceanensis	Eastern Bentwing-bat	V		1
Vespertilionidae	Scoteanax rueppellii	Greater Broad-nosed Bat	V		2
Flora			'		
Araliaceae	Astrotricha roddii	Rodd's Star Hair	E	E	12
Asteraceae	Rutidosis heterogama	Heath Wrinklewort	V	V	1
Fabaceae (Faboideae)	Swainsona sericea	Silky Swainson-pea	V		1
Myrtaceae	Eucalyptus caleyi subsp. ovendenii	Ovenden's Ironbark	V	V	1
Myrtaceae	Eucalyptus nicholii	Narrow-leaved Black Peppermint	V	V	1

Notes: V = Vulnerable; E = Endangered

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page B2

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development	Assessment Report
_	
APPENDIX C	MATTERS OF NATIONAL SIGNIFICANCE ASSESSMENT – PMST REPORT
	FWST REPORT



EPBC Act Protected Matters Report

This report provides general guidance on matters of national environmental significance and other matters protected by the EPBC Act in the area you have selected.

Information on the coverage of this report and qualifications on data supporting this report are contained in the caveat at the end of the report.

Information is available about <u>Environment Assessments</u> and the EPBC Act including significance guidelines, forms and application process details.

Report created: 08/01/19 09:21:50


Summary

Details

Matters of NES
Other Matters Protected by the EPBC Act
Extra Information


Caveat

<u>Acknowledgements</u>



This map may contain data which are ©Commonwealth of Australia (Geoscience Australia), ©PSMA 2010

Coordinates
Buffer: 5.0Km



Summary

Matters of National Environmental Significance

This part of the report summarises the matters of national environmental significance that may occur in, or may relate to, the area you nominated. Further information is available in the detail part of the report, which can be accessed by scrolling or following the links below. If you are proposing to undertake an activity that may have a significant impact on one or more matters of national environmental significance then you should consider the <u>Administrative Guidelines on Significance</u>.

World Heritage Properties:	None
National Heritage Places:	None
Wetlands of International Importance:	3
Great Barrier Reef Marine Park:	None
Commonwealth Marine Area:	None
Listed Threatened Ecological Communities:	4
Listed Threatened Species:	36
Listed Migratory Species:	11

Other Matters Protected by the EPBC Act

This part of the report summarises other matters protected under the Act that may relate to the area you nominated. Approval may be required for a proposed activity that significantly affects the environment on Commonwealth land, when the action is outside the Commonwealth land, or the environment anywhere when the action is taken on Commonwealth land. Approval may also be required for the Commonwealth or Commonwealth agencies proposing to take an action that is likely to have a significant impact on the environment anywhere.

The EPBC Act protects the environment on Commonwealth land, the environment from the actions taken on Commonwealth land, and the environment from actions taken by Commonwealth agencies. As heritage values of a place are part of the 'environment', these aspects of the EPBC Act protect the Commonwealth Heritage values of a Commonwealth Heritage place. Information on the new heritage laws can be found at http://www.environment.gov.au/heritage

A <u>permit</u> may be required for activities in or on a Commonwealth area that may affect a member of a listed threatened species or ecological community, a member of a listed migratory species, whales and other cetaceans, or a member of a listed marine species.

Commonwealth Land:	None
Commonwealth Heritage Places:	None
Listed Marine Species:	18
Whales and Other Cetaceans:	None
Critical Habitats:	None
Commonwealth Reserves Terrestrial:	None
Australian Marine Parks:	None

Extra Information

This part of the report provides information that may also be relevant to the area you have nominated.

State and Territory Reserves:	1
Regional Forest Agreements:	1
Invasive Species:	19
Nationally Important Wetlands:	None
Key Ecological Features (Marine)	None

Details

Fish

Maccullochella peelii Murray Cod [66633]

Matters of National Environmental Significance

Wetlands of International Importance (Ramsar)	[Resource Information]
Name	Proximity
Banrock station wetland complex	1100 - 1200km
Riverland	1100 - 1200km
The coorong, and lakes alexandrina and albert wetland	1300 - 1400km

Listed Threatened Ecological Communities		[Resource Information]	
For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.			
Name	Status	Type of Presence	
Natural grasslands on basalt and fine-textured alluvial plains of northern New South Wales and southern Queensland	Critically Endangered	Community likely to occur within area	
New England Peppermint (Eucalyptus nova-anglica) Grassy Woodlands	Critically Endangered	Community may occur within area	
Weeping Myall Woodlands	Endangered	Community may occur within area	
White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland	Critically Endangered	Community likely to occur within area	
Listed Threatened Species		[Resource Information]	
Name	Status	Type of Presence	
Birds			
Anthochaera phrygia Regent Honeyeater [82338]	Critically Endangered	Foraging, feeding or related behaviour likely to occur within area	
Calidris ferruginea			
Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area	
Erythrotriorchis radiatus			
Red Goshawk [942]	Vulnerable	Species or species habitat likely to occur within area	
Geophaps scripta scripta Squatter Pigeop (southern) [64440]	Vulnerable	Species or species habitat	
Squatter Pigeon (southern) [64440]	vuliterable	Species or species habitat likely to occur within area	
Grantiella picta			
Painted Honeyeater [470]	Vulnerable	Species or species habitat likely to occur within area	
Lathamus discolor			
Swift Parrot [744]	Critically Endangered	Species or species habitat likely to occur within area	
Rostratula australis	En de s		
Australian Painted-snipe, Australian Painted Snipe [77037]	Endangered	Species or species habitat may occur within area	

Vulnerable

Species or species

Name	Status	Type of Presence habitat known to occur
Mammals		within area
Chalinolobus dwyeri Large-eared Pied Bat, Large Pied Bat [183]	Vulnerable	Species or species habitat likely to occur within area
Dasyurus maculatus maculatus (SE mainland population) Spot-tailed Quoll, Spotted-tail Quoll, Tiger Quoll (southeastern mainland population) [75184]	i <mark>on)</mark> Endangered	Species or species habitat likely to occur within area
Nyctophilus corbeni Corben's Long-eared Bat, South-eastern Long-eared Bat [83395]	Vulnerable	Species or species habitat likely to occur within area
Petauroides volans Greater Glider [254]	Vulnerable	Species or species habitat may occur within area
Petrogale penicillata Brush-tailed Rock-wallaby [225]	Vulnerable	Species or species habitat known to occur within area
Phascolarctos cinereus (combined populations of Qld, Koala (combined populations of Queensland, New South Wales and the Australian Capital Territory) [85104] Pseudomys novaehollandiae	NSW and the ACT) Vulnerable	Species or species habitat likely to occur within area
New Holland Mouse, Pookila [96]	Vulnerable	Species or species habitat likely to occur within area
Pteropus poliocephalus Grey-headed Flying-fox [186]	Vulnerable	Foraging, feeding or related behaviour may occur within area
Plants		
Acacia pubifolia Velvet Wattle [19799]	Vulnerable	Species or species habitat may occur within area
Astrotricha roddii [56312]	Endangered	Species or species habitat known to occur within area
Boronia granitica Granite Boronia [18598]	Endangered	Species or species habitat may occur within area
Cadellia pentastylis Ooline [9828]	Vulnerable	Species or species habitat likely to occur within area
Callistemon pungens [55581]	Vulnerable	Species or species habitat likely to occur within area
<u>Dichanthium setosum</u> bluegrass [14159]	Vulnerable	Species or species habitat likely to occur within area
Eucalyptus caleyi subsp. ovendenii Ovenden's Ironbark [56193]	Vulnerable	Species or species habitat known to occur within area
Eucalyptus mckieana McKie's Stringybark [20199]	Vulnerable	Species or species habitat likely to occur within area
Eucalyptus nicholii Narrow-leaved Peppermint, Narrow-leaved Black Peppermint [20992]	Vulnerable	Species or species habitat known to occur within area

Name	Status	Type of Presence
Homopholis belsonii Belson's Panic [2406]	Vulnerable	Species or species habitat may occur within area
Lepidium peregrinum Wandering Pepper-cress [14035]	Endangered	Species or species habitat may occur within area
Leucopogon confertus Torrington Beard-heath [14417]	Endangered	Species or species habitat may occur within area
Prasophyllum sp. Wybong (C.Phelps ORG 5269) a leek-orchid [81964]	Critically Endangered	Species or species habitat may occur within area
Rutidosis heterogama Heath Wrinklewort [13132]	Vulnerable	Species or species habitat known to occur within area
Thesium australe Austral Toadflax, Toadflax [15202]	Vulnerable	Species or species habitat likely to occur within area
Tylophora linearis [55231]	Endangered	Species or species habitat may occur within area
Reptiles		
Delma torquata Adorned Delma, Collared Delma [1656]	Vulnerable	Species or species habitat may occur within area
Furina dunmalli Dunmall's Snake [59254]	Vulnerable	Species or species habitat may occur within area
Uvidicolus sphyrurus Border Thick-tailed Gecko, Granite Belt Thick-tailed Gecko [84578]	Vulnerable	Species or species habitat known to occur within area
Wollumbinia belli Bell's Turtle, Western Sawshelled Turtle, Namoi River Turtle, Bell's Saw-shelled Turtle [86071]	Vulnerable	Species or species habitat may occur within area
Listed Migratory Species * Species is listed under a different scientific name on the second	the EPBC Act - Threatened	[Resource Information Species list.
Name	Threatened	Type of Presence
Migratory Marine Birds <u>Apus pacificus</u>		
Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Migratory Terrestrial Species		
Hirundapus caudacutus White-throated Needletail [682]		Species or species habitat likely to occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat may occur within area
Myiagra cyanoleuca Satin Flycatcher [612]		Species or species habitat likely to occur within area
Rhipidura rufifrons Rufous Fantail [592]		Species or species habitat likely to occur within area
Migratory Wetlands Species		

Name	Throotopod	Type of Dreserve
Name	Threatened	Type of Presence
Actitis hypoleucos		
Common Sandpiper [59309]		Species or species habitat may occur within area
Calidris acuminata		
Sharp-tailed Sandpiper [874]		Species or species habitat may occur within area
Calidris ferruginea		
Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Calidris melanotos		
Pectoral Sandpiper [858]		Species or species habitat may occur within area
Gallinago hardwickii		
Latham's Snipe, Japanese Snipe [863]		Species or species habitat may occur within area
Pandion haliaetus		
Osprey [952]		Species or species habitat may occur within area

Other Matters Protected by the EPBC Act

Other Matters Protected by the EPBC Act		
Listed Marine Species		[Resource Information]
* Species is listed under a different scientific name on t	he EPBC Act - Threatened	Species list.
Name	Threatened	Type of Presence
Birds		
Actitis hypoleucos		
Common Sandpiper [59309]		Species or species habitat may occur within area
Apus pacificus		
Fork-tailed Swift [678]		Species or species habitat likely to occur within area
Ardea alba		
Great Egret, White Egret [59541]		Species or species habitat likely to occur within area
Ardea ibis		
Cattle Egret [59542]		Species or species habitat may occur within area
Calidris acuminata		
Sharp-tailed Sandpiper [874]		Species or species habitat may occur within area
Calidris ferruginea		
Curlew Sandpiper [856]	Critically Endangered	Species or species habitat may occur within area
Calidris melanotos		
Pectoral Sandpiper [858]		Species or species habitat may occur within area
Chrysococcyx osculans		
Black-eared Cuckoo [705]		Species or species habitat likely to occur within area
Gallinago hardwickii		
Latham's Snipe, Japanese Snipe [863]		Species or species habitat may occur within area

Name	Threatened	Type of Presence
Haliaeetus leucogaster White-bellied Sea-Eagle [943]		Species or species habitat may occur within area
Hirundapus caudacutus White-throated Needletail [682]		Species or species habitat likely to occur within area
<u>Lathamus discolor</u> Swift Parrot [744]	Critically Endangered	Species or species habitat likely to occur within area
Merops ornatus Rainbow Bee-eater [670]		Species or species habitat may occur within area
Motacilla flava Yellow Wagtail [644]		Species or species habitat may occur within area
Myiagra cyanoleuca Satin Flycatcher [612]		Species or species habitat likely to occur within area
Pandion haliaetus Osprey [952]		Species or species habitat may occur within area
Rhipidura rufifrons Rufous Fantail [592]		Species or species habitat likely to occur within area
Rostratula benghalensis (sensu lato) Painted Snipe [889]	Endangered*	Species or species habitat may occur within area

Extra Information

Extra Information							
State and Territory Reserves	[Resource Information]						
Name	State						
Crooked Creek	NSW						
Regional Forest Agreements	[Resource Information]						
Note that all areas with completed RFAs have been included.							
Name	State						
North East NSW RFA	New South Wales						
Invasive Species	[Resource Information]						
Weeds reported here are the 20 species of national significance (WoNS), along with other introduced plants that are considered by the States and Territories to pose a particularly significant threat to biodiversity. The following feral animals are reported: Goat, Red Fox, Cat, Rabbit, Pig, Water Buffalo and Cane Toad. Maps from Landscape Health Project, National Land and Water Resouces Audit, 2001.							

Name	Status	Type of Presence
Birds		
Columba livia		
Rock Pigeon, Rock Dove, Domestic Pigeon [803]		Species or species habitat likely to occur within area
Passer domesticus		
House Sparrow [405]		Species or species

Name	Status Type of Presence
Sturnue vulgarie	habitat likely to occur within area
Sturnus vulgaris Common Starling [389]	Species or species habitat likely to occur within area
Frogs	
Rhinella marina	
Cane Toad [83218]	Species or species habitat may occur within area
Mammals	
Bos taurus	
Domestic Cattle [16]	Species or species habitat likely to occur within area
Canis lupus familiaris	
Domestic Dog [82654]	Species or species habitat likely to occur within area
Capra hircus	
Goat [2]	Species or species habitat likely to occur within area
Felis catus	
Cat, House Cat, Domestic Cat [19]	Species or species habitat likely to occur within area
Feral deer	
Feral deer species in Australia [85733]	Species or species habitat likely to occur within area
Oryctolagus cuniculus	
Rabbit, European Rabbit [128]	Species or species habitat likely to occur within area
Sus scrofa	
Pig [6]	Species or species habitat likely to occur within area
Vulpes vulpes	Species or species habitat
Red Fox, Fox [18]	likely to occur within area
Plants	
Genista sp. X Genista monspessulana	
Broom [67538]	Species or species habitat may occur within area
Nassella neesiana	
Chilean Needle grass [67699]	Species or species habitat likely to occur within area
Opuntia spp.	
Prickly Pears [82753]	Species or species habitat likely to occur within area
Pinus radiata	
Radiata Pine Monterey Pine, Insignis Pine, Wilding Pine [20780]	Species or species habitat may occur within area
Rubus fruticosus aggregate	
Blackberry, European Blackberry [68406]	Species or species habitat likely to occur within area
Salix spp. except S.babylonica, S.x calodendron & S	.x reichardtii
Willows except Weeping Willow, Pussy Willow and Sterile Pussy Willow [68497]	Species or species habitat likely to occur within area
Solanum elaeagnifolium	
Silver Nightshade, Silver leaf Nightshade, Tomata Wood	Species or species habitat
Horse Nettle, Silver-leaf Nightshade, Tomato Weed, White Nightshade, Bull-nettle,	likely to occur within area

Prairie-berry, Satansbos, Silver-leaf Bitter-apple, Silverleaf-nettle, Trompillo [12323]

Caveat

The information presented in this report has been provided by a range of data sources as acknowledged at the end of the report.

This report is designed to assist in identifying the locations of places which may be relevant in determining obligations under the Environment Protection and Biodiversity Conservation Act 1999. It holds mapped locations of World and National Heritage properties, Wetlands of International and National Importance, Commonwealth and State/Territory reserves, listed threatened, migratory and marine species and listed threatened ecological communities. Mapping of Commonwealth land is not complete at this stage. Maps have been collated from a range of sources at various resolutions.

Not all species listed under the EPBC Act have been mapped (see below) and therefore a report is a general guide only. Where available data supports mapping, the type of presence that can be determined from the data is indicated in general terms. People using this information in making a referral may need to consider the gualifications below and may need to seek and consider other information sources.

For threatened ecological communities where the distribution is well known, maps are derived from recovery plans, State vegetation maps, remote sensing imagery and other sources. Where threatened ecological community distributions are less well known, existing vegetation maps and point location data are used to produce indicative distribution maps.

Threatened, migratory and marine species distributions have been derived through a variety of methods. Where distributions are well known and if time permits, maps are derived using either thematic spatial data (i.e. vegetation, soils, geology, elevation, aspect, terrain, etc) together with point locations and described habitat; or environmental modelling (MAXENT or BIOCLIM habitat modelling) using point locations and environmental data layers.

Where very little information is available for species or large number of maps are required in a short time-frame, maps are derived either from 0.04 or 0.02 decimal degree cells; by an automated process using polygon capture techniques (static two kilometre grid cells, alpha-hull and convex hull); or captured manually or by using topographic features (national park boundaries, islands, etc). In the early stages of the distribution mapping process (1999-early 2000s) distributions were defined by degree blocks, 100K or 250K map sheets to rapidly create distribution maps. More reliable distribution mapping methods are used to update these distributions as time permits.

Only selected species covered by the following provisions of the EPBC Act have been mapped:

- migratory and
- marine

The following species and ecological communities have not been mapped and do not appear in reports produced from this database:

- threatened species listed as extinct or considered as vagrants
- some species and ecological communities that have only recently been listed
- some terrestrial species that overfly the Commonwealth marine area
- migratory species that are very widespread, vagrant, or only occur in small numbers

The following groups have been mapped, but may not cover the complete distribution of the species:

- non-threatened seabirds which have only been mapped for recorded breeding sites
- seals which have only been mapped for breeding sites near the Australian continent

Such breeding sites may be important for the protection of the Commonwealth Marine environment.

Coordinates

-29.190623 151.336471,-29.190324 151.336299,-29.190324 151.336299,-29.188525 151.345569,-29.190474 151.34677,-29.193471 151.346255,-29.195269 151.357242,-29.208156 151.357242,-29.208605 151.348659,-29.212651 151.349345,-29.212651 151.331493,-29.205459 151.331664,-29.205459 151.336642,-29.190623 151.336471

Acknowledgements

This database has been compiled from a range of data sources. The department acknowledges the following custodians who have contributed valuable data and advice:

- -Office of Environment and Heritage, New South Wales
- -Department of Environment and Primary Industries, Victoria
- -Department of Primary Industries, Parks, Water and Environment, Tasmania
- -Department of Environment, Water and Natural Resources, South Australia
- -Department of Land and Resource Management, Northern Territory
- -Department of Environmental and Heritage Protection, Queensland
- -Department of Parks and Wildlife, Western Australia
- -Environment and Planning Directorate, ACT
- -Birdlife Australia
- -Australian Bird and Bat Banding Scheme
- -Australian National Wildlife Collection
- -Natural history museums of Australia
- -Museum Victoria
- -Australian Museum
- -South Australian Museum
- -Queensland Museum
- -Online Zoological Collections of Australian Museums
- -Queensland Herbarium
- -National Herbarium of NSW
- -Royal Botanic Gardens and National Herbarium of Victoria
- -Tasmanian Herbarium
- -State Herbarium of South Australia
- -Northern Territory Herbarium
- -Western Australian Herbarium
- -Australian National Herbarium, Canberra
- -University of New England
- -Ocean Biogeographic Information System
- -Australian Government, Department of Defence
- Forestry Corporation, NSW
- -Geoscience Australia
- -CSIRO
- -Australian Tropical Herbarium, Cairns
- -eBird Australia
- -Australian Government Australian Antarctic Data Centre
- -Museum and Art Gallery of the Northern Territory
- -Australian Government National Environmental Science Program
- -Australian Institute of Marine Science
- -Reef Life Survey Australia
- -American Museum of Natural History
- -Queen Victoria Museum and Art Gallery, Inveresk, Tasmania
- -Tasmanian Museum and Art Gallery, Hobart, Tasmania
- -Other groups and individuals

The Department is extremely grateful to the many organisations and individuals who provided expert advice and information on numerous draft distributions.

Please feel free to provide feedback via the Contact Us page.

GAIA BONSHAW SOLAR EIS Revised Biodiversity Developmen	nt Assessment Report
APPENDIX D	LIKELIHOOD OF OCCURRENCE AND RISK ANALYSIS

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

In order to complement identification of likely threatened species likely to occur within the Development Site, the NSW BioNet and the PMST were undertaken. A likelihood of occurrence and risk assessments were undertaken for all species. Where results identify additional species to those generated in the BCD Calculator as "Ecosystem Species Credits" or "Species Credit Species", those species were added in the surveys design.

Likelihood of Occurrence

Species identified in the NSW BioNet atlas and the PMST were collated into a table where an assessment of the likelihood of occurrence of that threatened biodiversity was undertaken. In making this determination, the following factors were considered:

- habitat quality within and adjacent to the Development Site as determined through review of regional vegetation mapping and the results of the September survey;
- breeding habitat/resources present assists with identification of the importance of habitat to the species;
- dispersal ability based on known ecology whether the species have an ability to disperse to new areas of habitat following disturbance; and
- local records in similar habitat/distance/connectivity to the Development Site.

This allows for assessment of cryptic or seasonal species that are unlikely to be readily identified during brief site inspections and/or due to seasonal constraints. The likelihood of each species occurring was categorised as known, potential or unlikely to occur based on the definitions provided in Table D-1. Results of likelihood of occurrence is presented in Table D.3 for TECs and Table D.4 for threatened species.

Table D.1 Definitions of Likelihood of Occurrence

Category	Description
Known	 the ecological community/species/matter has been recorded in the Development Site during field surveys; or database records demonstrate that the ecological community/species has been known to occur in the Development Site within the last 10 year period.
Potential	 the ecological community/species' known distribution includes the Development Site, and suitable habitat is present within it, or, database records demonstrate that the ecological community/species has been known to occur in the Development Site, however has not been recorded within the last 10 years, or the species is a wide ranging flying species which may 'fly-over' the Development Site, regardless of the habitat types present and has been recorded within the 10 km locality surrounding the Development Site.
Unlikely	 the ecological community/species has not been recorded within 10 km locality of the Development Site and suitable habitat does not occur within the Development Site, or the Development Site is not within the TEC/species' known distribution, or sufficient field surveys have been conducted to conclude that the species is likely to be absent.

The following considerations were made in assessing habitat suitability and distribution: Habitat quality within and adjacent to the Site.

Breeding habitat/resources present – assists with identification of the importance of habitat to the species. The species' ability to disperse to new areas of habitat following disturbance.

Local records in similar habitat/distance/connectivity to the Site.

www.erm.com Version: 1.0 Project No.: 0470861 Client: GAIA Australia 26 July 2019 Page D1

Risk Assessment

A risk assessment was undertaken using the definitions of Species Sensitivity and Consequence to assign a relative risk ranking for each listed ecological value (Low, Medium, High or Very High, as shown in Table D.2). Impacts to ecological values with potential to occur that were assessed as having a Low risk was not further assessed. Results of risk assessment is presented in Table D.3 for TECs and Table D.4 for threatened species.

Impacts to ecological values with potential to occur that were assessed as having a Medium, High or Very High risk were further assessed in accordance with the requirements of the EPBC Act and BC Act, including the preparation of detailed Assessments of Significance (see Appendix E).

Species sensitivity rankings are based on the species conservation status under the EPBC Act, FM Act and BC Act. Where the conservation status differs between listings, the conservation status with higher sensitivity is used.

Table D.2 Risk Assessment Matrix

	Consequence							
		Negligible	Minor	Moderate	Major			
	Ecological value not listed as threatened	Low	Low	Medium	High			
Sensitivity	Ecological value listed as Vulnerable or Migratory	Low	Medium	Medium	High			
Se	Ecological value listed as Endangered	Low	Medium	High	Very High			
	Ecological value listed as Critically Endangered	Medium	High	Very High	Very High			

Consequence Definitions

Negligible: No impacts or removal of ecological community. Effect on species is within the likely normal

range of variation. No removal of specific breeding habitat features.

Minor: Indirect impacts to listed ecological community which may affect a small proportion of the

ecological community. Effects a small proportion of a population and Project-related mortality of a small number of individuals may occur, but does not substantially affect other species dependent on it, or the populations of the species itself. No removal of specific breeding habitat features.

on it, or the populations of the openior from the removal of opening habitat features.

Moderate: Direct removal of a portion of a listed ecological community. Effects a sufficient proportion of a

species population that it may bring about a substantial change in abundance and/or reduction in distribution over one or more generations, but does not threaten the long term viability of that

population or any population dependent on it.

Major: Direct removal of a listed ecological community. Effects an entire population or species at

sufficient scale to cause a substantial decline in abundance and/or change in distribution beyond with natural recruitment (reproduction, immigration from unaffected areas) may not return that population or species, or any population or species dependent upon it, to its former level within

several generations, or when there is no possibility of recovery.

Species sensitivity definitions

Species sensitivities refer to listed under either the EPBC Act or BC Act. Where listings differ, the higher sensitivity is used.

riigher concilivity is acca.

www.erm.com Version: 1.0 Project No.: 0470861 Client: GAIA Australia 26 July 2019 Page D2

Table D.3 Threatened Ecological Communities: Likelihood of Occurrence and Risk Assessment

TEC name	EPBC Act	BC Act	Community Description	Likely of TEC Occurring in the Site	Potential Impacts	Mitigation Measures	Risk Rating
Natural Grasslands on Basalt and Fine- textured alluvial plains of northern New South Wales and southern Queensland Source: PMST	CE		The Natural grasslands on basalt and fine-textured alluvial plains of northern New South Wales (NSW) and southern Queensland are native grasslands typically composed of perennial native grasses. They are found on soils that are fine textured (often cracking clays) derived from either basalt or alluvium on flat to low slopes (< 1 degree). A tree canopy is usually absent, but when present, comprises ≤ 10% projective foliage cover. The distribution of the ecological community is strongly reliant on soil type as it is associated with fine textured, often cracking clays derived from either basalt or quaternary alluvium. Temperate grasses and grassland forbs comprise the ground layer. Tussock grasses within this vegetation community can be dominated by species of the genera Austrodanthonia, Austrostipa, Bothriochloa, Chloris, Enteropogon, or Themeda. In the Darling Downs component of the community, Bluegrass (Dichanthium sericeum) tends to dominate, whereas the Plains Grass (Austrostipa aristiglumis) tend to dominate in the Liverpool Plains component. The herbaceous cover includes species within genera Desmodium, Glycine, Lotus and Rhynchosia. The shrub layer is generally a minimum component of the community, it can include Mimosa (Acacia farnesiana), Pittosporum phylliraeoides, Pimelea spp. and Sclerolaena spp. A tree canopy is typically absent. Where trees are present, they are of variable species composition and comprise less than 10% of projective crown cover. Tree species that may be present as scattered individuals include: Acacia pendula (Weeping Myall), Eucalyptus albens (White Box), E. conica (Fuzzy Box), E. coolabah (Coolabah), E. melliodora (Yellow Box), E. populnea (Poplar Box) or E. tereticornis (Forest Red Gum). This ecological community occurs from the Darling Downs in Queensland to Dubbo in NSW and incorporates the Liverpool and Moree Plains. This ecological community occurs within the Brigalow Belt South Bioregion and Border Rivers-Gwydir, Central West, Namoi, Condamine, Burnett Mary and Fitzroy Ba	Unlikely The Development Site is located immediately south form the Darling Downs, Queensland. Cracking clays were observed in the southern portion of the Development Site, however, the site is highly disturbed and the characteristic species in the northern portion of the distribution of this TEC, i.e. Queensland Bluegrass (<i>Dichanthium sericeum</i>) were not recorded. Therefore, it is considered unlikely this CEEC occurs.	NA	NA	Low
New England Peppermint (Eucalyptus nova- anglica) Grassy Woodlands Source: PMST	CE	CE	The New England Peppermint (<i>Eucalyptus nova-anglica</i>) Grassy Woodlands ecological community occurs in northern NSW and southern Queensland, in the New England Tablelands, NSW North Coast and Nandewar IBRA Bioregions. The tree canopy is typically dominated (>50%) or co-dominated (>30%) by the tree species <i>Eucalyptus nova-anglica</i> (New England Peppermint). A range of other associated tree species may be present, and may be co-dominant in the ecological community, but do not dominate it by themselves, in particular <i>E. pauciflora</i> (Snow Gum) and <i>E. dalrympleana</i> subsp. <i>heptantha</i> (Mountain Gum). The understorey is usually made up of a dense, species-rich ground layer of grasses and herbs. Shrubs are typically sparse to absent. The main tree species in the community are New England Peppermint (<i>Eucalyptus nova-anglica</i>). Other tree species include Snow Gum (<i>E. pauciflora</i>), Black Sallee (<i>E. stellulata</i>), Mountain Gum (<i>E. dalrympleana subsp. heptantha</i>), Blakely's Red Gum (<i>E. blakelyi</i>) and Fuzzy Box (<i>E. conica</i>).	Unlikely No tree elements characteristic of the vegetation community were observed within the Development Site during surveys.	NA	NA	Low
Weeping Myall Woodlands	Е		The Weeping Myall Woodlands ecological community occurs on the inland alluvial plains west of the Great Dividing Range in NSW and Queensland. It occurs in the Riverina, NSW South Western Slopes, Darling Riverine Plains, Brigalow Belt South, Brigalow Belt North, Murray-Darling Depression, Nandewar and Cobar	Unlikely No elements characteristic of the vegetation community were observed within the Development Site during surveys.	NA	NA	Low

TEC name	EPBC Act	BC Act	Community Description	Likely of TEC Occurring in the Site	Potential Impacts	Mitigation Measures	Risk Rating
Source: PMST			Peneplain IBRA Bioregions. The ecological community currently occurs in small pockets throughout this range. The Weeping Myall Woodlands occur in a range from open woodlands to woodlands, generally 4-12 m high, in which Weeping Myall (<i>Acacia pendula</i>) trees are the sole or dominant overstorey species. Weeping Myall trees often occur in monotypic stands, however other vegetation may also occur in the ecological community, though not as dominant species. These include: Western Rosewood (<i>Alectryon oleifolius subsp. elongatus</i>); Poplar Box (<i>Eucalyptus populnea</i>); or Black Box (<i>Eucalyptus largiflorens</i>). Grey Mistletoe (<i>Amyema quandang</i>) commonly occurs on the branches of Weeping Myall trees throughout the ecological community's range. Weeping Myall goes through regular cycles of senescence (aging and death) and regeneration. Weeping Myall trees are also susceptible to defoliation by Bagshelter Moth (<i>Ochrogaster lunifer</i>) caterpillars and are often lopped for domestic stock fodder. Therefore, the ecological community can be dominated by Weeping Myall trees that are in a living, defoliated or dead state. The understorey of Weeping Myall Woodlands often includes an open layer of shrubs above an open ground layer of grasses and herbs, though the ecological community can exist naturally either as a shrubby or a grassy woodland.				
White Box – Yellow Box – Blakely's Red Gum Grassy Woodland and Derived Native Grassland Source: PMST	CE	E	Box – Gum Grassy Woodlands and Derived Grasslands are characterised by a species-rich understorey of native tussock grasses, herbs and scattered shrubs, and the dominance, or prior dominance, of White Box, Yellow Box or Blakely's Red Gum trees. In the Nandewar Bioregion, Grey Box (<i>Eucalyptus microcarpa</i> or <i>E. moluccana</i>) may also be dominant or codominant. The tree-cover is generally discontinuous and consists of widely-spaced trees of medium height in which the canopies are clearly separated. Associated, and occasionally co-dominant, trees include, but are not restricted to: Grey Box (<i>Eucalyptus microcarpa</i>), Fuzzy Box (<i>E. conica</i>), Apple Box (<i>E. bridgesiana</i>), Red Box (<i>E. polyanthemos</i>), Red Stringybark (<i>E. macrorhyncha</i>), White Cypress Pine (<i>Callitris glaucophylla</i>), Black Cypress Pine (<i>C. enderlicheri</i>), Long-leaved Box (<i>E. gonicalyx</i>), New England Stringybark (<i>E. calignosa</i>), Brittle Gum (<i>E. mannifera</i>), Candlebark (<i>E. rubida</i>), Argyle Apple (<i>E. cinerea</i>), Kurrajong (<i>Brachychiton populneus</i>) and Drooping She-oak (<i>Allocasuarina verticillata</i>). This ecological community occurs in areas where rainfall is between 400 and 1200 mm per annum, on moderate to highly fertile soils at altitudes of 170 metres to 1200 metres. The White Box – Yellow Box – Blakely's Red Gum grassy woodlands that existed prior to European settlement now exists as remnants in three different states. The three states are: i) An overstorey of eucalypt trees exists, but there is no substantial native understorey. ii) A native understorey exists, but there is no substantial native understorey ii) A native understorey exists, but there shave been cleared; and iii) Both a native understorey and an overstorey of eucalypts exist in conjunction. In order for an area to be included in the EPBC listed ecological community, a patch must have a predominantly native understorey. The size and life-form of understorey species are such that viable populations can exist in very small areas. Therefore, in order to be the listed ecological	Potential An area likely to conform to the Box-Gum community was recorded in the southern portion of the Subject Land. An individual tree, potentially identified as Blakely's Red Gum (Eucalyptus blakely), was observed during the September 2018 survey. The tree was later confirmed as Eucalyptus dealbata by ecologists Joanne Woodhouse and Tom Cotter. No elements of this vegetation community were found across the Development Site although it is likely to occur within the broader Project Boundary.	Negligible	Avoidance	Low

Table D.4 Threatened Species: Likelihood of Occurrence and Risk Assessment

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Fauna Reptilia							
Delma torquata Adorned Delma, Collared Delma Source: PMST		V	Adorned Delma is the smallest of the legless lizards. The Collared Delma normally inhabits eucalypt dominated woodland and open forest where it is associated with suitable micro-habitats (exposed rocky outcrops). The ground cover is predominantly native grasses, such as Kangaroo Grass (<i>Themeda triandra</i>), Barbed-wire Grass (<i>Cymbopogon refractus</i>), Wiregrass (<i>Aristida</i> sp.) and <i>Lomandra</i> . The Collared Delma is known from the western suburbs of Brisbane, Queensland, and the following sites: Bunya Mountains, Blackdown Tableland National Park (NP), Bullyard Conservation Park, D'Aguilar Range NP, Expedition NP, Naumgna and Lockyer Forest Reserves, Western Creek near Millmerran and the Toowoomba Range. It occurs within the South East Queensland, Condamine, Burnett Mary and Fitzroy (Queensland) Natural Resource Management regions.	Unlikely No records of the species are known within a 10km radius of the Site. The species was not recorded during target reptile surveys within the rocky outcrops in December 2018.	NA NA	NA NA	Low
Furina dunmalli Dunmall's Snake		V	Dunmall's Snake preferred habitat is Brigalow forest and woodland with fallen timber and ground litter, growing on cracking clay soils and clay loam soils. It also occurs in eucalypt and <i>Callitris</i> woodland with fallen timber and ground litter. The species is nocturnal. Dunmall's Snake occurs in south-east interior of Queensland, including the Darling Downs, and is thought to potentially extend into inland north-eastern NSW. Most locality records are between 200 and 500 m elevation	Unlikely No records of the species are known within a 10km radius of the Site. The species was not recorded during target reptile surveys in December 2018.	NA	NA	Low
Uvidicolus sphyrurus Border Thick-tailed Gecko Source: BioNet, PMST	V	V	The Border Thick-tailed Gecko is a small lizard up to 10 cm long (average 7 cm). It often occurs on steep rocky or scree slopes, especially granite. Recent records from basalt and metasediment slopes and flats indicate its habitat selection is broader than formerly thought and may have extended into areas that were cleared for agriculture. Favours forest and woodland areas with boulders, rock slabs, fallen timber and deep leaf litter. Occupied sites often have a dense tree canopy that helps create a sparse understorey. These Geckos are active at night and shelter by day under rock slabs, in or under logs, and under the bark of standing trees. It is found only on the tablelands and slopes of northern NSW and southern Queensland, reaching south to Tamworth and west to Moree. Most common in the granite country of the New England Tablelands. Occurs at sites ranging from 500 to 1100 m elevation. Populations are mostly fragmented, with over 50 discrete sites currently known that are separated by at least 2km.	Potential Habitat resources for the species are present within the Development Site. The only record within the 10km locality is at approximately 2.4km south from the site and was recorded over 10 years ago (in 1999) The species was not recorded during target reptile surveys within the rocky outcrops in December 2018.	Negligible A total of eleven rocky areas were recorded within the Subject Land. Five of the rocky outcrops have been avoided. Only two outcrops occur in the middle of the cleared paddocks and cannot be avoided by the development footprint. The remaining four may also be avoided during the detailed design and survey of the required riparian buffers and perimeter road.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation and habitat features to be retained within the adjacent habitat.	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Wollumbinia belli Bell's Turtle	E	V	Bell's Turtle occurs in shallow to deep pools in upper reaches or small tributaries of major rivers in granite country. Occupied pools are most commonly less than 3 m deep with rocky or sandy bottoms and patches of vegetation. Most typically uses narrow stretches of rivers 30 - 40 m wide. Most surrounding habitat has been converted to grazing land. Nests are dug out in riverbanks of sand or loam during late September to January. Eggs take 80 days to hatch and are thus vulnerable to nest predation for an extended period. Primarily a vegetarian, eating both aquatic plants and terrestrial leaves that fall into the watercourse. Also takes invertebrates ranging from insects to crayfish, other small animals and carrion. In NSW, currently found in four disjunct populations in the upper reaches of the Namoi, Gwydir and Border Rivers systems, on the escarpment of the North West Slopes.	Unlikely No deep pools occur within any of the unnamed creeks or their tributaries.	Negligible	NA .	Low
Aves							
Anthochaera phrygia Regent Honeyeater Source: PMST	CE	CE	The Regent Honeyeater mainly inhabits temperate woodlands and open forests, particularly Box – Ironbark woodland and riparian forests of River Sheoak. The species inhabits woodlands that support a significantly high abundance and species richness of birds. These type of woodlands have significantly large numbers of mature trees, high canopy cover and abundance of mistletoes. The species can also be found in drier coastal woodlands and forests in some years. Non-breeding flocks of the species can be seen foraging in flowering coastal Swamp Mahogany and Spotted Gum forests. Although the species is a generalist forager, it feeds mainly on the nectar from a small number of eucalypts that produce high volumes of nectar (e.g. Mugga Ironbark, Yellow Box, White Box and Swamp Mahogahy).	Potential Limited foraging habitat is present within the woodland and riparian corridor . This woodland bird species forages in woodlands with significantly large numbers of mature trees, high canopy cover and abundance of mistletoes. No habitat is available within the disturbed grasslands. The Regent Honeyeater has not been recorded within the Site.	Negligible. The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation and habitat features to be retained within the adjacent habitat.	Low
Artamus cyanopterus cyanopterus Dusky Woodswallow	V		The Dusky Woodswallow primarily inhabit dry, open eucalypt forests and woodlands, including mallee associations, with an open or sparse understorey of eucalypt saplings, acacias and other shrubs, and ground-cover of grasses or sedges and fallen woody debris. It has also been recorded in shrublands, heathlands and very occasionally in moist forest or rainforest. Also found in farmland, usually at the edges of forest or woodland. The species forages on invertebrates, mainly insects, which are captured whilst hovering or sallying above the canopy or over water. It builds an open, cup-shape nest made of twigs, grass, fibrous rootlets and occasionally casuarina needles. Generally, nests are located on shrubs or low trees, living or dead, horizontal or upright forks in braches, spouts, hollow stumps or logs, behind loose bark or in a hollow in the top of a wooden fence post.	Potential. Potential foraging habitat is present in most woodland vegetation zones, excluding grasslands, cleared and disturbed land No records of the species exist for the Site and the nearest location is at approximately 4 km south-east where it was recorded in 1995.	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	NA NA	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Calyptorhynchus lathami Glossy Black-Cockatoo Source: BioNet	V		The Glossy Black-cockatoo inhabits open forest and woodlands of the coast and the Great Dividing Range where stands of sheoak occur. Black Sheoak (Allocasuarina littoralis) and Forest Sheoak (A. torulosa) are important foods. Inland populations feed on a wide range of sheoaks, including Drooping Sheoak, Allocasuaraina diminuta, and A. gymnathera. Belah is also utilised and may be a critical food source for some populations. In the Riverina, birds are associated with hills and rocky rises supporting Drooping Sheoak, but also recorded in open woodlands dominated by Belah (Casuarina cristata). Feeds almost exclusively on the seeds of several species of she-oak (Casuarina and Allocasuarina species), shredding the cones with the massive bill.	Unlikely. Limited breeding habitat for the species is present within the Development Site. However, no suitable feeding habitat is present as its preferred feed tree species have not been recorded within the Site. It is unlikely the sepcies would breed in hollow-bearing trees at the site as the cost to travel long distances to feed will be too high. The nearest location is at approximately 3.6 km to the east, where one record from 1999 exists.	Negligible	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Climacteris picumnus victoriae Brown Treecreeper (eastern subspecies) Source: BioNet	V		The Brown Treecreeper (eastern subspecies) is found in eucalypt woodlands (including Box-Gum Woodland) and dry open forest of the inland slopes and plains inland of the Great Dividing Range; mainly inhabits woodlands dominated by stringybarks or other rough-barked eucalypts, usually with an open grassy understorey, sometimes with one or more shrub species; also found in mallee and River Red Gum (<i>Eucalyptus camaldulensis</i>) Forest bordering wetlands with an open understorey of acacias, saltbush, lignum, cumbungi and grasses; usually not found in woodlands with a dense shrub layer; fallen timber is an important habitat component for foraging; also recorded, though less commonly, in similar woodland habitats on the coastal ranges and plains. It is considered resident of areas where it occurs and is usually observed in pairs or small groups of 8 to 12 birds. It forages on trunks and branches of trees and among fallen timber. Hollows in standing dead or live trees and stumps are essential for nesting.	Potential This species was recorded adjacent to the Subject Land during potential offset site investigations. Suitable habitat for the species is considered to occur in the woodland PCTs. Grasslands and cleared land (disturbed grasslands) do not represent suitable habitat for the species	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Chthonicola sagittata Speckled Warbler Source: BioNet	V		The Speckled Warbler lives in a wide range of Eucalyptus dominated communities that have a grassy understorey, often on rocky ridges or in gullies. Typical habitat would include scattered native tussock grasses, a sparse shrub layer, some eucalypt regrowth and an open canopy. Large, relatively undisturbed remnants are required for the species to persist in an area. The diet consists of seeds and insects, with most foraging taking place on the ground around tussocks and under bushes and trees. Pairs are sedentary and occupy a breeding territory of about ten hectares, with a slightly larger home-range when not breeding. The rounded, domed, roughly built nest of dry grass and strips of bark is located in a slight hollow in the ground or at the base of a low dense plant, often among fallen branches and other litter. A side entrance allows the bird to walk directly inside. A clutch of 3-4 eggs is laid, between August and January, and both parents feed the nestlings. Speckled Warblers often join mixed species feeding flocks in winter, with other species such as Yellow-rumped, Buff-rumped, Brown and Striated Thornbills.	Potential This species was recorded adjacent to the Subject Land during potential offset site investigations. Suitable habitat for the species is considered to occur in the woodland PCTs. Grasslands and cleared land (disturbed grasslands) do not represent suitable habitat for the species.	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low

Biodiversity Development Assessment Report

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Daphoenositta chrysoptera Varied Sittella Source: BioNet	V		The Varied Sittella Inhabits eucalypt forests and woodlands, especially those containing rough-barked species and mature smooth-barked gums with dead branches, mallee and <i>Acacia</i> woodland. It feeds on arthropods gleaned from crevices in rough or decorticating bark, dead branches, standing dead trees and small branches and twigs in the tree canopy. It builds a cup-shaped nest of plant fibres and cobwebs in an upright tree fork high in the living tree canopy, and often re-uses the same fork or tree in successive years. Generation length is estimated to be 5 years. The Varied Sittella is sedentary and inhabits most of mainland Australia except the treeless deserts and open grasslands. Distribution in NSW is nearly continuous from the coast to the far west.	Potential Potential foraging habitat is present in most woodland vegetation zones. The nearest record of the species dates back in 1995 and was located at approximately 4 km south from the Site.	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Erythrotriorchis radiatus Red Goshawk Source: PMST	CE	V	Red Goshawks inhabit open woodland and forest, preferring a mosaic of vegetation types, a large population of birds as a source of food, and permanent water, and are often found in riparian habitats along or near watercourses or wetlands. In NSW, preferred habitats include mixed subtropical rainforest, Melaleuca swamp forest and riparian Eucalyptus forest of coastal rivers.	Unlikely Limited habitat for the species exists within the Development Site. Riparian corridor was dry at the time of surveys and highly fragmented. No existing records of the species within the 10 km locality exist. The species was not recorded during surveys.	Negligible	NA	Low
Geophaps scripta scripta Squatter Pigeon Source: PMST	CE	V	Squatter Pigeons are medium-sized ground-dwelling pigeons. They are found in grassy woodlands and plains, preferring sandy areas and usually close to water. Feed on the ground, on seeds of grasses, herbs and shrubs, as well as insects. Nest on the ground. The species is found from north Queensland to the North West Slopes of NSW and extending down to the Liverpool Plains and Dubbo. Today they are very rare in the southern parts of their range.	Potential Substandard foraging habitat is present within the Development Site. No existing records of the species within the 10 km locality exist. The species was not recorded during surveys.	Negligible	NA	Low
Glossopsitta pusilla Little Lorikeet Source: BioNet	V		The Little Lorikeet is a small parrot distributed widely across the coast and Great Divide regions. The species forages primarily in the canopy of open Eucalyptus forests and woodland. It also forages in Angophora, Melaeuca and other species including paddock, roadside remnants and urban trees. It feeds mainly on nectar and pollen, occasionally on native fruits.	Potential. Limited suitable foraging habitat for the species is present in Eucalypt trees within the remnant trees and vegetation. The nearest records of the species dates back in 1995 (two records) and were located at approximately 1 km east from the Site.	Negligible. Any clearing of Eucalypt trees has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Develop fauna clearance protocol that includes procedures to be followed should any injured fauna be encountere	Low

www.erm.com Version: 1.0 Project No.: 0470861 Client: GAIA Australia 26 July 2019 Page D8

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Grantiella picta Painted Honeyeater Source: PMST	V	V	The Painted Honeyeater inhabits Boree/ Weeping Myall (<i>Acacia pendula</i>), Brigalow (<i>A. harpophylla</i>) and Box-Gum Woodlands and Box-Ironbark Forests. A specialist feeder on the fruits of mistletoes growing on woodland eucalypts and acacias. Prefers mistletoes of the genus <i>Amyema</i> . Insects and nectar from mistletoe or eucalypts are occasionally eaten. Nest from spring to autumn in a small, delicate nest hanging within the outer canopy of drooping eucalypts, she-oak, paperbark or mistletoe branches.	Potential Limited foraging resources for the species are available in Box and Ironbark trees across the Subject Land. No existing records of the species within the 10 km locality exist. The species was not recorded during surveys.	Negligible. Any clearing of Eucalypt trees has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Neophema pulchella Turquoise Parrot Source: BioNet	V		The Turquoise Parrot's range extends from southern Queensland through to northern Victoria, from the coastal plains to the western slopes of the Great Dividing Range (OEH 2019). The Turquoise Parrot lives on the edges of eucalypt woodland adjoining clearings, timbered ridges and creeks in farmland. Usually seen in pairs or small, possibly family, groups and have also been reported in flocks of up to thirty individuals. It prefers to feed in the shade of a tree and spends most of the day on the ground searching for the seeds or grasses and herbaceous plants, or browsing on vegetable matter. Forages quietly and may be quite tolerant of disturbance. However, if flushed it will fly to a nearby tree and then return to the ground to browse as soon as the danger has passed. It nests in tree hollows, logs or posts, from August to December. It lays four or five white, rounded eggs on a nest of decayed wood dust (OEH 2019).	Potential Suitable habitat for the species includes woodlands and native grasslands. Roosting and breeding habitat for this woodland bird are hollowbearing trees in woodlands. It forages in woodlands and native grasslands. The nearest record of the species dates back in 1995 and was located at approximately 1 km south-east from the Site.	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Ninox connivens Barking Owl	V		The Barking Owl is found throughout continental Australia except for the central arid regions. Inhabits woodland and open forest, including fragmented remnants and partly cleared farmland. It is flexible in its habitat use, and hunting can extend in to closed forest and more open areas. Sometimes able to successfully breed along timbered watercourses in heavily cleared habitats (e.g. western NSW) due to the higher density of prey on these fertile riparian soils. Roost in shaded portions of tree canopies, including tall midstorey trees with dense foliage such as Acacia and Casuarina species. During nesting season, the male perches in a nearby tree overlooking the hollow entrance. Requires very large permanent territories in most habitats due to sparse prey densities. Monogamous pairs hunt over as much as 6000 hectares, with 2000 hectares being more typical in NSW habitats (OEH 2019).	Potential Hollow bearing trees occur within woodland habitats and also as isolated trees within highly disturbed vegetation zones on the Subject Land.	Negligible. Any clearing of Eucalypt trees has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Petroica boodang Scarlet Robin Source: BioNet	V		The Scarlet Robin lives in dry eucalypt forests and woodlands. The understorey is usually open and grassy with few scattered shrubs. This species lives in both mature and regrowth vegetation. It occasionally occurs in mallee or wet forest communities, or in wetlands and tea-tree swamps.o Its habitat usually contains abundant logs and fallen timber: these are important components of its habitat. It breeds on ridges, hills and foothills of the western slopes, the Great Dividing Range and eastern coastal regions; this species is occasionally found up to 1000 metres in altitude. It breeds between July and January. In autumn and winter many Scarlet Robins live in open grassy woodlands, and grasslands or grazed paddocks with scattered trees. It builds nests in the fork of branches, usually more than 2 metres above the ground.	Potential Limited suitable habitat for the species occurs in woodlands with fallen logs and timber in the Subject Land. The nearest record of the species dates back in 1995 and was located at approximately 4 km south-east from the Site.	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Pomatostomus temporalis temporalis Grey-crowned Babbler (eastern subspecies) Source: BioNet	V		The Grey-crowned Babbler inhabits open Box-Gum Woodlands on the slopes, and Box-Cypress-pine and open Box Woodlands on alluvial plains. Woodlands on fertile soils in coastal regions. Flight is laborious so birds prefer to hop to the top of a tree and glide down to the next one. Birds are generally unable to cross large open areas. It lives in family groups that consist of a breeding pair and young from previous breeding seasons. A group may consist of up to fifteen birds. All members of the family group remain close to each other when foraging. It is insectivorous and it forages on the trunks and branches of eucalypts and other woodland trees or on the ground, digging and probing amongst litter and tussock grasses. It builds nests that are used as dormitory and roosting and uses them all year round. It breeds between July and February. Territory ranges from one to 50 hectares (usually ten hectares) and are defended all year. Grey-crowned Babblers are communal breeders that form a family group, in which offspring from the previous season and other unrelated birds help to raise the current's year's brood. In some populations, breeding success is related to the number of helpers. Young birds staty with the family group for at least one year after fledging and may remain for two or more years acting as non-breeding helpers. As breeding spaces become available in the population, some helpers may disperse to establish their own breeding group. Population viability studies in Victoria suggests that a viable population is likely to contain more than ten family groups, while populations with less than ten family groups are likely to have high rate of extinction.	Known A breeding population of Greycrowned have been recorded on site. The nearest BioNet record of the species dates back in 2010 and was located at approximately 500 m east from the Development Site.	Moderate A breeding population of Grey-crowned have been recorded on site. The movement of this species is largely confined to the riparian corridors as this species has limited flying capacity. These corridors (with a 20-30m buffer) have been avoided although nests will be removed within the development footprint	Approved clearing limits to be clearly delineated with temporary fencing or similar prior to construction commencing. Removal of trees with nests will be included in the vegetation clearing protocol including any seasonal constraints to avoid impacting any juveniles or unfledged chicks. Removal of trees with nests will be supervised by an experienced fauna catcher or ecologist. A portion of felled trees will be salvaged as habitat for fauna and translocated in suitable areas in the remainder of the Project Boundary.	Medium

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Rostratula australis Australian Painted-snipe Source: PMST	E	E	The Australian Painted Snipe is small freshwater wader. Prefers fringes of swamps, dams and nearby marshy areas where there is a cover of grasses, lignum, low scrub or open timber. Nests on the ground amongst tall vegetation, such as grasses, tussocks or reeds. The nest consists of a scrape in the ground, lined with grasses and leaves. Breeding is often in response to local conditions; generally occurs from September to December. Incubation and care of young is all undertaken by the male only. Forages nocturnally on mud-flats and in shallow water. Feeds on worms, molluscs, insects and some plant-matter.	Unlikely No preferred habitat is available. No existing records of the species within the 10 km locality exist. The species was not recorded during surveys.	NA	NA	Low
Stagonopleura guttata Diamond Firetail Source: BioNet	V		Diamond Firetails are found in open grassy woodland, heath and farmland or grassland with scattered trees. Diamond Firetails feed on the ground and generally eat ripe or partially ripe seeds and can be seen hopping around on the ground. They occasionally eat insects and their larvae. The Diamond Firetail builds a nest with green grass blades and stems and lines it with fine grasses and feathers. The nest can be found in trees and shrubs with dense foliage and has sometimes been known to build in the base of a hawk's nest.	Potential Potential roosting and breeding habitat for the species occurs in woodlands, foraging habitat for the species occurs across the entire Subject Land The nearest BioNet record of the species dates back in 1995 and was located at approximately 3 km south-east from the Site.	Negligible The second and third order watercourses will be avoided through detailed design. Riparian buffers will be applied to either side of the streams, measured from the edge of the high bank. The distances applied to this BDAR are 20m either side of the second order stream and 30m either side of the third order streams. This will significantly reduce the risk of the potential impacts.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Tyto novaehollandiae Masked Owl	V	-	Extends from the coast where it is most abundant to the western plains. Overall records for this species fall within approximately 90% of NSW, excluding the most arid north-western corner. It lives in dry eucalypt forests and woodlands from sea level to 1100 m. A forest owl, but often hunts along the edges of forests, including roadsides. The typical diet consists of tree-dwelling and ground mammals, especially rats. Pairs have a large homerange of 500 to 1000 hectares. Roosts and breeds in moist eucalypt forested gullies, using large tree hollows or sometimes caves for nesting(OEH 2019).	Potential Hollow bearing trees occur within woodland habitats and also as isolated trees within highly disturbed vegetation zones on the Subject Land.	Negligible. Any clearing of Eucalypt trees has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Calidris ferruginea Curlew Sandpiper Source: PMST	E	CE, Mi	The Curlew Sandpiper is a migratory bird inhabiting coastal habitats and sometimes freshwater wetlands. It also occurs in non-tidal swamps, lakes and lagoons on the coast and sometimes inland.	Unlikely No preferred habitat is available. No existing records of the species within the 10 km locality exist. The species was not recorded during surveys.	NA	NA	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Mammals							
Dasyurus maculatus Spotted-tailed Quoll Source: PMST	V	E	The Spotted-tailed Quoll is recorded across a range of habitat types, including rainforest, open forest, woodland, coastal heath and inland riparian forest, from the sub-alpine zone to the coastline. Individual animals use hollow-bearing trees, fallen logs, small caves, rock outcrops and rocky-cliff faces as den sites. Females occupy home ranges up to about 750 hectares and males up to 3500 hectares. Are known to traverse their home ranges along densely vegetated creeklines	Unlikely The high level of disturbance at the Development Site suggests no suitable habitat for the species is present. No known records of the species exists within the 10 km locality. The species was not recorded during surveys in September and December 2018.	NA	NA	Low
Petaurus norfolcensis Squirrel Glider Source: BioNet	V		The species is widely though sparsely distributed in eastern Australia, from northern Queensland to western Victoria. Inhabits mature or old growth Box, Box-Ironbark woodlands and River Red Gum forest west of the Great Dividing Range and Blackbutt-Bloodwood forest with heath understorey in coastal areas. Prefers mixed species stands with a shrub or <i>Acacia</i> midstorey. Live in family groups of a single adult male one or more adult females and offspring. Require abundant tree hollows for refuge and nest sites. Diet varies seasonally and consists of Acacia gum, eucalypt sap, nectar, honeydew and manna, with invertebrates and pollen providing protein.	Unlikely. No suitable habitat for the species occurs within the Site. Only one record of the species exist within the 10km locality. It is located at approximately 3.5 km to the south from the Site and was recorded in 1997.	Negligible	NA	Low
Petauroides volans Greater Glider Source: PMST		V	The greater glider is restricted to eastern Australia, occurring from the Windsor Tableland in north Queensland through to central Victoria (Wombat State Forest), with an elevational range from sea level to 1200 m above sea level. The greater glider is an arboreal nocturnal marsupial, largely restricted to eucalypt forests and woodlands. During the day it shelters in tree hollows, with a particular selection for large hollows in large, old trees. The greater glider is considered to be particularly sensitive to forest clearance.	Unlikely Limited substandard foraging habitat for the species occurs in remnant vegetation within the Development Site. No rsheltering habitat is present and the species has not been recorded within the 10km locality.	Negligible	NA	Low
Petrogale penicillata Brush-tailed Rock-wallaby Source: BioNet PMST	V		In NSW they occur from the Queensland border in the north to the Shoalhaven in the south, with the population in the Warrumbungle Ranges being the western limit. Occupy rocky escarpments, outcrops and cliffs with a preference for complex structures with fissures, caves and ledges, often facing north. Browse on vegetation in and adjacent to rocky areas eating grasses and forbs as well as the foliage and fruits of shrubs and trees. Shelter or bask during the day in rock crevices, caves and overhangs and are most active at night.	Unlikely. Limited habitat for the species occurs within the Site. Only one record of the species exist within the 10km locality. It is located at approximately 2.2 km to the south-west from the Site and was recorded in 1997.	Negligible	NA	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Phascolarctos cinereus Koala Source: BioNet, PMST	V	V	The Koala is an arboreal marsupial that inhabits eucalypt woodlands and forests. The species feed on the foliage of more than 70 species of eucalypt and 30 non-eucalypt species.	Potential Suitable feeding trees (e.g. E. albens) are present within the Subject Land No known records exists within the site. The nearest record is located at approximately 3 km south-east and is dated 1995.	Negligible. Any clearing of koala feed tree species has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Pseudomys novaehollandiae New Holland Mouse, Pookila Source: PMST		V	The New Holland Mouse has a fragmented distribution across Tasmania, Victoria, New South Wales and Queensland. Known to inhabit open heathlands, woodlands and forests with a heathland understorey and vegetated sand dunes It is a social animal, living predominantly in burrows shared with other individuals Distribution is patchy in time and space, with peaks in abundance during early to mid stages of vegetation succession typically induced by fire.	Unlikely. Limited substandard woodland habitat for the species is present within the Development Site. No records of the species are known within the 10km locality.	Negligible	NA	Low
Pteropus poliocephalus Grey-headed Flying-fox Source: PMST	V	V	Grey-headed Flying-foxes are generally found within 200 km of the eastern coast of Australia, from Rockhampton in Queensland to Adelaide in South Australia. Occur in subtropical and temperate rainforests, tall sclerophyll forests and woodlands, heaths and swamps as well as urban gardens and cultivated fruit crops. Roosting camps are generally located within 20 km of a regular food source and are commonly found in gullies, close to water, in vegetation with a dense canopy. Feed on the nectar and pollen of native trees, in particular Eucalyptus, Melaleuca and Banksia, and fruits of rainforest trees and vines. Also forage in cultivated gardens and fruit crops.	Potential Potential foraging habitat is present in canopy of Eucalypt trees. Grasslands and cleared land excluded as potential foraging habitat No records of the species exist within the 10km locality. The species was not recorded during surveys.	Negligible. Any clearing of Eucalypt trees has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Microchiropteran Bats							
Chalinolobus dwyeri Large-eared Pied Bat, Large Pied Bat Source: PMST	V	V	The Large-eared Pied Bat is found mainly in areas with extensive cliffs and caves, from Rockhampton in Queensland south to Bungonia in the NSW Southern Highlands. It is generally rare with a very patchy distribution in NSW. There are scattered records from the New England Tablelands and North West Slopes. The species is found in well-timbered areas containing gullies. The species roosts in caves (near their entrances), crevices in cliffs, old mine workings and in the disused, bottle-shaped mud nests of the Fairy Martin (<i>Petrochelidon ariel</i>), frequenting low to mid-elevation dry open forest and woodland close to these features. Females have been recorded raising young in maternity roosts (c. 20-40 females) in roof domes in sandstone caves and overhangs. They remain loyal to the same cave over many years.	Potential. Potential foraging/hunting habitat for cave dependant microchiropteran bats is present in the remnant trees and vegetation. No suitable roosting habitat is available and the microchiropteran bats	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential foraging habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Breeding habitat for these species is not present within the Development Site and avoidance measures for breeding habitat is not required.	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risi Rating
Falsistrellus tasmaniensis Eastern False Pipistrelle Source: BioNet	V		The Eastern False Pipistrelle is found on the south-east coast and ranges of Australia, from southern Queensland to Victoria and Tasmania (OEH 2019). The species prefers moist habitats, with trees taller than 20 m. It generally roosts in eucalypt hollows, but has also been found under loose bark on trees or in buildings. It hunts beetles, moths, weevils and other flying insects above or just below the tree canopy. Hibernates in winter. Females are pregnant in late spring to early summer (OEH 2019). Only one record of the species exist within the 10km locality. It is located at approximately 3.2 km to the south-west from the Site and was recorded in 1997.	Potential foraging/hunting and roosting habitat for hollow dependant microchiropteran bats is present in the remnant trees and vegetation.	Minor Any clearing of native vegetation has the potential to add to the incremental decline of potential foraging habitat available within the region. Removal of important habitat features e.g. 37 hollow bearing trees are to be removed	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Trees will be planted at a rate of 2:1 with two new trees planted for each HBT removed. This results in planting of 74 new trees within the riparian corridor. As naturally formed tree hollows will take many decades to develop, nest-boxes suitable for hollow dependent microbats will be installed prior to HBT clearance and will be monitored during the lifespan of the solar farm	Medium
Miniopterus schreibersii (orinae) oceanensis Eastern Bentwing-bat (Large Bentwinged-bat) Source: BioNet	V		Eastern Bentwing-bats occur along the east and northwest coasts of Australia. Caves are the primary roosting habitat, but also use derelict mines, storm-water tunnels, buildings and other man-made structures. Hunt in forested areas, catching moths and other flying insects above the tree tops. Form discrete populations centred on a maternity cave that is used annually in spring and summer for the birth and rearing of young. At other times of the year, populations disperse within about 300 km range of maternity caves. The nearest known record is located at approximately 150m north-west from the site and was recorded in 2006.	Known. Potential foraging/hunting habitat for cave dependant microchiropteran bats is present in the remnant trees and vegetation. No suitable roosting habitat is available and the microchiropteran bats assessed are unlikely to be dependent on any of the resources available.	Minor Any clearing of native vegetation has the potential to add to the incremental decline of potential foraging habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Breeding habitat for these species is not present within the Development Site and avoidance measures for breeding habitat is not required.	Low
Nyctophilus corbeni Corben's Long-eared Bat, South- eastern Long-eared Bat Source: PMST	V	V	Corben's Long-eared Bat inhabits a variety of vegetation types, including mallee, bulloke <i>Allocasuarina leuhmanni</i> and box eucalypt dominated communities, but it is distinctly more common in box/ironbark/cypress-pine vegetation that occurs in a north-south belt along the western slopes and plains of NSW and southern Queensland. It roosts in tree hollows, crevices, and under loose bark. Slow flying agile bat, utilising the understorey to hunt non-flying prey - especially caterpillars and beetles - and will even hunt on the ground. Mating takes place in autumn with one or two young born in late spring to early summer. Corben's Long-eared Bat has not been recorded within the 10 km locality.	Known. Potential foraging/hunting and roosting habitat for hollow dependant microchiropteran bats is present in the remnant trees and vegetation.	Minor Any clearing of native vegetation has the potential to add to the incremental decline of potential foraging habitat available within the region. Removal of important habitat features e.g. 37 hollow bearing trees are to be removed	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Trees will be planted at a rate of 2:1 with two new trees planted for each HBT removed. This results in planting of 74 new trees within the riparian corridor. As naturally formed tree hollows will take many decades to develop, nest-boxes suitable for hollow dependent microbats will be installed prior to HBT clearance and will be monitored during the lifespan of the solar farm	Medium

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Scoteanax rueppellii Greater Broad-nosed Bat Source: BioNet	V		The Greater Broad-nosed Bat is found mainly in the gullies and river systems that drain the Great Dividing Range, from north-eastern Victoria to the Atherton Tableland. It extends to the coast over much of its range. In NSW it is widespread on the New England Tablelands, however does not occur at altitudes above 500 m. Utilises a variety of habitats from woodland through to moist and dry eucalypt forest and rainforest, though it is most commonly found in tall wet forest. Open woodland habitat and dry open forest suits the direct flight of this species as it searches for beetles and other large, slow-flying insects; this species has been known to eat other bat species. Although this species usually roosts in tree hollows, it has also been found in buildings. Only two records of the species exist within the 10km locality. The nearest record is located at approximately 3.2 km to the south from the Site and was recorded in 1997.	Potential foraging/hunting and roosting habitat for hollow dependant microchiropteran bats is present in the remnant trees and vegetation.	Minor Any clearing of native vegetation has the potential to add to the incremental decline of potential foraging habitat available within the region. Removal of important habitat features e.g. 37 hollow bearing trees are to be removed	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Trees will be planted at a rate of 2:1 with two new trees planted for each HBT removed. This results in planting of 74 new trees within the riparian corridor. As naturally formed tree hollows will take many decades to develop, nest-boxes suitable for hollow dependent microbats will be installed prior to HBT clearance and will be monitored during the lifespan of the solar farm	Medium
Mormopterus eleryi Bristle-faced Free-tailed Bat	V		Distributed from the southern half of the Northern Territory to central Queensland and north-western NSW. In NSW, the species has been recently recorded from only three disjunct locations: thirteen individuals from Gundabooka National Park, south of Bourke; one individual from Dhinnia Dthinawan Nature Reserve (formerly Bebo State Forest), north of Warialda two individuals near Bonshaw. Knowledge of the ecology of the Hairy-nosed Freetail Bat is limited, however evidence suggests that the species depends on hollows and tree fissures for roosting sites. All other Australian species from the same family generally roost in tree hollows and fissures. Appears to be extremely rare throughout its range. Nationally, it has been recorded from only 15 locations.	Potential foraging/hunting and roosting habitat for hollow dependant microchiropteran bats is present in the remnant trees and vegetation.	Minor Any clearing of native vegetation has the potential to add to the incremental decline of potential foraging habitat available within the region. Removal of important habitat features e.g. 37 hollow bearing trees are to be removed	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat. Trees will be planted at a rate of 2:1 with two new trees planted for each HBT removed. This results in planting of 74 new trees within the riparian corridor. As naturally formed tree hollows will take many decades to develop, nest-boxes suitable for hollow dependent microbats will be installed prior to HBT clearance and will be monitored during the lifespan of the solar farm	Medium
Fish							
Maccullochella peelii Murray Cod Source: PMST		V	The Murray Cod is a large freshwater fish endemic to the Murray-Darling Basin, from south east Queensland, through NSW, into Vitoria and South Australia. The species can grow to 100 kg in the wild. The species requires permanent streams and is highly dependent on instream woody structures for habitat, is highly territorial and very aggressive.	Unlikely No suitable habitat for the species is present within the Development Site.	Negligible	NA	Low
Flora							
Acacia pubifolia Velvet Wattle Source: PMST	Е	V	Velvet Wattle is a shrub or small tree 3 - 8 m tall with golden yellow flowers and dark-grey bark. It occurs in NSW and Qld. In NSW it is known from two main populations, one north of Emmaville and the other near Warrabah National Park. The species generally grows in dry shrubby woodland on granite and metasediment soils.	Unlikely No suitable habitat for the species occurs within the Development Site. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible	NA	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Astrotricha roddii Rodd's Star Hair Source: BioNet, PMST	E	E	Rodd's Star Hair is an upright, sparsely-branched shrub 1 - 3 m tall. Rodd's Star Hair usually grows in low dry woodland and shrublands on granite and acid volcanic outcrops, often in rock crevices. The species occurs in NSW in the Ashford area north of Inverell, including Kwiambal and Kings Plains National Parks, Severn River Nature Reserve and Severn River State Forest, and has also been recorded at one site in southern Queensland.	Potential Limited potential habitat for the species occurs within the Subject Land. A total of 12 records of the species exist within the 10 km locality. No records exist within the Development Site and the species was not recorded within the development footprint. The nearest record is located at approximately 4.3 km to the south and was recorded in 1999.	Negligible A total of eleven rocky areas were recorded within the Subject Land. Five of the rocky outcrops have been avoided. Only two outcrops occur in the middle of the cleared paddocks and cannot be avoided by the development footprint. The remaining four may also be avoided during the detailed design and survey of the required riparian buffers and perimeter road.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation and habitat features to be retained within the adjacent habitat.	Low
Boronia granitica Granite Boronia Source: PMST	V	E	Granite Boronia is a medium-sized shrub 0.6 - 2 m tall. It flowers from July to October. It grows on granitic soils amongst rock outcrops, often in rock crevices, and in forests and woodlands on granite scree and shallow soils. Important site characteristics include low precipitation and high levels of solar radiation. This semi-arid soil environment will have selected the more xerophytic species from the available regional assemblage of rainforest species. The largely barren substrate (e.g. granite) may help to control too frequent fire, thus allowing maturity and seed set. Granite Boronia occurs in scattered localities on the New England Tablelands and North West Slopes north from the Armidale area to the Stanthorpe district in southern Queensland. It can be locally common in appropriate habitat (e.g. Torrington).	Potential Suitable substrate for the species is present within the Subject Land. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible A total of eleven rocky areas were recorded within the Subject Land. Five of the rocky outcrops have been avoided. Only two outcrops occur in the middle of the cleared paddocks and cannot be avoided by the development footprint. The remaining four may also be avoided during the detailed design and survey of the required riparian buffers and perimeter road.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation and habitat features to be retained within the adjacent habitat.	Low
Cadellia pentastylis Ooline Source: PMST	V	V	Ooline is a medium-sized spreading tree usually about 10 m tall, and rarely to 25 m. Appears to flower spasmodically, during a general flowering period of October to January. Dispersal of fruit and seed is probably by "passive fall" or by birds. Seeds showed a high rate of infertility at all sites, although they have been successfully germinated and established after heat application. Forms a closed or open canopy mixing with eucalypt and cypress pine species. There appears to be a strong correlation between the presence of Ooline and low- to medium-nutrient soils of sandy clay or clayey consistencies, with a typical soil profile having a sandy loam surface layer, grading from a light clay to a medium clay with depth. Has the capacity to resprout from rootstock and coppice vigorously from stumps, a feature which may be critical for the species survival in a fire-prone environment. Populations display a variety of age classes including large mature trees, suckering regrowth and seedlings. The total area occupied by Ooline is only about 1200 hectares, with remaining populations in NSW still threatened to various degrees by clearing for agriculture and grazing pressures.	Potential The development Site is located within known distribution of the species although it was not recorded on site during seasonal surveys. surveys. Noo known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Callistemon pungens Source: PMST		V	Callistemon pungens is a distinct shrub or small tree ranging from 2-5 m tall. The species inhabits a range from riparian areas dominated by Casuarina cunninghamiana subsp. cunninghamiana to woodland and rocky shrubland. Often in rocky watercourses, usually with sandy granite (occasionally basalt) creek beds. Flowers over spring and summer, mostly in November.	Unlikely No suitable habitat is present within the Development Site. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible	NA	Low
Dichanthium setosum Bluegrass Source: PMST	V	V	Bluegrass is an upright grass less than 1 m tall. Flowering time is mostly in summer. Associated with heavy basaltic black soils and red-brown loams with clay subsoil. Often found in moderately disturbed areas such as cleared woodland, grassy roadside remnants and highly disturbed pasture. (Often collected from disturbed open grassy woodlands on the northern tablelands, where the habitat has been variously grazed, nutrient-enriched and water-enriched). It is open to question whether the species tolerates or is promoted by a certain amount of disturbance, or whether this is indicative of the threatening processes behind its depleted habitat. Associated species include Eucalyptus albens, Eucalyptus melanophloia, Eucalyptus melliodora, Eucalyptus viminalis, Myoporum debile, Aristida ramosa, Themeda triandra, Poa sieberiana, Bothriochloa ambigua, Medicago minima, Leptorhynchos squamatus, Lomandra aff. longifolia, Ajuga australis, Calotis hispidula and Austrodanthonia, Dichopogon, Brachyscome, Vittadinia, Wahlenbergia and Psoralea species. Bluegrass occurs on the New England Tablelands, North West Slopes and Plains and the Central Western Slopes of NSW, extending to northern Queensland. It occurs widely on private property, including in the Inverell, Guyra, Armidale and Glen Innes areas.	Potential. Suitable habitat elements for the species occur within the Development Site. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Eucalyptus caleyi subsp. ovendenii Ovenden's Ironbark Source: BioNet, PMST	V	V	Ovenden's Ironbark grows in grassy woodland on dry, shallow soils of moderate fertility. Its preferred altitudes are 610 to 820 m, on granitic substrates. Ovenden's Ironbark occupies a higher geographical range than that of subspecies caleyi, occurring on the crests of broad high ridges and replacing subspecies caleyi inhabiting the lower slopes in the same general area. Associated species include Eucalyptus melliodora, Eucalyptus dealbata, Eucalyptus albens, Eucalyptus melanophloia and Geijera parviflora. Flowering occurs from July to September, with fruits having a distinctly square cross-section. Can be locally abundant within its grassy woodland habitat. Juveniles were present in about half the sampled sites within Torrington State Conservation Area, indicating good recruitment	Potential The species was not recorded within the Development Site during surveys in September and December 2018. Several species with which Ovenden's Ironbark is associated were recorded within the Development Site, including Eucalyptus dealbata, Eucalyptus albens and Geijera parviflora. Only one record of the species exist within the 10 km locality and at approximately 5.3 km to the south-west from the Development Site. The species was recorded in 1999.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Eucalyptus mckieana MicKie's Stringybark Source: PMST	V	V	MicKie's Stingybark is a medium sized tree about 25 m tall. The species flowers are white, with a flowering period of March to May. The species is remarkable for its very narrow and numerous sucker leaves, the narrowest of all the stringybarks and which persist to a height of 2 to 4 metres. Eucalyptus mckieana is found in grassy open forest or woodland on poor sandy loams, most commonly on gently sloping or flat sites. Associated species at Northern Tablelands sites include Angophora floribunda, Eucalyptus amplifolia, Eucalyptus andrewsii, Eucalyptus bridgesiana, Eucalyptus youmanii, Eucalyptus nicholii, Eucalyptus blakelyi and Eucalyptus conica, and at North Western Slopes sites Eucalyptus andrewsii, Eucalyptus stannicola, Eucalyptus prava and Angophora floribunda. Confined to the drier western side of the New England Tablelands of NSW, from Torrington to Bendemeer. Most populations occur on private property, but it does occur in Kings Plain National Park, Torrington State	Unlikely. Very limited potential habitat is present. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible	NA NA	Low
Eucalyptus nicholii Narrow-leaved Black Peppermint Source: BioNet, PMST	V	V	The Narrow-leaved Black Peppermint typically grows in dry grassy woodland, on shallow soils of slopes and ridges. Found primarily on infertile soils derived from granite or metasedimentary rock. Seedling recruitment is common, even in disturbed soils, if protected from grazing and fire. It tends to grow on lower slopes in the landscape. This species is sparsely distributed but widespread on the New England Tablelands from Nundle to north of Tenterfield, being most common in central portions of its range. Found largely on private property and roadsides, and occasionally in conservation reserves. Planted as urban trees, windbreaks and corridors.	Potential The species was not recorded during surveys in September and December 2018. Only one record of the species exist within the 10 km locality and at approximately 5.5 km to the south-west from the Development Site. The species was recorded in 2000.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Homopholis belsonii Belson's Panic Source: PMST		V	Belson's Panic is a perennial grass growing to 0.5m high. It is known to occur in dry woodland habitats on poor soils, such as those derived from basalt. It occurs on rocky hills supporting White Box (<i>Eucalyptus albens</i>) and in Wilga (<i>Geijera parviflora</i>) woodland; flat to gently undulating alluvial areas supporting Belah (<i>Casuarina cristata</i>) forest; and soils and plant communities of Poplar Box (<i>Eucalyptus populnea</i>) woodlands. It may also be associated with shadier areas of Brigalow (<i>Acacia harpophylla</i>), Myall (<i>A. melvillei</i>), and Weeping Myall (<i>A. pendula</i>) communities; in Mountain Coolibah (<i>Eucalyptus orgadophila</i>) communities; and on roadsides. It is generally found among fallen timber at the base of trees or shrubs, among branches and leaves of trees hanging to ground level or along the bottom of netting fences. It is known to occur within the southern Brigalow belt, Queensland and on the north-western slopes and plains of NSW. In NSW, this species occurs between Wee Waa, Goondiwindi and Glen Innes.	Potential Limited suitable habitat for the species occurs within the Development Site. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Lepidium peregrinum Wandering Pepper-cress Source: PMST		E	The Wandering Pepper-cress is a spreading soft-stemmed perennial herb to sub-shrub 10 - 80 cm tall but sometimes ascending to 2 m in surrounding vegetation. Flowers from January to April. The largest population of Wandering Pepper Cress occurs in an open riparian forest on the banks of the Tenterfield creek at Clifton. Sandy alluvium is the main soil type at the site. Associated species at the Clifton site are dominated by Eucalyptus camaldulensis and Casuarina cunninghamiana, with a variably dense shrubby understorey of Hymenanthera dentata, Bursaria spinosa, Acacia fimbriata, Acacia floribunda, Callistemon viminalis and Leptospermum brachyandrum. Lepidium peregrinum was most abundant in the tussock grassland fringe of the riparian open forest, comprising Poa species, Lomandra longifolia and Paspalum dilatatum. After 2001, the species was found near Tenterfield and south-eastern Queensland.	Unlikely No suitable habitat for the species is present. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible	NA	Low
Leucopogon confertus Torrington Beard-heath Source: PMST		E	Leucopogon confertus is known from only one collection along Silent Grove Road, near Torrington, in the NSW Northern Tablelands. It is possible that the species is extinct. Little is known of the habitat of <i>L. confertus</i> , but it may occur in open forest or woodland in rocky granite areas. This species occurs within the Border Rivers—Gwydir (NSW) Natural Resource Management Region. The distribution of this species overlaps with the "White Box-Yellow Box-Blakely's Red Gum Grassy Woodland and Derived Native Grassland".	Potential Limited potential habitat is present within the Subject Land. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible	NA	Low
Polygala linariifolia Native Milkwort	E	-	Native Milkwort is an annual or perennial herb recorded to occur north from Copeton Dam and the Warialda area to southern Queensland; also found on the NSW north coast near Casino and Kyogle, and there is an isolated population in far western NSW near Weebah Gate, west of Hungerford. It occurs in sandy soils in dry eucalypt forest and woodland with a sparse understorey. The species has been recorded from the Inverell and Torrington districts growing in dark sandy loam on granite in shrubby forest of Eucalyptus caleyi, Eucalyptus dealbata and Callitris, and in yellow podsolic soil on granite in layered open forest. Flowers from spring to summer (OEH 2019).	Potential Limited potential habitat is present within the Subject Land. No known records of the species exist within the 10 km locality. The species was not recorded during surveys.	Negligible	NA NA	Low
Prasophyllum sp. Wybong (C.Phelps ORG 5269) A leek Orchid Source: PMST		CE	Prasophyllum sp. Wybong (C. Phelps ORG 5269) is a terrestrial orchid that grows to approximately 30 cm high. A perennial orchid, appearing as a single leaf over winter and spring. Flowers in spring and dies back to a dormant tuber over summer and autumn. Known to occur in open eucalypt woodland and grassland. Endemic to NSW, it is known from near Ilford, Premer, Muswellbrook, Wybong, Yeoval, Inverell, Tenterfield, Currabubula and the Pilliga area. Most populations are small, although the Wybong population contains by far the largest number of individuals.	Potential Suitable habitat for the species is present within the Development Site. No known records of the species exist within the 10 km locality. The species was not recorded during surveys in September and December 2018.	Negligible	NA	Low

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Rutidosis heterogama Heath Wrinklewort Source: BioNet, PMST	V	V	The Heath Wrinklewort is a small perennial herb to 30 cm tall. It grows in heath on sandy soils and moist areas in open forest, and has been recorded along disturbed roadsides. It flowers mainly in Autumn. The species has been recorded from near Cessnock to Kurri Kurri with an outlying occurence at Howes Valley. On the Central Coast it is located north from Wyong to Newcastle. There are north coast populations between Wooli and Evans Head in Yuraygir and Bundjalung National Parks. It also occurs on the New England Tablelands from Torrington and Ashford south to Wandsworth south-west of Glen Innes.	Potential Limited potential habitat for the species might occur within the Subject Land. Surveys were undertaken outside the flowering period for the species. Only one record of the species exist within the 10 km locality and at approximately 5.4 km to the south from the Development Site. The species was recorded in 1999.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Swainsona sericea Silky Swainson-pea Source: BioNet	V		The Silky Swainson-pea is a prostrate or erect perennial, growing to 10 cm tall. The species flowers in spring and produces hairy pods. It is found in Natural Temperate Grassland and Snow Gum <i>Eucalyptus pauciflora</i> Woodland on the Monaro. It is also found in Box-Gum Woodland in the Southern Tablelands and South West Slopes. It is sometimes found in association with cypress-pines <i>Callitris</i> spp. Habitat on plains unknown. Regenerates from seed after fire.	Potential. Cypress pines, Whtie Cypress Pine (Callitris glaucophylla), was recorded within the Development Site. The Development Site is highly disturbed due to historical land use and current land use for grazing. Therefore, it is considered the species has a very low potential to occur within the Development Site. The species was not recorded during surveys in Spring and Summer. Only one record of the species exist within the 10 km locality and at approximately 3.5 km to the south-west from the Development Site. The species was recorded in 2003.	Negligible. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. Future development is unlikely to exacerbate the existing degree of edge effects or degradation of any retained habitats.	Limit clearing through delineation of designated construction areas. This will help to protect native vegetation to be retained within the Development Site and adjacent habitat.	Low
Thesium australe Austral Toadflax, Toadflax Source: PMST	V	V	The Austral Toadflax is a small, straggling herb to 40 cm tall. It is found in very small populations scattered across eastern NSW, along the coast, and from the Northern to Southern Tablelands. It is also found in Tasmania and Queensland and in eastern Asia. Although originally described from material collected in the SW Sydney area, populations have not been seen in a long time. It may persist in some areas in the broader region. It occurs in grassland on coastal headlands or grassland and grassy woodland away from the coast. It is often found in association with Kangaroo Grass (<i>Themeda australis</i>). The species is a root parasite that takes water and some nutrient from other plants, especially Kangaroo Grass.	Potential Limited suitable habitat for the species is present within the Development Site. No records of the species exist within the 10 km locality and the species was not recorded during the September and December 2018 surveys.	Negligible	NA	Low

GAIA BONSHAW SOLAR EIS

Biodiversity Development Assessment Report

Scientific Name Common Name	BC Act	EPBC Act	Species and Habitat Information	Likelihood of species occurring in the Site	Potential Impacts	Mitigation Measures	Residual Risk Rating
Tylophora linearis	V	E	Tylophora linearis is a slender, almost hairless twiner	Potential	Negligible	NA	Low
			with a clear sap. It grows in dry scrub and open forest.	Limited suitable habitat for the			
			Recorded from low-altitude sedimentary flats in dry	species is present within the			
Source: PMST			woodlands of Eucalyptus fibrosa, Eucalyptus	Development Site.			
			sideroxylon, Eucalyptus albens, Callitris endlicheri,	No records of the species exist			
			Callitris glaucophylla and Allocasuarina luehmannii.	within the 10 km locality and			
			Also grows in association with Acacia hakeoides, Acacia	the species was not recorded			
			lineata, Melaleuca uncinata, Myoporum species and	during the September and			
			Casuarina species. It flowers in spring, with flowers	December 2018 surveys.			
			recorded in November or May with fruiting probably 2 to				
			3 months later.				

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development	Assessment Report
APPENDIX E	TEST OF SIGNIFICANCE AND SIGNIFICANT IMPACT ASSESSMENT

Test of Significance

The following Test of Significance (ToS) have been prepared in accordance with Section 7.3 of the BC Act and OEH (2018) *Threatened Species Test of Significance Guidelines*.

Grey-crowned Babbler (Pomatostomus temporalis) – Vulnerable

Species Overview

The Grey-crowned Babbler is a bird listed as vulnerable under the BC Act.

The Grey-crowned Babbler inhabits open Box-Gum Woodlands on the slopes, and Box-Cypress-pine and open Box Woodlands on alluvial plains. Woodlands on fertile soils in coastal regions. Its flight is laborious so birds prefer to hop to the top of a tree and glide down to the next one. Birds are generally unable to cross large open areas. It lives in family groups that consist of a breeding pair and young from previous breeding seasons. A group may consist of up to fifteen birds. All members of the family group remain close to each other when foraging. It is insectivorous and it forages on the trunks and branches of eucalypts and other woodland trees or on the ground, digging and probing amongst litter and tussock grasses. It builds nests that are used as dormitory and roosting and uses them all year round. It breeds between July and February. Territory ranges from one to 50 hectares (usually ten hectares) and are defended all year.

Grey-crowned Babblers are obligate communal breeders that form a family group, in which offspring from the previous season and other unrelated birds help to raise the current's year's brood. Young birds stay with the family group for at least one year after fledging and may remain for two or more years acting as non-breeding helpers. As breeding spaces become available in the population, some helpers may disperse to establish their own breeding group. In some populations, breeding success is related to the number of helpers. Population viability studies in Victoria suggests that a viable population is likely to contain more than ten family groups, while populations with less than ten family groups are likely to have high rate of extinction. In NSW, the species breeds between July and February (OEH 2019).

It has been suggested that cooperative breeder species, such as the Grey-crowned Babbler, are more sensitive to habitat fragmentation and loss (including loss due to fire) as availability of resources for breeding decreases (Fischer 2011). Habitat fragmentation and predation by introduced species being the major threats to the species. The species is capable to survive in disturbed landscapes, such as urban areas, where proportion of ground cover and leaf litter provides sufficient food (Lambert and Ford 2016, Stevens et. al. 2015).

Results of the NSW BioNet atlas indicated that a total of three records of the Grey-crowned Babbler exists within the 10 km locality. None of those known (i.e. BioNet) records are within the Development Site. The nearest BioNet record of the species dates back to 2010 and was located approximately 500 m east of the Development Site.

The Grey-crowned Babbler was observed during surveys in September and December 2018, heard during the March 2019 survey and observed more recently during surveys in November 2019 and January 2020. In December 2018, the Grey-crowned Babbler was recorded at 20 locations on the Subject Land and a total of 21 bird nests likely to belong to the species were observed (see Figure 5.2). The records included six individuals observed, 21 calls heard with an additional two potential calls heard at distance. A pair was also observed adding twigs to a nest located on the lower branches of a Silver-leaved Ironbark (*Eucalyptus melanophloia*). At least 12 individuals were observed on and adjacent to the Subject Land during more recent surveys in November 2019 and January 2020.

The Grey-crowned Babbler was observed in the following vegetation types:

- 594_Moderate (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion). This vegetation zone covers a total of 4.10 ha on the Subject Land;
- 596_Moderate (Tumbledown Red Gum White Cypress Pine Silver-leaved Ironbark shrubby woodland mainly
 in the northern Nandewar Bioregion). This vegetation zone covers a total of 11.15 ha are on the Subject Land;
- 544_Low (Rough-barked Apple White Cypress Pine Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion). This vegetation zone covers a total of 0.83 ha on the Subject Land
- 516_Very Low (Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion). This vegetation zone covers a total of 3.00 ha on the Subject Land; and
- 516_Disturbed Grasslands (Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion). This vegetation zone covers a total of 7.39 ha on the Subject Land.

Test of Significance

(a) in the case of a threatened species, whether the proposed development or activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction.

The Grey-crowned Babbler individuals observed within the Development Site are likely to be part of a single family group utilising the grassy woodlands and derived grasslands. Based on published information, the home range of a local population can range from one to 50ha (usually 10 ha). It is likely the family group range extends beyond the development footprint, as two potential Grey-crowned Babbler nests were recorded to the north-east from the Development Site. The movement of this species is expected to be largely confined to the woodland habitats as this species has limited flying capacity. The riparian corridors (with a 20-30m buffer) have been avoided although nests will be removed within the development footprint and any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region.

The removal of 8 nests (38 % of the 21 nests recorded on site) will reduce the availability of roosting dormitories for the species, and will likely reduce available nesting sites for current breeding adults and for young individuals sexually mature seeking to establish breeding nests. Removal of trees with nests will be included in the vegetation clearing protocol including any seasonal constraints to avoid impacting any juveniles or unfledged chicks. The vegetation clearance protocol should also consider that nests are salvaged and translocated to a suitable location within the riparian corridor where possible.

A portion of felled trees will be salvaged as foraging habitat for fauna and translocated to suitable areas in the remainder of the Project Boundary.

Tree replacement is also proposed as a mitigation measure for loss of hollow bearing trees. Replacement of trees at a rate of 2:1 (i.e. two trees will be planted to replace each hollow bearing tree removed). This results in planting of 68 new trees within the riparian corridor. In the event that a newly-planted tree dies during the lifespan of the solar farm, the client has the responsibility to replace it in order to achieve a 100% recruitment of 68 trees. This is expected to improve the current fragmented condition of the riparian corridor and assist movement of the Grey-crowned Babbler along creek-lines outside of the Development Site and beyond the solar panel fields. Monitoring of the revegetation works would also include monitoring of the local Babbler population including population numbers and evidence on ongoing breeding (active nests).

Based on the application of these mitigation measures the proposed solar farm is unlikely to result in an adverse effect on the life cycle of the species such that the local population might be placed at risk of extinction.

- (b) in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
 - '(i) Is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or

Not applicable

'(ii) Is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction.

Not applicable

Test of	Significance
(c)	in relation to the habitat of a threatened species or ecological community:
	(i) the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and
	The movement of this species is largely confined to the woodland habitats and riparian corridors as this species has limited flying capacity. Habitat to be removed includes 29.17 ha of woodland zones and 11.80 ha of Derived Native Grasslands. This constitutes 27.5 % of the Development Site.
	The riparian corridors (with a 20-30m buffer) have been avoided although 8 nests (38% of the 21 nests recorded on site) will be removed within the development footprint and any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region. No vegetation removal will occur outside of the Development Site and the remainder of the Property Boundary will not be modified as result of the current proposal.
	(ii) whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed action, and
	The removal of native vegetation has the potential to result in fragmentation of fauna habitat, with resultant effects on fauna species movement, reproduction and gene flow. The impact of this vegetation clearance in an already highly modified landscape is anticipated to be negligible for species such as the Greycrowned Babbler given that no significant fauna movement corridors currently exist within the development site (excluding the riparian corridor).
	The movement of this species is largely confined to the woodland habitats and riparian corridors as this species has limited flying capacity. The riparian corridors (with a 20-30m buffer) have been avoided and the proposed replacement tree planting is expected to improve the current fragmented condition of the riparian corridor and assist movement of the Grey-crowned Babbler along creek-lines outside of the Development Site and beyond the solar panel fields. Monitoring of the plantings would also include monitoring of the local Babbler population including population numbers and evidence on ongoing breeding (active nests).
	It is considered that the development of this site will not further isolate or fragment any known habitat linkages for the local population of Grey-crowned Babbler.
	(iii) the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality
	The Development Site has been selected as being a suitable site for a solar plant based on its mostly cleared landscape with minimal vegetation removal required. Habitat to be removed includes 29.17 ha of woodland zones and 11.80 ha of Derived Native Grasslands (27.5 % of the Development Site). Although localised and short term impacts may be observed, the local population of Babbler would not be dependent on this habitat for their long term survival, particularly given the retention (and enhancement) of the riparian corridors which will assist movement of the Grey-crowned Babbler along creek-lines outside of the Development Site and beyond the solar panel fields.
(d)	whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly)
	The proposed development is not located within areas listed as Critical Habitat Declarations in the Register of Declared Areas of Outstanding Biodiversity Value in NSW.

Test of Significance

(e) whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key of a key threatening process.

The proposed development of this site is likely to include the following key threatening processes (KTP):

- Infestation of habitat by invasive weed exotic perennial grasses.
- Aggressive exclusion from forest and woodland habitat by over abundant Noisy Miners.
- Nest predation by species such as ravens and butcherbirds in the fragmented landscape.

Mitigation measures such as weed management, monitoring and control of invasive and nuisance species will be required during the construction and operational phase of the proposed solar farm development..

Conclusion

Although localised and short term impacts may be observed, the local population of the Grey-crowned Babbler would not be dependent on development site for their long term survival, particularly given the retention (and enhancement) of the riparian corridors which will assist movement of the Grey-crowned Babbler along creek-lines outside of the Development Site and beyond the solar panel fields.

Loss of habitat for the Grey-crowned Babbler within the Development Site will require offsetting as per the BC Act. The possibility to offset the direct impact on habitat of the Grey-crowned Babbler by active management and/or enhancement of retained areas of habitat may be considered a viable option in consultation with BCD.

It is considered that with the implementation of mitigation measures, the proposed solar farm development will not result in any long term significant effect on the population.

References

- DSE (2003) Grey-crowned Babbler. Flora & Fauna Guarantee Action Statement No. 34. Victoria's Department of Sustainability and Environment, Victoria.
- Ellis M.V. and Taylor J.E. (2013) Birds in remnant woodland vegetation in the central wheatbelt of New South Wales during drought declared years 2005 to 2009. Australian Zoologist 36(3): 332-348
- Fischer S.E. (2011) The effect of fire on the breeding ecology of the Grey-crowned Babbler (*Pomatostomus temporalis*) in the Australian Monsoon Tropics. Bachelor of Science (Honours Thesis), Charles Darwin University, Darwin.
- Fragar L. (2009) Husbandry Manual Grey-crowned Babbler Pomastostomus temporalis. Delungra, NSW.
- Harrison C., Lloyd H. and Field C. (2016) Evidence review of the impact of solar farms on birds, bats and general ecology. Manchester Metropolitan University, UK.
- Lambert K.T.A. and Ford H. (2016) Habitat use by grey-crowned babbler, Pomatostomus temporalis, in urban and peri-urban environments. Pacific Conservation Biology 8pp.
- OEH (2019) Grey-crowned Babbler Profile. NSW Office of Environment and Heritage. On-line resource accessed via: https://www.environment.nsw.gov.au/threatenedspeciesapp/profile.aspx?id=10660
- Stevens K.P. (2015) Species conservation in fragmented landscapes: Implications for the Grey-crowned Babbler. PhD Thesis, Deakin University.
- Stevens K.P., Holland G.J., Clarke R.H., Cooke R. and Bennett A.F. (2015) What determines habitat quality for a declining woodland bird in a fragmented environment: The grey-crowned babbler Pomatostomus temporalis in South-Eastern Australia? PLOS ONE 10(6): e0130738.
- Vesk P.A., Robinson D., van der Ree R., Wilson C.M., Saywell S. and McCarthy M.A. (2015) Demographic effects of habitat restoration for the Grey-crowned Babbler Pomatostomus temporalis, in Victoria, Australia. PLOS ONE 10(7): e1030153.

Bristle-faced Free-tailed Bat (Mormopterus eleryi) – Endangered

Species Overview

The Bristle-faced Free-tailed Bat or Hairy-nosed Freetail Bat (*Mormopterus eleryi* Reardon and McKenzie, 2008) (Reardon et. al. 2008) or *Mormopterus (Setirostris) eleryi* (Reardon et. al. 2014) is a microchiropteran bat species listed as endangered under the BC Act. The species was collected and referred to as *Mormopterus* 'Species 6' (e.g. Churchill 1988, Adams *et. al.* 1988) in publications prior to its formal description. The species is identified as data deficient in the action plan for Australian Bats (Duncan *et. al.* 1999).

The Bristle-faced Free-tailed Bat is distributed from the southern half of the Northern Territory to central Queensland and north-western NSW. The species appears to be extremely rare throughout its range. Nationally, it has been recorded from only 15 locations. In NSW, the species has been recently recorded from only three disjunct locations (NSW SC 2004, Pennay *et. al.* 2011, OEH 2019): thirteen individuals were recorded on 21 November 1998 at Yanda Creek in Gundabooka National Park, south of Bourke (Ellis 2001, Pennay 2006 in Scotts 2012); one individual was recorded in 2001 from Dhinnia Dthinawan Nature Reserve (formerly Bebo State Forest) (Pennay 2002); two individuals at and near Maroomba State Conservation Area (near Bonshaw) (Arden 2004 in Scotts 2012) and one individual captured on 25th March 2010 at the McIntyre River in Kwiambal National Park (Scotts 2012), Kwiambal National park and Maroomba State Conservation Area are located at approximately 37km south-west and 27km north-west from the proposed Bonshaw solar farm.

Knowledge of the ecology of the Bristle-faced Free-tailed Bat is limited, however evidence suggests that the species depends on hollows and tree fissures for roosting sites (NSW SC 2004, OEH 2019). All other Australian species from the family Molossidae, which includes the genus *Mormopterus*, generally roost in tree hollows and fissures (NSW SC 2004, OEH 2019) and Ellis (2001) recorded the species along with other six species known to roost in three hollows. Ellis and Wilson (1992) indicate that species within genus *Mormopterus* are aerial insectivorous with very high flight speed and limited manoeuvrability. They forage typically well-above the canopy and only coming low in relatively open areas. Three species within the genus *Mormopterus*, *M. nofolkensis*, *M. planiceps* and *Mormopterus* sp 1 were classified as an assemblage of vegetation dependent bats that generally fly over the canopy of any vegetation type when feeding; and which are dependent on hollows, usually in larger trees, for roosting and breeding (Ellis 2006).

At Yanda Creek (Gundabooka National Park), three individuals of the species were captured in a harp trap set under a *Eucalypt* tree on the margins of a dry creek (Ellis 2001), a similar habitat where the species' calls were recorded in the Bonshaw Solar Farm's assessment area. The habitat where the species was recorded by Ellis (2001) was a River Red Gum (*Eucalyptus camadulensis*) and Poplar Box (*E. populnea*) lined watercourse set in Poplar Box woodlands with areas of Mulga (*Acacia aneura*) shrublands in the vicinity. The species has also being regarded as using tree hollows for roosting with a "significant bias" for foraging and roost at locations along riparian habitats (Pennay 2006 in Scotts 2012). At Kwiamba National Park (Scotts 2012), the species was captured on a forested terrace adjacent to the McIntyre River where vegetation comprised a mixed over storey of Blakely's Red Gum (*Eucalyptus blakelyi*), Silver-leaved Ironbark (*E. melanophloia*) and Roughbarked Apple (*Angophora floribunda*). The mid-storey was generally sparse but with patch thickets of tea tree (*Leptospermum brevipes*) and White Cypress Pine (*Callitris glaucophylla*). A grassy understorey dominated the more open areas.

Lactating females had been recorded in November (Ellis 2001).

The calls of the Bristle-faced Free-tailed Bat were recorded in five of the eight SongMeters located within the assessment area. The records were analysed by Mr. Greg Ford (Balance!, see call report in Appendix I). Mr Ford is one of the most experience bat call analysis experts in Australia. Following consultation with BCD, Mr Greg Ford was requested to re-analyze the bat calls to ensure the records of the Bristle-faced Free-tailed Bat were confident. Regarding the Bristle-faced Free-tailed Bat, he noted that the calls of the species are similar to the non-listed Inland Broad-nosed Bat (*Scotorepens greyii*) but that the species can be differentiated based on their feeding buzzes (Ford 2018). A total of 53 definite calls of the species were recorded from S4 (11 definite calls) S2 (5 definite calls), S5 (25 definite calls), S6 (one definite call) and S7 (12 definite calls) (see

Figure 5.2). The locations with the highest number of calls were S5 (25 calls), S7 (12 calls) and S4 (11 calls). These locations were near creek lines (S5 adjacent to a second order creek line with water, whereas S4 and S7 are located at/near a dry first order creek line).

Based record of the Bristle-faced Free-tailed Bat in the assessment area is considered to be a genuine record of this rare microbat species.

The Bristle-faced Free-tail Bat was recorded in the following vegetation types:

- 594_Moderate (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion). A total of 18.78 ha of this vegetation zone is present within the Development Site.
- 594_Low (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion). A total of 7.02 ha of this vegetation zone is present within the Development Site.
- 594_Disturbed Grassland is cleared land allocated as PCT 594 (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion) for assessment purposes. A total of 45.95 ha of this vegetation zone is present within the Development Site.

Test of Significance

(a) in the case of a threatened species, whether the proposed development or activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction.

Little is known about the species and it has been assumed that similar to other species within the genus *Mormopterus*, it relies on tree hollows for roosting and maternity sites. It appears that the species is rare in NSW and no information is currently available regarding its habitat range, reproduction period and social activities although many of the existing records are along riparian habitats. This is also consistent with the results of the surveys within the Project Boundary, within a higher number of calls recorded near creek lines and water bodies. Within the Project Boundary, the riparian corridors (with a 20-30m buffer) have been avoided.

A total of 60 hollow bearing trees were recorded across the assessment area, including 34 HBT within the development footprint. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region and the removal of trees will be included in the vegetation clearing protocol including any seasonal constraints and monitoring of the hollows prior to removal to avoid impacting any breeding females or juveniles. If this species is confirmed utisiling any of the hollow bearing trees, the trees will be left undisturbed until further advice is sought from BCD and a suitably recognised bat expert.

Tree replacement is also proposed as a mitigation measure for loss of hollow bearing trees. Replacement of trees at a rate of 2:1 (i.e. two trees will be planted to replace each hollow bearing tree removed). This results in planting of 68 new trees within the riparian corridor. In the event that a newly-planted tree dies during the lifespan of the solar farm, the client has the responsibility to replace it in order to achieve a 100% recruitment of 68 trees. This is expected to improve the current fragmented condition of the riparian corridor. As naturally formed tree hollows will take many decades to develop, nest-boxes suitable for hollow dependent microbats will be installed prior to HBT clearance and will be monitored during the lifespan of the solar farm

Based on the recommended avoidance and mitigation measures outlined above, including the monitoring of the hollows prior to removal and consultation with BCD in the event that the species is confirmed utilising any of the hollows, it is unlikely that activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction.

Speci	es Overview
(b)	in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
	'(i) Is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or
	Not applicable
	'(ii) Is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction.
	Not applicable
(c)	in relation to the habitat of a threatened species or ecological community:
	(i) the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and
	The Development Site is located in a highly disturbed environment that has undergone extensive clearing. Approximately 34 hollow-bearing trees (HBTs) will be removed across the Development Site, including at least three HBTs along first order streams. No trees will be removed along the second and third order creeklines, where echolocation calls of the Bristle-faced Free-tailed Bat is more frequently recorded.
	Also, it is estimated that a larger number of trees, particularly rough bark <i>Eucalyptus</i> and <i>Callitris</i> are available in vegetation to be retained within the Project Boundary and wider locality. Therefore the removal of approximately 34 trees will represent a small portion of trees available as breeding and roosting habitat for the Bristle-faced Free-tailed Bat.
	Loss of approximately 34 HBTs represent net-loss of suitable habitat for the Bristle-faced Free-tailed Bat and other hollow dependent fauna. However, on a landscape scale, loss of approximately 34 HBTs within a mostly cleared area is not considered significant when considering currently available tree resources elsewhere in the Project Boundary and broader locality.
	(ii) whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed action, and
	The removal of native vegetation has the potential to result in fragmentation of fauna habitat, with resultant effects on fauna species movement, reproduction and gene flow. The impact of this vegetation clearance in an already highly modified landscape is anticipated to be negligible given that no significant fauna movement corridors currently exist within the development site (excluding the riparian corridor which will be avoided).
	As part of mitigation measures to compensate for loss of trees and vegetation, replacement of trees will be planted along the second and third order streams within the Project Boundary. This will increase the vegetation density along the creek-lines and enhance its value as a linking corridor with remnant vegetation fragments and paddock trees beyond the Development Site. This will also enhance corridors for mobile fauna such as the Bristle-faced Free-tailed Bat and other microbats recorded within the Project Boundary.
	Therefore, it is considered that the development of this site will not further isolate or fragment the existing landscape.

(iii) the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality

The Development Site has been selected as being a suitable site for a solar plant based on its mostly cleared landscape with minimal vegetation removal required. It appears that the species is rare in NSW and no information is currently available regarding its habitat range, reproduction period and social activities although many of the existing records are along riparian habitats. This is also consistent with the results of the surveys within the Project Boundary, within a higher number of calls recorded near creek lines and water bodies. Within the Project Boundary, the riparian corridors (with a 20-30m buffer) have been avoided.

The importance of hollow bearing for this species within NSW is also not known however for the purposes of this assessment they have been assumed to be used for roosting and breeding. A total of 60 hollow bearing trees were recorded across the assessment area, including 34 HBT within the development footprint. Any clearing of native vegetation has the potential to add to the incremental decline of potential habitat available within the region and the removal of trees will be included in the vegetation clearing protocol including any seasonal constraints and monitoring of the hollows prior to removal to avoid impacting any breeding females or juveniles. If this species is confirmed utisiling any of the hollow bearing trees, the trees will be left undisturbed until further advice is sought from BCD and a suitably recognised bat expert.

Based on the recommended avoidance and mitigation measures outlined within this assessment, including the monitoring of the hollows prior to removal and consultation with BCD in the event that the species is confirmed utilising any of the hollows, the proposal is unlikely to remove any key habitats that will impact on the long-term survival of the local population.

(d) whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly)

The proposed development is not located within areas listed as Critical Habitat Declarations in the Register of Declared Areas of Outstanding Biodiversity Value in NSW.

(e) whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key of a key threatening process.

Future development of this site is likely to include the following KTP:

- Clearing and removal of hollow bearing trees as a consequence of firewood collection and agricultural and forestry practices. The proposal will remove HBT to give way to installation of solar panels.
- Loss of habitat is exacerbated by its apparent low population numbers.
- Lack of understanding about the species' ecology, distribution and habitat preferences.
- Loss and degradation of foraging and roosting habitat, including changes in vegetation structure due to weed invasion.
- Pesticides and herbicides may reduce the availability of insects, or result in the accumulation of toxic residues in individual's fat stores.

The following mitigation measures will be implemented to minimise the potential of the above listed threats having negative effects on the species:

- A weed management plan will be required to be implemented during the construction and operation phases of the solar farm.
- Chemical use, such as pesticides and herbicides, will be controlled (or avoided) to prevent reduction of insect populations, a feeding resource for the species. This will also eliminate the possibility of trophic induced toxicity due to bioaccumulation in the bat species.

Loss of hollow-bearing trees will be mitigated by replacement of nest boxes (short term mitigation measure) and replacement of trees suitable to develop hollows at their mature stage. It is noted that mature trees (i.e. with more than 80cm DBH) were rare at the site. It has been proposed that periods of time in the range of 180 to 238 years (e.g. soft bark Eucalypt spp.) or up to 324 years (e.g. ironbarks) are required for hollows suitable for fauna (e.g. bats) to form naturally on trees (e.g. Parnaby et. al. 2011). The development footprint area contains mature trees, but not old growth (i.e. trees with DBH greater than 100cm (DEC 2004)). Therefore, replanting trees suitable to naturally develop hollows, e.g. Eucalypt spp., has the potential to replace tree hollows in the long-term and provide natural habitat for hollow-dependent species, such as the Bristle-faced Free-tailed Bat. Evidence suggest that lineal planting and remnants are suitable habitat for bats in disturbed landscapes (Lentini et. al. 2012).

Conclusion

The conservation value of locations where the Bristle-faced Free-tailed Bat had been recorded is unknown due to the lack of knowledge on the biology and ecological requirements for the species. It is known that the species is rare and, based on the ecology of other species within the same genus, it has been assumed the Bristle-faced Free-tailed Bat to be tree hollow dependent. The pattern of the species being more frequently recorded along creek-lines than in other habitats in the landscape has emerged and the riparian corridors (with a 20-30m buffer) have been avoided.

Based on the recommended avoidance and mitigation measures outlined above, including the monitoring of the hollows prior to removal and consultation with BCD in the event that the species is confirmed utilising any of the hollows, it is unlikely that activity is likely to have an adverse effect on the life cycle of the species such that the local population of the species is likely to be placed at risk of extinction although it is noted that any clearing of native vegetation, including paddock trees has the potential to add to the incremental decline of potential habitat available within the region and will require offsetting in accordance with the offset scheme under the BC Act. The possibility to offset the impact on site by active management and/or enhancing retained areas of habitat might be considered as a viable option in consultation with BCD.

References

- Andren M.J. (2004) Nandewar Biodiversity surrogates: vertebrate fauna. Report for the resource and conservation assessment council (RACAC). NSW Western Regional Assessments, coordinated by NSW Department of Infrastructure, Planning and Natural Resources, Project No. NAND05. Department of Environment and Conservation, Coffs Harbour.
- Adams M., Reardon T.R., Baverstock P.R. and Watts C.H.S. (1988) Electrophoretic resolution of species boundaries in Australian Microchiroptera. IV. The Molossidae (Chiroptera). Australian Journal of Biological Sciences 41: 315-326
- Churchill S. (2008) Australian Bats. Second Edition. Jacana Books, Crows Nest.
- DEC (2004) Natural Resource Management Advisory Series: Note 10: Old Growth Forests. NSW Department of Environment and Conservation, Coffs Harbour.
- Duncan A., Baker G.B. and Montgomery N. (1999) The Action Plan for Australian Bats. Environment Australia. On-line resource accessed via: http://155.187.2.69/biodiversity/threatened/publications/action/bats/5.html
- Ellis M. (2001) The first record of the Hairy-nosed Freetail Bat in New South Wales. Australian Zoologist 31(4): 608-609.
- Ellis M. (2006) An assemblage classification of the native vertebrate species of the Western Catchments of NSW.

- Ellis M. and Wilson P. (1992) An overview of the vertebrate fauna of the Brigalow Belt North-east of Bourke, N.S.W. Report prepared by the Royal Zoological Society of NSW for the Western Lands Commission, Bourke.
- Ford G. (2018) Getting the buzz on call identification: an exploration of the usefulness of feeding buzzes for species differentiation. Australian Bat Society Newsletter 50(early 2018): 48.
- Law B.S. (2018) Long-term research on forest bats: we have the technology. Australian Zoologist 39(4): 658-668
- Lentini P., Gibbons P., Fischer J., Law B., Hanspach J. and Martin T. (2012) Bats in a farming landscape benefit from linear remnants and unimproved pastures. PLoS ONE 7(11): e48201, DOI: 10.1371/journal.pone.0048201
- Lumsden L.F. (2004) The ecology and conservation of insectivorous bats in rural landscapes. PhD Thesis, Deakin University.
- NSW SC (2004) Hairy-nosed Freetail Bat endangered species listing. NSW Scientific Committee final determination. On-line resource accessed via: https://www.environment.nsw.gov.au/determinations/HairynosedFreetailBatEndSpListing.htm
- OEH (2019) Bristle-faced free-tailed bat, Hairy-nosed Freetail Bat Profile. NSW Office of Environment and Heritage. On-line resource accessed via: https://www.environment.nsw.gov.au/threatenedspeciesapp/profile.aspx?id=10923
- Parnaby H. and Hamilton-Smith E. (2004) The remarkable "Adaptable Bat": a challenge to ecological concepts in the management of Australian forest bats. Pp 81-93. In: Conservation of Australia's Forest Fauna (Second Edition), Lunney D, (editor). Royal Zoological Society of New South Wales, Mosman.
- Parnaby H., Lunney D. and Fleming M. (2011) Four issues influencing the management of hollowusing bats of the Pilliga forest of inland New South Wales. Pp 399-420 *In*: The Biology and Conservation of Australasian Bats, Law B., Eby P., Lunney D. and Lumsden L. (editors). Royal Zoological Society of NSW, Mosman, NSW, Australia.
- Pennay M., Law B. and Lunney D. (2011) Review of the distribution and status of the bat fauna of New South Wales and the Australian Capital Territory. Pp. 226-256. *In*: The Biology and Conservation of Australasian Bats, Law B., Eby P., Lunney D. and Lumsden L. (editors). Royal Zoological Society of NSW, Mosman, NSW, Australia.
- Reardon T. (2001) the unusual penis morphology of the hairy-nosed freetail bat. Aust. Bat Soc. Newsl. 17: 53-54.
- Reardon T., Adams M., McKenzie N., and Jenkins P. (2008) A new species of Australian Freetail Bat *Mormopterus eleryi* Sp. Nov. (Chiroptera: Molossidae) and a taxonomic reappraisal of *M. norfolkensis* (Gray). Zootaxa 1875: 1-31.
- Reardon T.B., McKenzie N.L., Cooper S.J.B., Appleton B., Carthew S. and Adams M. (2014) A molecular and morphological investigation of species boundaries and phylogenetic relationships in Australian free-tailed bats *Mormopterus* (Chiroptera: Molossidae). Australian Journal of Zoology 62: 109-136.
- Scotts D. (2012) A record of the Bristle-faced Free-tailed Bat Mormopterus eleryi from Kwianbal National park. Australian Zoologist 36(1): 49-54
- Young R.A. and Ford G.I. (2000) Bat fauna of a semi-arid environment in central western Queensland, Australia. Wildlife Research 27: 203-215

Eastern Cave Bat (Vespadelus troughtoni) - Vulnerable

Species Overview

The Eastern Cave Bat (Vespadelus troughtoni) is a microchiropteran bat species listed as vulnerable under the BC Act.

The Eastern Cave Bat is found in a broad band on both sides of the Great Dividing Range from Cape York to Kempsey, with records from the New England Tableland and the upper north coast of NSW. The western limit appears to be the Warrumbungle Range, and there is a single record from southern NSW, east of the ACT (OEH 2019).

The profile for the species (OEH 2019) indicates the following with regards to the habitat and ecology of the species:

- Very little is known about the biology of this uncommon species.
- A cave-roosting species that is usually found in dry open forest and woodland, near cliffs or rocky overhangs; has been recorded roosting in disused mine workings, occasionally in colonies of up to 500 individuals.
- Occasionally found along cliff-lines in wet eucalypt forest and rainforest.
- Little is understood of its feeding or breeding requirements or behaviour.

The Eastern Cave Bat was recorded in the following vegetation types:

- 594_Moderate (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion). A total of 4.10 ha of this vegetation zone is present within the Development Site.
- 594_Low (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion). A total of 9.54 ha of this vegetation zone is present within the Development Site.
- 594_Disturbed Grassland is cleared land allocated as PCT 594 (Silver-leaved Ironbark White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion) for assessment purposes. A total of 49.81 ha of this vegetation zone is present within the Development Site.

In accordance with the TBDC, the Eastern Cave Bat is also found in the other three PCTs recorded at the Development Site, i.e. PCT 516, 544 and 596.

(a) in the case of a threatened species, whether the proposed development or activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction. There is not suitable roosting or breeding habitat for the Eastern Cave Bat within the Development Site. The species is known to occur in the four PCTs recorded at the Development Site (i.e. PCT 594, 596, 544 and 516) and is likely that foraging resources for the species are present therein. Clearing within the Development Site will likely reduce potential foraging resources for the species. However, it is considered this is unlikely that activity is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction.

Species	S Overview
(b)	in the case of an endangered ecological community or critically endangered ecological community, whether the proposed development or activity:
	'(i) Is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or
	Not applicable
	'(ii) Is likely to substantially and adversely modify the composition of the ecological community such that its local occurrence is likely to be placed at risk of extinction.
	Not applicable
(c)	in relation to the habitat of a threatened species or ecological community:
	(i) the extent to which habitat is likely to be removed or modified as a result of the proposed development or activity, and
	The Development Site is located in a highly disturbed environment that has undergone extensive clearing. Approximately 29.17 ha of woodland and 11.80 ha of derived grasslands will be cleared within this disturbed environment. None of the vegetation to be cleared or the land where this will occur includes suitable roosting or breeding habitat for the Eastern Cave Bat.
	Therefore, the project will result in loss of a total of 40.97 ha of potential foraging habitat. However, on a landscape scale, loss of 40.97 ha within a mostly cleared area is not considered significant when considering currently available vegetated resources elsewhere in the Project Boundary and broader locality.
	(ii) whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed action, and
	The removal of native vegetation has the potential to result in fragmentation of fauna habitat, with resultant effects on fauna species movement, reproduction and gene flow. The impact of this vegetation clearance in an already highly modified landscape is anticipated to be negligible given that no significant fauna movement corridors currently exist within the development site (excluding the riparian corridor which will be avoided).
	As part of mitigation measures to compensate for loss of trees and vegetation, replacement of trees will be planted along the second and third order streams within the Project Boundary. This will increase the vegetation density along the creek-lines and enhance its value as a linking corridor with remnant vegetation fragments and paddock trees beyond the Development Site. This will also enhance corridors for mobile fauna such as the Eastern Cave Bat and other microbats recorded within the Project Boundary.
	Therefore, it is considered that the development of this site will not further isolate or fragment the existing landscape.

(iii) the importance of the habitat to be removed, modified, fragmented or isolated to the long-term survival of the species or ecological community in the locality

The Development Site has been selected as being a suitable site for a solar plant based on its mostly cleared landscape with minimal vegetation removal required. The Development Site does not contain suitable roosting or breeding habitat for the Eastern Cave Bat, only potential foraging habitat is present therein.

When considered at a larger spatial scale, loss of 40.97 ha of potential foraging ground in a disturbed land is not significant as more foraging habitat and at better condition is present in the broader locality. Therefore, the proposal is unlikely to remove any key habitats that will impact on the long-term survival of the local population.

(d) whether the proposed development or activity is likely to have an adverse effect on any declared area of outstanding biodiversity value (either directly or indirectly)

The proposed development is not located within areas listed as Critical Habitat Declarations in the Register of Declared Areas of Outstanding Biodiversity Value in NSW.

(e) whether the proposed development or activity is or is part of a key threatening process or is likely to increase the impact of a key of a key threatening process.

Future development of this site is likely to trigger the following KTP:

- Pesticides and herbicides may reduce the availability of insects, or result in the accumulation of toxic residues in individual's fat stores.
- Probable predation by cats and foxes

The following mitigation measures will be implemented to minimise the potential of the above listed threats having negative effects on the species:

- Chemical use, such as pesticides and herbicides, will be controlled (or avoided) to prevent reduction of insect populations, a feeding resource for the species. This will also eliminate the possibility of trophic induced toxicity due to bioaccumulation in the bat species.
- Monitoring and management of invasive pest species. This will minimize the possibility of predation on native species, such as microbats.

Conclusion

There is no suitable roosting and breeding habitat for the Eastern Cave Bat within the Development Site. The Development Site contains vegetation with potential foraging resources for the species. Given that little is known about the feeding behaviour of the Eastern Cave Bat, the conservation value of locations where the species had been recorded is unknown. Given that microchiropteran bats are considered to be insectivorous, use of pesticides and herbicides at the site will be avoided to prevent indirect impacts on insect populations and bioaccumulation on predators, such as the Eastern Cave Bat.

Based on the recommended avoidance and mitigation measures outlined above, including the monitoring of the invasive pest species and avoidance in use of herbicides and pesticides, it is unlikely that activity is likely to have an adverse effect on the life cycle of the species such that the local population of the species is likely to be placed at risk of extinction. Although it is noted that any clearing of native vegetation, including paddock trees has the potential to add to the incremental decline of potential habitat available within the region and will require offsetting in accordance with the offset scheme under the BC Act. The possibility to offset the impact on site by active management and/or enhancing retained areas of habitat might be considered as a viable option in consultation with BCD.

References

OEH (2019) Eastern Cave Bat – Profile. NSW Office of Environment and Heritage. On-line resource accessed via:

https://www.environment.nsw.gov.au/threatenedspeciesapp/profile.aspx?id=10829

www.erm.com Version: 2.0 Project No.: 0470861 0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

Significant Impact Assessment

The following Significant Impact Assessment (SIA) have been prepared in accordance with DoE (2013) *Matters of National Environmental Significance. Significant Impact Guidelines 1.1 under the Environment Protection and Biodiversity Conservation Act 1999.*

Cicadabird (Coracina tenuirostris) - EPBC Act status: Marine

Species Overview

The Cicadabird (*Coracina tenuirostris*) is listed as Marine under the EPBC Act (DoEE 2019), whereas the subspecies Melville Cicadabird (*Coracina tenuirostris melvillensis*) is listed as Migratory. Given that Significant Impact Criteria for marine species is not provided in DoE (2013), the SIA for Cicadabird has been based on criteria for migratory species. The Cicadabird is listed as least concern (ALA 2019, BirdLife International 2017).

The Common Cicadabird (*Edolisoma tenuirostre* synonym *Coracina tenuirostris*) is a full migrant bird species found in Australia, Indonesia, New Guinea and the Solomon Islands (BirdLife International 2017, Wikipedia 2019). In Australia, the Cicadabird inhabits the northernmost part of the NT and WA. It also inhabits the eastern coast of Queensland, New South Wales and Victoria (see Screen Shot below). The species is considered native to the Northern Territory (ALA 2019).



Screen shot 1 Distribution of Cicadabird (source BirLife International 2017)

The natural habitat of the Cicadabird are temperate forests and subtropical or tropical moist lowland forests (Wikipedia 2019). The species is found in several habitats, including forest, artificial/terrestrial, savannah and shrubland. Of those habitats, temperate forests and subtropical/tropical moist lowland forests are habitats of major importance (BirdLife International 2017).

The Cicadabird's generation length is 4.6 years (BirdLife International 2017). The Cicadabrid is considered part of the frugivorous and insectivorous group of birds which has been observed breeding in February and December (Lavery 1985).

One individual of the Cicadabird was recorded in PCT 594 – Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion moderate condition adjacent to Disturbed Land.

Significant Impact Assessment - Significant Impact Criteria for Migratory Species

(a) An action is likely to have a significant impact on a migratory species if there is a real chance or possibility that it will substantially modify (including by fragmenting, altering fire regimes, altering nutrient cycles or altering hydrological cycles), destroy or isolate an area of important habitat for a migratory species.

The sole individual of the Cicadabird was recorded in a moderate condition forest within the Development Site. The Cicadabird has a wide distribution along the eastern coast of Australia and it is of least concern. Regarding the habitat where the species was recorded, the Development Site has undergone a long history of disturbance due to clearing, agriculture and grazing practices, including the vegetation where the species was recorded. The disturbed nature of the site along with the fact that only one individual of the Cicadabird was observed suggests that the species would not be a frequent visitor, or be dependant on any of the resources available. Therefore, it is not considered that the proposed development will substantially modify, destroy or isolate an area of important habitat for the Cicadabird.

(b) An action is likely to have a significant impact on a migratory species if there is a real chance or possibility that it will result in an invasive species that is harmful to the migratory species becoming established in an area of important habitat for the migratory species, or

Information regarding invasive species harmful to the Cicadabird are not readily available (DoEE 2019). It has been assumed that predatory species feeding on birds (e.g. European Red Fox, Cats and Dogs) or species with aggressive behaviour (e.g. Noisy Miner) have potential to have negative effects on individuals and/or population of birds, such as the Cicadabird.

It is not considered that the proposal will result in the establishment of invasive species. Therefore, it is not expected that significant negative effects on the Cicadabird due to interactions with invasive species will result from the proposed solar farm.

(c) An action is likely to have a significant impact on a migratory species if there is a real chance or possibility that it will seriously disrupt the lifecycle (breeding, feeding, migration or resting behaviour) of an ecologically significant proportion of the population of a migratory species.

Very little is known about the breeding, feeding, migratory and resting behaviour of the Cicadabird. Given its widespread distribution in eastern Australia, it is considered sufficient feeding and breeding resources for the species occur.

Due to the Development Site being in a disturbed condition, it is not considered that critical feeding or breeding resources for the species are found. Therefore, it is not considered that the proposed solar farm will result in a significant impact on feeding or breeding resources for the species, nor will it result in disrupting its lifecycle or migratory behaviour.

Conclusion

The Cicadabird is widespread in eastern Australia and is a species of least concern. In spite of little knowledge about the species' feeding, breeding, roosting and migratory behaviour being available, the species continues to persist along the eastern coast of Australia, suggesting sufficient resources are present.

The Development Site is a disturbed environment with cleared and disturbed vegetation. Remnant vegetation patches are disturbed (moderate to very low condition) and are not considered to represent critical habitat for the species. Therefore, it is considered that the proposed solar farm will not result in significant impacts on the species and that a referral to the Commonwealth is not deemed necessary as significant impacts to migratory/marine Matters of National Significance will not occur.

References

- ALA (2019) Coracina (Edolisoma) tenuirostris (Jardine, 1831) Cicadabird. Atlas of Living Australia. On-line resource accessed on 15 April 2019 via:

 https://bie.ala.org.au/species/urn:lsid:biodiversity.org.au:afd.taxon:fbda0192-014f-46ff-a0e0-0617fb687f02
- BirdLife International (2017) Edolisoma tenuirostre. The IUCN Red List of Threatened Species 2017. On-line resource accessed on 15 April 2019 via: https://www.iucnredlist.org/species/103702470/118729711
- DoEE (2019) SPRAT Profile: Coracina tenuirostris Cicadabird. Australian Government,
 Department of Environment and Energy. On-line profile accessed via:
 http://www.environment.gov.au/cgi-bin/sprat/public/publicspecies.pl?taxon_id=649
- Lavery H.J. (1985) Breeding seasons of birds in north-eastern Australia, First Supplement, 1967-74. EMU 86 (Short Communications): 111-113
- Pedersen M.P., Irestedt M., Joseph L., Rahbek C. and Jonsson K.A. (2018) Phylogeography of a 'great speciator' (Aves: *Edolisoma tenuirostre*) reveals complex dispersal and diversification dynamics across the Indo-Pacific. Journal of Biogeography 45(4): 826-837.
- Wikipedia (2019) Common Cicadabird. On-line resource accessed via: https://en.wikipedia.org/wiki/Common_cicadabird

Assessment of Significance

The following Assessment of Significance (AoS) have been prepared in accordance with DPI (2008) Threatened Species Assessment Guidelines. The DPI (2008) guideline, provide a series of criteria to assess whether a proposed development is likely to have significant effects on aquatic threatened species listed under the FM Act. An AoS has been prepared for three aquatic threatened species whose indicative distributional maps suggest they were likely to occur in creeks within the Development Site. The species are:

- Southern Purple Spotted Gudgeon (Endangered Species)
- Eel Tailed Catfish (Endangered Population)
- Olive Perchlet (Endangered Population)


Southern Purple Spotted Gudgeon, Eel Tailed Catfish & Olive Perchlet

Species Overview

Southern Purple Spotted Gudgeon (Mogunda adspersa) - Endangered Species

The Southern Purple Spotted Gudgeon occur in inland drainages of the Murray-Darling basin as well as coastal drainages of northern NSW and Queensland. The western population of the Southern Purple Spotted Gudgeon was previously widespread in the Murray, Murrumbidgee and Lachlan River systems and tributaries of the Darling, but has experienced a significant decline in recent times. Southern Purple Spotted Gudgeons are now extremely rare in inland NSW, having been recorded from this area only once since 1983 (DPI 2019).

The Southern Purple Spotted Gudgeon generally grow to 15 cm in length. The species is found in slow moving or still waters of rivers, creeks and billabongs, often amongst weeds, rocks and snags. They feed mainly on insect larvae, but also consume worms, tadpoles, small fish and some plant matter. Female *Mogurnda adspersa* may lay several batches of eggs per season (30-1,300 per batch). The eggs are deposited in clusters on solid objects such as rocks, wood or broad-leafed plants. The male guards and fans the eggs until hatching (3-8 days). The species is part of the Endangered Aquatic Ecological Community in the Natural Drainage System of the Lower Murray River Catchment, the Endangered Aquatic Ecological Community in the Natural Drainage System of the Lowland Catchment of the Darling River and the Endangered Aquatic Ecological Community in the Natural Drainage System of the Lowland Catchment of the Lowland Catchment of the Lachlan River (FSC 2008a).



DPI (2019) Indicative Distribution Map for the Southern Purple Spotted Gudgeon

A close review of DPI's indicative distribution map for the Southern Purple Spotted Gudgeon (see Figure 5.4) indicates that none of the creeks within the Project Boundary are mapped as habitat for the species.

Eel Tailed Catfish (Tandanus tandanus) - Endangered Population

The Murray-Darling Basin (MDB) population of Eel Tailed Catfish is listed as an Endangered population under the FM Act.

Eel Tailed Catfish is an Australian endemic fish species (DPI 2015). It is naturally distributed throughout the Murray-Darling Basin and in the Eastern drainages NSW north of Newcastle. Eel Tailed Catfish numbers in the Murray-Darling Basin have declined due to a range of impacts including invasive species, habitat degradation, cold water pollution and fishing pressures and are now virtually absent from the Murray, Murrumbidgee and Lachlan catchments (DPI 2019).


Eel Tailed Catfish is a medium-sized fish with a large head and a compressed rear portion of the body. It has a relatively long life span, living for at least 8 years (DPI 2015). It can grow up to 900mm in length and 7kg, however fish over 2kg are exceptional (DPI 2015, 2019). Colour ranges from grey to brown dorsally or laterally, usually mottled with dark brown to black blotchings with a whitish underbelly. Larger fish have less mottlings and can be greener in colour fading to white below (DPI 2019). Catfish are predominantly opportunistic carnivores, feeding mainly on small fish, freshwater prawns, yabbies, zooplankton and insects.

Individuals are sexually mature at 3-5 years of age and spawn in spring/summer when water temperatures are 20-24°C (DPI 2015). Eel Tailed Catfish builds a nest in areas of still water to breed, and their reproduction is not temperature reliant. Males construct and defend a nest up to 2 metres in diameter, made from pebbles and gravel.

Eel Tailed Catfish is a non-migratory, benthic (bottom dwelling) species. It is relatively sedentary and adults typically only move within a 5 km range. Individuals are more active at night compared with during the day (DPI 2015, 2019). The species lives in a wide range of habitats including rivers, creeks, lakes, billabongs and lagoons, and although it inhabits flowing streams, prefers sluggish or still waters. It can be found in clear to turbid waters, and over substrates ranging from mud to gravel and rock. It is rare in natural riverine habitats but can be found in farm dams through-out inland NSW and southern Queensland (FSC 2008b). The species is relatively inactive and do not migrate for spawning, unlike other inland species such as Golden Perch or Murray Cod (DPI 2015, 2019).

Duncan et. al. (2017) indicated that Eel Tailed Catfish populations were most likely to be present in reaches of the MDB if there is a high cover of Cobble/Gravel, submerged macrophytes and low daily flow. Catfish are thought to prefer slow-moving or still waters and are found in greater abundance in lakes and backwaters. Breeding is more likely to occur at sites with a high abundance of Bedrock/Boulders, Riffle/Rapids, Emergent Macrophytes and low Daily Flow. They also note that in reaches of tributaries of the MDB where the species is currently extinct, it is unlikely that the species would naturally re-colonize and establish itself due to it being a not highly migratory species. Catfish have a relatively high minimum spawning temperature threshold relative to other native fishes in the MDB. Thermal pollution may partly explain their disappearance from some rivers in the MDB subject to cold-water releases from large impoundments.

The indicative distributional map for the species indicates the Project Boundary is within the portion of the Murray-Darling Basin where the species had been recorded (see DPI's image below). A zoom into the Project Boundary (see Figure 5.4) shows that the creeks within the Project Boundary are not mapped as habitat for the Eel Tailed Catfish. Duncan et. al. (2017) indicated that the species has disappeared from the Dumaresq River.



DPI (2019) Indicative Distribution Map for the Eel Tailed Catfish

Olive Perchlet (Ambassis agassizii) - Endangered Population

The Western Population of the Olive Perchlet in the Murray-Darling Basin (MDB) is listed as an Endangered population under the FM Act.


Olive Perchlets are a small native fish that occur in both eastern (coastal) and western (Murray-Darling) drainages, but these populations may be genetically distinct. The western population of the Olive Perchlet was once widespread throughout the Murray-Darling system of South Australia, Victoria, western New South Wales and southern Queensland. This population has suffered a serious decline and is now found only at a few sites in the Darling River drainage (DPI 2019).

Olive Perchlets have an oval shaped body with a moderately large mouth, very large eyes and a forked tail. They are usually semi-transparent, with dark-edged scales forming a distinct pattern. The fins are generally clear, although there is often a broad, blackish band along the edges of the pelvic and anal fins. Olive Perchlet can grow to about 7080 mm but are more commonly less than 40 mm (DPI 2013, 2019).

Males and females reach sexual maturity at one year of age, and live for 2-4 years. Spawning occurs between October and December when the water temperature reaches 23°C. Females lay 200-700 eggs which attach to aquatic plants and rocks on the streambed. The eggs are scattered among vegetation. Both sexes reach maturity in one year, with some males surviving and breeding for two years and some females breeding in their third year (DPI 2009).

Olive Perchlet inhabit rivers, creeks, ponds and swamps. They are usually found in slow-flowing or still waters. They are usually found in sheltered areas such as overhanging vegetation, aquatic macrophyte beds, logs, dead branches and boulders during the day, and disperse to feed during the night. Olive Perchlet feed on a range of zooplankton and aquatic and terrestrial insects (DPI 2013, 2019).

The indicative distributional map for the species in the Murray-Darling Basin indicates the Project Boundary is within the portion of the Murray-Darling Basin where the species had been recorded (see DPI's image below (DPI 2019)). The current distribution of the species in the MDB has decreased (DPI 2013). A zoom into the Project Boundary (see Figure 5.4) shows that the creeks within the Project Boundary are not mapped as habitat for the Olive Perchlet.



DPI (2019) Indicative Distribution Map for the Olive Perchlet

Assessment of Significance

(a) In the case of threatened species, whether the action proposed is likely to have an adverse effect on the life cycle of the species such that a viable local population of the species is likely to be placed at risk of extinction.

The six first order creeks to be lost as part of the proposed development are currently dry, as such, they do not represent suitable habitat for any of these aquatic fish species. The third, second and first order creeks within the Project Boundary to be retained were also dry at the time of surveys and are isolated from the broader riparian system due to the presence of Bruxner Way.

During period of heavy rain and increase surface flow, the possibility exists for aquatic fauna, such as fish, to occasionally occupy these creeks in association with flooding events whereby they can be carried into the creeks by the flow and remained 'trapped' as waters recede. When such floodinduced occasional use of the creeks might occur, it is unlikely those individuals would establish a local population due to the lack of connectivity with the broader riparian system and ephemerality of the creeks.

Species	Overview
	The proposed development is unlikely to result in any impacts to any local population of these aquatic species nor it is predicted that the project or activities associated with it will put the species at risk of extinction.
(b)	in the case of an endangered population, whether the action proposed is likely to have an adverse effect on the life cycle of the species that constitutes the endangered population such that a viable local population of the species is likely to be placed at risk of extinction.
	None of the creeks within the Subject Land provide habitat for any of the species assessed and the proposed development is unlikely to result in any impacts to any local population of these species or have any effect on the life cycle of any local population.
(c)	 in the case of an endangered ecological community or critically endangered ecological community, whether the action proposed: (i) is likely to have an adverse effect on the extent of the ecological community such that its local occurrence is likely to be placed at risk of extinction, or (ii) is likely to substantially and adversely modify the composition of the ecological community
	such that its local occurrence is likely to be placed at risk of extinction.
	NA
(d)	 in relation to the habitat of a threatened species, population or ecological community: (i) the extent to which habitat is likely to be removed or modified as a result of the action proposed, and (ii) whether an area of habitat is likely to become fragmented or isolated from other areas of habitat as a result of the proposed action, and (iii) the importance of the habitat to be removed, modified, fragmented or isolated to the long-
	term survival of the species, population or ecological community in the locality
	None of the creeks within the Subject Land provide habitat for any of the species assessed and the proposed development is unlikely to result in any impacts or remove any habitat critical to the survival of any local population of these species.
(e)	whether the action proposed is likely to have an adverse effect on critical habitat (either directly or indirectly)
	No critical habitat for the three threatened fish species is present at the Development Site.
(f)	whether the action proposed is consistent with the objectives or actions of a recovery plan or threat abatement plan
	Southern Purple Spotted Gudgeon
	No recovery plan or threat abatement plan is available for the Southern Purple Spotted Gudgeon. The final determination for the species indicates that threatening processes associated with the decline of the species are:
	predation by introduced fish such as gambusia and redfin perch,
	 habitat loss, rapid fluctuations in water levels (due to water regulation) that have deleterious effects on successful reproduction and recruitment.
	A Priorities Action Statement for the species is available in DPI (2019). The proposed development do not contravene any of the provided actions for recovery of the Southern Purple Spotted Gudgeon.

Eel Tailed Catfish:

No recovery plan or threat abatement plan is available for the Eel Tailed Catfish. The final determination for the species and its PrimaFace (DPI 2015) indicates that threatening processes associated with the decline of the species are uncertain, but that probably include:

- Historic commercial fishing;
- Loss of habitat (lakes, billabongs, lagoons) through river regulation;
- Interactions with introduced species, such as carp (Cyprinus carpio) and Redfin Perch (Perca fluviatilis);
- Loss of habitat and spawning sites through siltation;
- Reduced success of spawning and recruitment;
- Loss of habitat due to alterations to flow patterns and flooding regimes;
- Reduced habitat and loss of temperature spawning cues due to cold-water discharge from the base of large dams and high-level weirs;
- Loss of aquatic plants;
- Chemical pollution, including agricultural pesticides.

A Priorities Action Statement for the species is available in DPI (2019). The proposed development do not contravene any of the provided actions for recovery of the Eel Tailed Catfish.

Olive Perchlet:

No recovery plan or threat abatement plan is available for the Olive Perchlet. The final determination for the species and its PrimaFace (DPI 2013) indicates that threatening processes associated with the decline of the species may include:

- Predation by introduced fish such as Mosquitofish (Gambusia holbrooki) and Redfin perch (Perca fluviatilis).
- Habitat degradation and loss, including the removal of vegetation, logs and snags.
- Rapid fluctuations in water levels (due to river regulation) that have deleterious effects on successful reproduction and recruitment.
- Cold water pollution from impoundment water release restricting spawning.
- Loss of instream aquatic vegetation through the impacts of river regulation and carp (Cyprinus carpio).

A Priorities Action Statement for the species is available in DPI (2019). The proposed development do not contravene any of the provided actions for recovery of the Western Population of the Olive Perchlet.

Conclusion

None of the creeks within the Subject Land provide habitat for any of the species assessed and the proposed development is unlikely to result in any impacts or remove any habitat critical to the survival of any local population of these species.

References

- DPI (2013) **PrimeFact: Olive Perchlet (western population)** *Ambassis agassizii*. NSW Department of Primary Industries, Port Stephens.
- DPI (2015) **PrimeFact: Eel-tailed Catfish population in the Murray-Darling Basin,** *Tandanus* **tandanus.** NSW Department of Primary Industries, Port Stephens.
- DPI (2018) Protecting Eel-Tailed Catfish in Western NSW a guide for fishers and land managers. NSW Department of Primary Industries.
- DPI (2019) **Threatened Species and Maps**. On-line resources accessed via:

 https://www.dpi.nsw.gov.au/fishing/species-protection/threatened-species-distributions-in-nsw
- Duncan M., Gilligan D. and Robinson W. (2017) **Freshwater catfish (***Tandanus tandanus***) habitat requirements**. NSW Department of Primary Industries, Narrandera NSW.
- FSC (2008a) Final Determination: *Mogurnda adspersa* Purple Spotted Gudgeon. Listing by the Fisheries Scientific Committee under the NSW Fisheries Management Act 1994 (Ref No. FD35, File No. FSC 00/15, January 2008).
- FSC (2008b) Final Determination: The *Tandanus tandanus* Eel Tailed Catfish in the Murray/Darling Basin as an Endangered Population. Listing by the Fisheries Scientific Committee under the NSW Fisheries Management Act 1994 (Ref No.FD41, File No. FSC08/02).
- FSC (2009) Recommendation Ambassis agassizii (Olive Perchlet). Recommendation by the Fisheries Scientific Committee under the NSW Fisheries Management Act 1994 (Ref. No. RF17, File No. FSC 01/09).
- Rourke M. and Gilligan D. (2010) **Population genetic structure of freshwater catfish (***Tandanus tandanus***) in the Murray-Darling Basin and coastal catchments of New South Wales: Implications for future re-stocking programs**. Industry & Investment NSW, Narrandera Fisheries Centre, Narrandera NSW.

www.erm.com Version: 2.0 Project No.: 0470861 0470861 GAIA Bonshaw Solar BDAR Version 4.docx

Revised Biodiversity Development	Assessment Report
APPENDIX F	SURVEY METHODS AND EFFORT
AFFLINDIX	SORVET METHODS AND EFFORT

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

GAIA BONSHAW SOLAR EIS

Survey Method and Effort

Flora and fauna surveys were undertaken within the Development Site at locations shown in Figure 5.1. Environmental conditions are presented in Table F.1 below. Weather conditions are as per the Bureau of Meteorology's Applethorpe (Station 041175).

Fauna survey methods and effort are presented in Table F.1.

Targeted flora surveys methods and effort are presented in Table F.3.

Table F.1 Weather during Surveys

Date	Min Temp	Max Temp	Rainfall (mm)	Other
10/09/2018	3.1	22.6	0	
11/09/2018	6.0	20.3	0	
12/09/2018	5.1	21.7	0	
13/09/2018	5.1	24.2	0	
14/09/2018	5.2	25.4	0	
10/12/2018	14.0	30.7	0	
11/12/2018	16.0	27.4	0	
12/12/2018	14.2	30.3	0	
13/12/2018	14.2	29.2	0.2	
14/12/2018	17.0	27.7	26	
25/03/2019	19.6	29.9	0	
26/03/2019	19.1	24.1	0	
27/03/2019	14.6	17.8	1.2	
28/03/2019	14.0	21.7	1.2	
21/01/2020	24.4	37	0.2	
22/01/2020	20.5	38	0	

Table F.2 Summary of Fauna Survey Effort

Bird Surveys

Survey method: A 20 minute point survey was undertaken by one or two ecologists. Survey consisted of listening bird calling, observing birds activity using binoculars. Bird surveys are in general agreement with guidelines such as DEWHA (2010) Survey guidelines for Australia's threatened birds. Guidelines for detecting birds listed as threatened under the Environment Protection and Biodiversity Conservation Act 1999.

Date	Survey ID	Start Time	Finish Time	Ecologists	Effort (man hours)	Notes
12/09/2018	В0	10:35	11.00	JW	0.40	Diurnal
10/12/2018	B1	19:22	19:42	TC, ACM	0.67	Dusk
11/12/2018	B2	19:25	17:45	TC, ACM	0.67	Dusk
12/12/2018	В3	18:27	18:27	TC, ACM	0.67	Dusk
13/12/2018	B4	10:48	11:08	TC, ACM	0.67	Diurnal
13/12/2018	B5	18:22	18:42	TC	0.33	Dusk
14/12/2018	В6	6:08	6:28	TC	0.33	Dawn
14/12/2018	B7	6:57	7:17	TC	0.33	Dawn
	1	1	To	otal Survey Effort	4.07	

Ecologists: ACM - Adriana Corona Mothe; JW - Joanne Woodhouse; TC - Thomas Cotter

Frog Surveys

Survey method: Frog surveys were undertaken using playback of the call of Tusked Frog (*Adelotus brevis*) using a mobile phone and a speaker. Two or three cycles of two minutes playing the call recording followed by two minutes listening for any response.

Date	Survey ID	Start Time	Finish Time	Ecologists	No Playback events	Playback Duration
12/12/2018	F1	22:15	22:23	TC, ACM	2	2 min call, 2 min listen
13/12/2018	F2	18:58	19:10	TC, ACM	3	2 min call, 2 min listen
14/12/2018	F3	7:30	7:38	TC, ACM	2	2 min call, 2 min listen
Total effort (no call playback events)					7	Survey undertaken at suitable habitat for the targeted species.

Ecologists: ACM - Adriana Corona Mothe; TC - Thomas Cotter

Reptile Surveys

A total of seven reptile surveys were undertaken within rocky areas.

Reptile surveys were design to target the following:

- General reptile habitats
- Targeting habitat for the Zigzag Velvet Gecko (Amalosia rhombifer) in gum trees (e.g. Eucalyptus blakelyi) and Whyte Cypress Pine (Callitris glaucophyla).
- Targeting habitat for the Border Thick-tailed Gecko (Uvidicolus sphyrurus) in rocky habitats, coarse woody debris and leaf litter.
- Targeting habitat for the Pale Headed Snake (Hoplocephalus bitorquatus) treed areas, riparian areas and coarse woody debris.

Survey Method: Two ecologist undertook nocturnal spotlight surveys. Searches included turning rocks, searching underneath leaf litter piles and under logs. Spotlighting also targeted tree trunks. No recommended survey effort for the targeted species is readily available, therefore, survey effort was determined based on standard practice for reptiles, habitat availability at the Development site, species profiles and advice for similar species (e.g. geckos) as per DEWHA (2011) *Survey guidelines for Australia's threatened reptiles*.

Date	Survey ID	Start Time	Finish Time	Ecologists	Effort (Man hours)	Notes
10/12/2018	R1	20:21	20:41	TC, ACM	0.67	Rocky Area. Leaf litter, log, rock, granite. Eastern facing
10/12/2018	R2	19:57	10:17	TC, ACM	0.67	Rocky Area. Leaf litter (3cm depth), log and rock turn, granite, eastern slope
11/12/2018	R3	20:50	21:10	TC, ACM	0.67	Rocky Area. Leaf litter (2cm depth), east facing
11/12/2018	R4	21:26	21:46	TC, ACM	0.67	Rocky Area. Leaf litter (2cm depth), west facing
11/12/2018	R5	21:58	22:09	TC, ACM	0.33	Rocky Area
11/12/2018	R6	21:00	21:34	TC, ACM	1.13	Rocky Area
12/12/2018	R7	20:34	20:54	TC, ACM	0.67	Rocky Area
12/12/2018	R8	20:02	20:22	TC, ACM	0.67	Rocky Area
			To	otal Survey Effort	5.48	Survey effort for reptiles at the Development Site is considered sufficient given the extent of suitable habitat.

Ecologists: ACM - Adriana Corona Mothe; TC - Thomas Cotter

Camera Traps

A total of eight camera traps were installed across the Project Boundary. Camera traps used included:

- Four pocket camera SG565F-8M.
- Four UV565 cameras.

Survey method: Camera traps were set up targeting arboreal fauna, particularly that using tree hollows, such as Squirrel Glider (*Petaurus norfolcensis*). Camera traps were set up to record fauna during three or four nights. The cameras were timed to record fauna activity one hour before dusk to hour after dawn. Analysis of photographic records were undertaken by ERM's ecologists.

Unit ID	Date Set up	Date Removed	Recording Start Time	Recording Finish Time	Effort (Trap Nights)	Notes
C1	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Camera set up on an ironbark. Directed at a hollow in adjacent <i>Angophora floribunda</i> . Lure used: honey water spray
C2	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Camera set up on a <i>Eucalyptus albens</i> at 2m height. Directed at an adjacent <i>E. albens</i> .
C3	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Camera set up on a <i>Callitris glaucophylla</i> and at 1,5m high. Directed towards a <i>Eucalyptus blakelyi</i> which had some arboreal mammal scratch marks.
C4	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Camera set up on a <i>Hakea</i> sp. at 1.5m high. Directed towards a <i>Eucalyptus melanophloia</i> with hollow.
C5	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Camera set up on a <i>Eucalyptus melanophloia</i> and at 2 m high. Directed towards adjacent <i>Eucalyptus melanophloia</i> 's trunk located at the edge of a drainage feature with water present.
C6	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Camera set up on a <i>Dododanea</i> sp. and a 1.5 m high. Directed towards <i>Eucalyptus melanophloia</i> with hollows. Lure used: honey water spray.
C7	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Camera set up on a <i>Callitris glaucophylla</i> and at 1.75m high. Directed towards adjacent a <i>Corymbia dolichocarpa</i> with hollow. Lure used: honey water spray.
C8	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Camera set up on a stag at 1.5m high. Directed towards <i>Eucalyptus dealbata</i> trunk with hollows and scratch marks. Lure used: honey water spray.
				Total trap nights	28	Given that the Development Site has undergone intensive disturbance due to its historical and current land use, an intense level of survey (e.g. 14 night traps and using approximately 10 cameras per hectare (DSEWPC (2011) Survey guidelines for Australia's threatened mammals)) was not considered necessary. We targeted the most likely habitats were arboreal species might occur.

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page F3

SongMeters

Eight SongMeter model SM2+ (Wildlife Acoustics Inc.) were used. A total of 28 trap nights were recorded using eight instruments during three to four nights.

The microchiropteran bat recordings were submitted for analyses by recognised bat call analysis expert, Mr Greg Ford (Balance! Environmental). The report is provided in Appendix I.

Unit ID	Date Set up	Date Removed	Recording Start Time	Recording Finish Time	Effort (Trap Nights)	Notes
S1	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Unit set up on an <i>E. albens</i> ' stag, adjacent dam with water.
S2	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Unit set up on an E albens at 2 m high and directed to adjacent dam.
S3	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Unit set up on an <i>Angophora floribunda</i> , at 1.5m high and along dry drainage feature.
S4	10/12/2018	14/12/2018	18:00 hrs	6:00 hrs	4	Unit set up on an <i>E. melanophloia</i> , at 1.5 m high and along dry drainage feature.
S5	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Unit set up on an <i>A. floribunda</i> , at 2m high, facing drainage feature with water immediately adjacent. Lure used: honey water spray.
S6	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Unit set up on an <i>E. melanophloia</i> with hollows, at 1.75m high and adjacent to a dry drainage feature.
S7	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Unit set up on a <i>C. glaucophylla</i> , at 2m high, directed towards <i>C. glaucophylla</i> regrowth
S8	11/12/2018	14/12/2018	18:00 hrs	6:00 hrs	3	Unit set up on a stag, at 1.75m high, directed towards cleared land adjacent <i>Eucalyptus dealbata</i> with hollows.
				Total	28	A total of 28 trap nights were undertaken at the Assessment Area, including the Development Site (approx. 149.24 ha). In accordance with the OEH (2016) 'Species credit' threatened bats and their habitats. NSW survey guide for the Biodiversity Assessment Method. A minimum of 16 bat nights are required for a default 50ha site. For Subject Lands larger than 50 ha, survey effort is to be scaled up. Based on the highly disturbed and heavily modified nature of the Subject Land and the identification of 29.17 ha of Woodland Vegetation and 11.80 ha Derived Native Grasslands, this survey effort is considered appropriate. SongMeters were installed along riparian areas and areas most likely used by microchiropteran bats as potential flyways.

Ecologists: JW = Joanne Woodhouse; TC = Thomas Cotter; ACM = Adriana Corona Mothe.

Targeted flora survey method

Candidate flora species requiring survey were identified early on during Stage 1 of the project. Table F.3 shows the candidate species identified along with their flowering periods and habitats. Survey method consisted of the following:

- Random meander transects (RMT) were undertaken across the Development Site to undertake vegetation ground-truthing. Along RMT, where suitable habitat for a candidate flora species was identified, the corresponding species was searched for. This was undertaken during the spring and summer survey field trips.
- As part of vegetation mapping, habitat observations and vegetation community observation were undertaken at over 150 locations across the Development Site. At each of these locations, candidate species were searched for if suitable habitat was present.
- At BAM plot locations, candidate flora species were looked for within the BAM plot area. A total of 44 BAM plots were undertaken.

Table F.3 Candidate flora species identified during survey design

Common Name (Scientific	ВС	EPBC	Habitat Targeted	Stratum	Flowering Period	Survey period		
Name)	Act	Act		Targeted		Spring (September 2018)	Summer (December 2018)	Autumn (March 2019)
Native Milkwort (<i>Polygala linariifolia</i>)	V	-	Grassland, Woodlands with <i>Eucalyptus</i> spp. and <i>Callitris</i> spp.	Groundcover	Spring - Summer	х	x	
Scant Pomaderris (<i>Pomaderris</i> queenslandica)	E	-	Vegetated areas, particularly where gum trees (Eucalyptus) are present	Shrub	Spring - Summer	х	х	
Silky Swainson-pea (<i>Swainsona</i> sericea)	V	-	Grassland, Woodlands with <i>Eucalyptus</i> spp. and <i>Callitris</i> spp.	Groundcover	Spring	х		
Austral Toadflax (<i>Thesium australe</i>)	V	V	Grassland, Grassy Woodlands, areas with Kangaroo Grass (<i>Themeda australis</i>)	Groundcover	Spring	х		
Bluegrass (Dichanthium setosum)	V	V	Grassland, Grassy Woodlands	Groundcover	Summer		х	
Finger Panic Grass (Digitaria porrecta)	E	-	Grassland, Grassy Woodlands, areas with Eucalyptus albens	Groundcover	Summer		х	
Rodd's Star Hair (<i>Astrotricha roddii</i>)	Е	Е	Woodlands	Shrub	October to February	х	х	
Heath Wrinklewort (<i>Rutidosis</i> heterogama)	V	V	Open Forests, disturbed areas	Groundcover	Mainly Autumn			х

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page F5

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

Common Name (Scientific	ВС	EPBC	Habitat Targeted	Stratum	Flowering	Survey period		
Name)	Act	Act		Targeted	Period	Spring (September 2018)	Summer (December 2018)	Autumn (March 2019)
Ovenden's Ironbark (Eucalyptus caleyi subsp. ovendenii)	V	V	Grassy Woodlands, paddock trees	Canopy	July to September	х		
Narrow-leaved Black Peppermint (Eucalyptus nicholii)	V	V	Grassy Woodlands, paddock trees	Canopy	Autumn	x	Х	х

Notes: * Tree identification based on trunk and leave characteristics

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page F6

Assessment Report
BAM PLOTS DATA AND PHOTOGRAPHS

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

BAM Plot field data sheets

A total of 44 BAM Plots were undertaken as part of this study. Field data sheets used included:

- Hard Copy: Field data sheet printed in paper were used to collect data in a total of 35 BAM plots (i.e. plot P1 to P5 and P15 to P44). This section presents scanned field data sheets of those BAM plots.
- Electronic: Electronic field data sheets was created in Survey 123 (ArcGIS). A tablet was used in the field to collect data for nine BAM plots (i.e. plots P6 to P14). Data for BAM plots collected with electronic data sheet are presented in tabular form in this Appendix.

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page G1

BAM Plot - Field Survey Form

Date: 11 Sept 18

Project No: _____

Sheet A

Recorders: 10 + Adriana Plot ID:

Project Name: Bonshaw Solar Zone ID: ___

Page 1 of ____

	56.	IBRA region:	Nandow	test							
Datum:		IBRA sub-region:			W - C.	e dos					
Likely Vegeta	ation Class:	iotor sub region.	NOW IL TO	OF INDE	The In	week			Confi	dence	
, 6									н	M L	
Plant Comm	unity Type (PCT):	Silver Iron	care w	hile c	yoress		EEC:			dence M L	
		BAM Neste	Walk Take	1.16.1.0	0.				LASAN	is Mas	
(km)						lots			ANY - IL Alit!		
		1				1015	_				
X mark	LP1	LP3			LP5		20n	n x 50m		(Salte) red	
		-				1-1					
						[]	20n	1 x 20m			
am					midline						
		m 1					1m	x 1m			
com-		LP2	LP	4							
		1									
Om											
			-								
Umi		20.00			35 m						
ot 20m x 50r	n - midline	201111									
	ii - iiiidiiiie		-1	_		-		- 1	_		
Point on	Way Point	Easting	Norti	hing	Orientatio	n*	Photo - H	Horizontal	Photo	- Vertical	
midline	110,10111	20011118					T HOLO TIONZONIA		100000000000000000000000000000000000000		
0 m	ATS				86 E.		-		V		
50 m	A76.										
Aagnetig bearing	obtained with compas a	long the midline									
	Variable Committee				1						
AM Attribute	(1m x 1m nlots)										
AM Attribute	(1m x 1m plots)	1	2		3	4		5			
Jak	LP	1 / 0 i	2	/n	3	101		5		Average	
AM Attribute Attribut	e LP WP	1 LOI	2 LP2	493		LP 4		5		Average	
Attribut	e LP WP Photo	401	42	7	3	4				Average	
Attribut	e <u>LP</u> WP Photo	5	42	70	3 -	80		20		Average	
Attribut itter Cover (% are Ground (9	e LP WP Photo)	5 85	42 35 50	70	3 -	4				Average	
Attribut itter Cover (% are Ground (% ryptogam Co	e WP Photo) (6) Ver (%)	5 85 2	42 35 50 3	70 20	3 -	194 80 15		20 70		Average	
Attribut itter Cover (% are Ground (% ryptogam Co	e WP Photo) (6) Ver (%)	5 85	42 35 50	70	3 -	80		20		Average	
Attribut itter Cover (% are Ground (9 ryptogam Co ock Cover (%	e	5 85 2	42 35 50 3 5	70 20 5	3 .	194 80 15		20 70	line of the		
Attribut itter Cover (% are Ground (% ryptogam Co ock Cover (% iter cover is assess	e	5 95 2 8 entage ground cover of li	42 35 50 3 5	70 20 5	3 .	194 80 15		20 70	line of the		
Attribut itter Cover (% are Ground (% ryptogam Co ock Cover (% iter cover is assess	e	5 35 2 8 entage ground cover of lint;	35 50 3 5 tter recorded from	70 20 5 5 5 of five 1m x 1m	n plots centered at	80 15 5 5, 15, 25,	35 and 45n	20 70 10 -	line of the		
Attribut tter Cover (% are Ground (9 ryptogam Covock Cover (% ter cover is assess = Litter Plot (1m)	e	5 8 entage ground cover of lint; graphy + site features	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m	n plots centered at	80 15 5 5, 15, 25,	35 and 45n	20 70 10 -	line of the		
Attribut itter Cover (% are Ground (9 ryptogam Cov ock Cover (% iter cover is assess e = Litter Plot (1m	e	5 35 2 8 entage ground cover of lint;	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m	n plots centered at	80 15 5 5, 15, 25,	35 and 45n	20 70 10 -	line of the		
Attribut itter Cover (% are Ground (9 ryptogam Cov ock Cover (% tter cover is assess = Litter Plot (1m orphological Typ	e	5 8 entage ground cover of lint; graphy + site features	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m	n plots centered at	80 15 5 5, 15, 25,	35 and 45n Zone (opi	20 70 10 - n along the mid	line of the		
Attribut itter Cover (% are Ground (9 ryptogam Cov ock Cover (% tter cover is assess = Litter Plot (1m orphological Typ	e	5 8 entage ground cover of lint; graphy + site features Landform Element	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m Landfor	n plots centered at	80 15 5 5, 15, 25,	35 and 45n Zone (op)	20 70 10 - n along the mid			
Attribut tter Cover (% are Ground (9 ryptogam Covock Cover (% ter cover is assess = Litter Plot (1m orphological Typ hology	e	5 8 entage ground cover of lint; graphy + site features Landform Element	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m Landfor	n plots centered at	80 15 5 5, 15, 25,	35 and 45n Zone (op) M Sc	20 70 10 - n along the mid tional) iicrorelief bil Depth			
Attribut tter Cover (% are Ground (9 ryptogam Covock Cover (% ter cover is asses = Litter Plot (1m prphological Typ hology	e	5 8 entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m Landfor Soil Col	n plots centered at	80 15 5 5, 15, 25,	35 and 45n Zone (op) M Sc	20 70 10 - n along the mid			
Attribut tter Cover (% are Ground (% ryptogam Co ock Cover (% ter cover is asses = Litter Plot (1m orphological Typ hology	e	sentage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m Landfor Soil Col	n plots centered at	80 15 5 5, 15, 25,	35 and 45n Zone (op) M Sc	20 70 10 - n along the mid tional) iicrorelief bil Depth istance to near			
Attribut itter Cover (% are Ground (% ryptogam Co ock Cover (% tter cover is asses = Litter Plot (1m forphological Typ thology	e LP WP Photo) (6) Ver (%)) sed as the average percox 1m plot); WP =Waypoi Physion	sentage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	35 50 3 5 tter recorded from	70 20 5 5 of five 1m x 1m Landfor Soil Col	n plots centered at	80 15 5 5, 15, 25,	35 and 45n Zone (op) M Sc	20 70 10 - n along the mid tional) iicrorelief bil Depth istance to near			

Other
Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Soil Erosion

Fire damage Storm damage Weediness

Firewood | CWD removal Grazing (identify native/stock)

BAM Plot - Field Survey Form

Sheet B

Page 2 of ____

Date: 11/9/18	Recorders: Jo + Adriana	Plot ID:
Project No:	Project Name: Bonshaw Solar	Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree	Stems Count #	Stems with Hollows				
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^			
80 ⁺ cm							
50 - 79 cm	1		1				
30 - 49 cm		1					
20 - 29 cm		WH 1 = 6		1			
10 - 19 cm		20+10 = 30					
5 - 9 cm		40					
<5 cm ^b		collistris V	n/a	n/a			
ngth of logs (m): (≥10 cm diameter, >50c	min 120 540 540	230 366 270	120 =2186			

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is \leq 10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
/						
Callitris / stag	23	HI			1	-
silver ironbark	59	H2			10	-
staa	38	H3			6	
J						

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

00 m ²	plot: Sheet _\ of _	Survey Name	Plot Identifier		Re	ecorders	ACM /	JW
Date	11 09 2018	Bonshaw	Plot 1					
GF ode		each growth form group: Fu ic species: Full species nam	Il species name mandatory ne where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Collitris sp	(Cipresi) Callitris	glavcophylla (white Gpress	Din) N	30	2+28		
	Iron bark (Ev.	willy analom enterin	(Silver-bours brombare)	N	15	1.		
	SI - Asterce	ecc 36.710, 60. +10	0 0 30 70 30 5 17	*	0.5	7 1000		
	S2 Urtra	wens (Small Wel	tle) 0;00;10; 1x	*N	0.1	15		
	Grass I Thomas	du tempora (Kongaroo Gr	uss) 10, 10,00 ;50	N	0.2	410		
	Close L - Oxal	is percenans	1/ 10,50	N	0.1	62		
		SAL +1 . 000 0 : 1 . E		N	0. 2	2000		
	NI - Tribbur	to the franches and sale		オ製	0.1	14		
	Qualia? Opent	ia contraction (Ti	ger Pear) IN XXIX	*	6.1	A		
	WZ - Hypost	nacris radicata (Flatur	10 · /	*	6.1	112		
	Gross 2 Sport	rolos go Tripages	long lamie = 13, / MK	N	6.1	300		
	N2 Pimelea no	wa ranglica		N	0.1	1		
	& Close 2 = Trifal	ium· sp2.	. 20: 11 %	3K	6.	122		
	W3 - Solanum	sp. 1	10,1	*	0.1	12		
	Gross 3 Porms	so -		*	0.1	1		
		molle (Cranesbill G	examium) 1	*	0.1	1		
	A N3 Callibris of	How cophyla (regeneration or	nostly eaten) //	N	0.1	2		
	WS - Gamochael	50.1	20	便臣	0.1	26		
	N4 - E-oxdia	In Ams		N	0.1	1		
	1 Ortina or	ens (Simil Nottle		*	0.1	195		
	Scheroluena	birchii		. N	_	X		
								4.
							4	
	*							
							- 11	

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m Abundance: 1, 2, 3, ..., 10, 20, 30, ..., 100, 200, ..., 1000, ...

Sheet A

Page 1 of ____

Date:	09	18	_ Recorde	rs: <u>Jo</u>	, Adm	ana		_Plot ID:	2
Project No	o:		Project I	Name: 1	Bonsk	in Solo	XIK	_Zone ID:	
		ion Requ	uirements for BA	M Plot su	irvey				
Zone:	Do.		IBRA region:	Mond	A WAT				
Datum:			IBRA sub-region:	North	exim No	Then Come	lex		
Likely Vegeta	ation Cl	ass:			300	1			Confidence
Plant Comm	unity T	ma (DCT).							H M L Confidence
Plant Comm	unity iy	/pe (PC1):	Grey Box 1	Grassy We	re bullboo	open for	EEC:		H M L
10.01			BAM Neste	d Plots	Requirem	ent			
Plot 20m x 50r Point on midline		ine y Point	LP2 Easting	N	LP4 Jorthing	Orientation	-1 _! 2	Om x 50m Om x 20m m x 1m - Horizontal	Photo - Vertical
50 m		88	151, 33768°		20946	244°N		V	V
Magnetig bearing				1 47	2 20110				
BAM Attribute	(1m x	1m plots)							
Attribut	e	WP Photo	1 LP/	ip2		3 JP 10	1)	13	Average
Litter Cover (%	5)		100.	80.	6±		70.	60	2.
Bare Ground (9	%)	1/2							
Cryptogam Co									
Rock Cover (% Litter cover is asses LP = Litter Plot (1m	sed as the); WP =Waypo	centage ground cover of li pint; ography + site features						line of the 20m x 50m p
Morphological Typ	oe		Landform Element		Landfo	rm Pattern		Microrelief	
ithology			Soil Surface Texture		Soil Col	our		Soil Depth	
Slope			Aspect		Site Dra	ainage		Distance to near water and type	est
Plot D	Disturbano	ce	Severity Code A	ge Code			Observational Ev	idence	
earing (incl. loggin	g)								
ultivation (incl. past oil Erosion	ture)	V							
rewood CWD ren	noval								
razing (identify nat	tive/stock)	1							
re damage torm damage									
/eediness			(1 1						
ther									

Plot 20m x 50m: Tree Classes a

Sheet B Page 2 of ____

DBH Class#	Ti	ree Stems Count #		Stem	with Hollows			
DBH Class	Euc*	Non-	·Euc^	Euc*	No	n-Euc^		
80 ⁺ cm	1		1					
50 - 79 cm	11		1					
30 - 49 cm								
20 - 29 cm	1).	1)	-					
10 - 19 cm		1						
5 - 9 cm		8						
<5 cm ^b		40	40 n/a		n/a			
Euc - Record of living of Non-Euc - Record of li Count apply when the name of the largest living stem is	native eucalypt trees (include ving native non eucalypt tree umber of tree stems within s included in the count/estim	es a size class is ≤10. Estimates o nate. Tree Stems must be livir	can be used when >10 (eg	10, 200, 30,, 100, 200), 300). For a multi-ste	mmed tree, only		
SAM Attribut	DE	l WP	n x 50m) Easting	Northing	Hollow <20 cm	Hollow >		
Grey Bo	x 14				1111	1		

Date: 11/09/18 Recorders: 10 + Adnana Plot ID: 2

Project No: _____ Project Name: Bonshaw Solar . ____ Zone ID: _____

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

- Daniel (1971)	olot: Sheet _ of _	Survey Name	Plot Identifier		Re	ecorders		
Date	8103 PO Q1	0470861-Bonshau	2	ACM,	JW			
GF Code	Top 3 native species in All other native and exo	each growth form group: Full stic species: Full species name	species name mandatory where practicable	N, E or HTE	Cover	Abund	stratum	vouche
T	Grey Box Even	lyptus musocarea		2	30	13		
T		alyphos molomophila	Silver-leved frombare	N	20	- 1		
		us sp. 1		N	0.3	+		
	Small Tree 2 N	otalaza longifalia		N	30	47		
		otelaca muscarpa		N	0.1	1		
	NI small shoot	up to Robertlanh P: Hosposum	spinescens / 114	N	1.6	L		
	Grass 1 50 T	ripogen longiformis	0 19 10 /11	N	0.1	123		
	Clove I & flower	Oxalis executions	+; 5; 10; 11	N	0.1	119		
		posobolus «longutus		2	0.1	2		
	WID	Asterniane sp. 2	• 10	*-	0.1	110		
				*	0.1	30		
		Euphorbia sp. 1		13.*	0.1	21		
	N3 1	Pirmalea neo-anglica Melichrus sp. I	1//	N	8.1	4		
				N	1.0	1		
		itris glucophylla (U		N	0.1	3		-
	G1005 3 20 5	Trifolium sc. L	60: //	N	0.1	62		
	Close 2	Vine Parsonia sel	1 12	*	0.\	2		
-				- 12	0.1	124		
	Opention Come as	Phts Openha auran	itiaca /	E	0.1	1		
	VI & Elem	Hardenbergia violacene	6 /	N	6.1	11		
	V2	Pardorea gundorana	10 + /	N	0.1	11		
	N6 Mune	THE Orchondra se.	Α ///	N	0.1	3		
-								
			(9)					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ... 100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4×1.4 m, and $1\% = 2.0 \times 2.0$ m, $5\% = 4 \times 5$ m, $25\% = 10 \times 10$ m

Abundance: 1, 2, 3, ..., 10, 20, 30, ..., 100, 200, ..., 1000, ...

Sheet A Page 1 of ____

Project No		Project I	Name: <u>Boo</u>	show	Solar		Zone ID:		
riojective		_ 110,000	vanic.	OI TOTO			Lone ib.	-	-
Desktop Inf	ormation Requ	irements for BA	M Plot survey						_
Zone:		IBRA region:	Nondens	:V-					
Datum:		IBRA sub-region:		r North	a Camel				
Likely Vegeta	tion Class:		T OCM TELY (IST.)	1 144 1111	ii Sauta			Confidence	9
							-	н м	
Plant Commu	inity Type (PCT):	? Rough -b	inved Apple -	White Expess P	ino - Blukely	EEC:		Confidence H M	
		BAM Nest	ed Plots Req	uirement				wl	
(0 m =					Plots			4 mo	di
	LP1	LP3		LPS			n x 50m		
2 777							ii x soiii		
		1			i_		m x 20m		
(XIII)				m	idline	1 100	x 1m		
Vivia 1		LP2	LP4			Till	I X TIII		
		LPZ	LP4						
0.01									
		Section 1	(7)	- Albam					
Det		SOm.	m	-03-00	None				
			m	-4-0					
			Northi			Photo -	Horizontal	Photo - Ve	rtical
lot 20m x 50n Point on	n - midline Way Point	20 n		ing Oi	rientation*	25.25.55	Horizontal	Photo - Ve	rtical
Point on midline 0 m	way Point	Easting		ing Oi	50.00	25.25.55	CAN ROOM CITY		rtical
Point on midline 0 m	n - midline Way Point	Easting		ing Oi	rientation*	25.25.55	CAN ROOM CITY		rtical
Point on midline 0 m 50 m Magnetig bearing	way Point	Easting		ing O:	rientation*	25.25.55			rtical
Point on midline 0 m 50 m Magnetig bearing	Way Point 190 obtained with compas a (1m x 1m plots)	Easting 151. 3403 statements and the midline	29.210	ing Oi	rientation*	4	5		
Point on midline 0 m 50 m Magnetig bearing	way Point 190 obtained with compas a (1m x 1m plots) LP WP	Easting 151. 3403	5 29.210	ing O:	rientation*	4			rtical
Point on midline 0 m 50 m Magnetig bearing AM Attribute	way Point 190 obtained with compas a (1m x 1m plots) LP WP Photo	Easting 151. 3403 statements and the midline	2 20	3 21	rientation*	4	5 2 5		
Point on midline 0 m 50 m Magnetig bearing AM Attribute Attribute	way Point 190 obtained with compas a (1m x 1m plots) LP WP Photo)	Easting 151. 3403 statements and the midline	2 20 20 20	3 21 20	rientation*	4	5 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2		
Point on midline 0 m 50 m Magnetig bearing AM Attribute Attribute itter Cover (% are Ground (%)	way Point 190 obtained with compas a (1m x 1m plots) LP WP Photo)	Easting 151. 3403 statements and the midline	2 20	3 21	rientation*	4	5 2 5		
Point on midline 0 m 50 m Magnetig bearing AM Attribute Attribute itter Cover (% are Ground (% ryptogam Cover))	way Point 190 Obtained with compas a (1m x 1m plots) LP WP Photo Photo yer (%)	Easting 151. 3403 statements and the midline	2 20 20 20	3 21 20 40	rientation*	4	5 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2		
Point on midline 0 m 50 m Magnetig bearing AM Attribute Attribute itter Cover (% are Ground (% ryptogam Covock Cover (%))	way Point F90 obtained with compas a (1m x 1m plots) LP WP Photo) 6) ver (%)	Easting 151, 3403 statement of the midline 1 19 19 15 15 20	2 20 20 20 20 40 50	3 21 20 40	rientation* 24° NE	4	5 25 70 5	A	erag
Point on midline 0 m 50 m Magnetig bearing AM Attribute Attribute itter Cover (% are Ground (% ryptogam Covock Cover (%) tter cover is assessing to the cover is a cover in the cover is a cover in the cover in the cover in the cover is a cover in the cover in	way Point 190 Obtained with compas a (1m x 1m plots) LP WP Photo) (6) Ver (%) sed as the average perc x 1m plot); WP = Waypo	Easting 151, 3403 statement of the stat	2 20 20 30 20 itter recorded from f	3 21 20 40 ive 1m x 1m plots c	rientation* 24 ° NE	4 2 2 2 5, 35 and 45	5 90 5	A	erag
Point on midline 0 m 50 m Magnetig bearing AM Attribute Attribute itter Cover (% are Ground (% ryptogam Covock Cover (%)) tter cover is assess P = Litter Plot (1m)	way Point Tyo obtained with compas a (1m x 1m plots) E WP Photo) (6) yer (%) sed as the average perc x 1m plot); WP =Waypo Physio	Easting 151. 3403 : slong the midline 1 19 20 entage ground cover of lint; graphy + site feature	2 20 20 30 20 itter recorded from f	3 21 20 40 ive 1m x 1m plots c	rientation* 24 ° NE 10 50 4 entered at 5, 15, and Manageme	25, 35 and 45	5 90 5 m along the mic	A	erag
Point on midline 0 m 50 m Magnetig bearing Attribute Attribute itter Cover (%) are Ground (%) cryptogam Cover (%) tter cover (%)	way Point Tyo obtained with compas a (1m x 1m plots) E WP Photo) (6) yer (%) sed as the average perc x 1m plot); WP =Waypo Physio	Easting 151, 3403 statement of the stat	2 20 20 30 20 itter recorded from f	3 21 20 40 ive 1m x 1m plots c	rientation* 24 ° NE 10 50 4 entered at 5, 15, and Manageme	25, 35 and 45	5 90 5	A	erag

Morphological Type	Landform Element	Landform Pattern	Microrelief
Lithology	Soil Surface Texture	Soil Colour	Soil Depth
Slope	Aspect	Site Drainage	Distance to nearest water and type

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)			
Cultivation (incl. pasture)			
Soil Erosion		1 1 1 2	
Firewood CWD removal			
Grazing (identify native/stock)			
Fire damage			
Storm damage	- 14		
Weediness			
Other			

Seventy: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Recorders: ___

Sheet B

Page 2 of

Plot ID: 3

roject No: _	Proje	ct Name:	Zone ID:			
ot 20m x 50	m: Tree Classes ^a	*	8			
DBH Class#	Tree Ster	ns Count #	Stems wit	h Hollows		
DDIT Class	Euc*	Non-Euc^	Euc*	Non-Euc^		
80 ⁺ cm						
50 - 79 cm	1		1			
30 - 49 cm	1					
20 - 29 cm		t				
10 - 19 cm) .				
5 - 9 cm		L				
<5 cm ^b	1	40	n/a	n/a		
ngth of logs ((m): (≥ 10 cm diameter, >50cm in	530 150 40	00 90 620 33	= 371		

Date:

Count apply when the number of tree stems within a size class is \leq 10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
ronbark (creba?)	65	24			1	11

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

Lloing

00 m ²	plot: Sheet _ of _	Survey Name	Plot Identifier		Re	ecorders		
Date	10 09 2018	0470261- Benchaw	3	ACM	15W			
GF Code	Top 3 native species in All other native and exo	each growth form group: Full tic species: Full species nam	species name mandatory e where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Ironback (E-evel	(10)		N	25	1	Caner	
	Red from (E. Hlente	8 07 - 2019 T	in Colley continued the	N	20	112	Congey	
A	Callistus glavcophy	Ma (from < 70 can to Small	tion	N	0.2	1+ 43	7.50	
	1 N2 - Pinchea	was and lieu Jes	1/1		0.1	143.3	Sheb	
		a se. L		*	0.1	600	Grand	
	Fern 1		1 XXXXX		0.1	22	Growns	
	a Dountro (some)	Opentia warantiawa		*	0.1	2	Good	
	1 N1 Ozothum	nus cascinioides	50, 1//		0.1	17.3	Growd - Shoul	
		ostis sp. 1	lo	*	0.1	10	groord	
	WISO S	enecio sp. 1	/ ///	*	0-1	6	good	
		+2) & Tuldium		*	1.0	6	ground	
	Clove I (worker)	Schrobena birch	8; 30; 34	N	0.1	28	ground	
	W2 trus	Urtice urens	10 // XX	*	0.1	32	ground	
	N3 = @ but .	grazed to ground		N	6.\	1	ground	
		= Closed Patt) Oxulas ce		N	0.1	8	groond	
	W3 Plantag	o set		*	1.0		grunt	
		ge. 1	// ///	N	0.1	7	ground	
	WPasiall	ot 2 Astoracana se. 1	/	*	0.1	- 1	ground	

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'. Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ... 100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4×1.4 m, and $1\% = 2.0 \times 2.0$ m, $5\% = 4 \times 5$ m, $25\% = 10 \times 10$ m Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Occiall - base ground
So.l + Silly - growthy Shale froguests
Ant? holder
rathit

791 @ som

Sheet A

EEC:

Page 1 of ____


H M L

H M L

Date: 12-09 - 2018	Recorder	S: ACN / JW	Plot ID:
Project No: 0470861	Project Na	Zone ID:	
Desktop Information Re	quirements for BAN	Plot survey	
Zone:	IBRA region:	Nandenan	
Datum:	IBRA sub-region:	Nunderer Northern Condex	
Likely Vegetation Class:			Confidence

BAM Nested Plots Requirement

Cleared - Non-nutice



Plot 20m x 50m - midline

Plant Community Type (PCT):

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
50m	7.96	0338834	6768241		1	-
50 m ()	F96 797	0338827	6768189	10 N	J	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	WP Tag	PPF	800	801	602	Average
	Photo	V	V	V	V	V	
Litter Cover (%)		15	2	0	85	80	
Bare Ground (%)		60	95	890	5	10	
Cryptogam Cover (%	6)	5	5	5	D	0	
Rock Cover (%)	10.	0	2	D	0	0	

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determinng PCT and Management Zone (optional)

Morphological Type		Landform Element		Landform Pattern		Microrelief	Hat
Lithology		Soil Surface Texture		Soil Colour	light orange	Soil Depth	
Slope	gont Class	Aspect	N	Site Drainage		Distance to nearest water and type	Greek to bust

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)			
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)			
Fire damage			
Storm damage			
Weediness			
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Sheet B

n/a

Page 2 of ____

n/a

Date: 12-69 -	Reco	rders: ACM /JW		_ Plot ID:4
Project No:	Proje	ct Name: 0 470861		Zone ID:
Plot 20m x 50m:	Tree Classes ^a			
DBH Class [#]	Tree Ste	ms Count #	Stems w	rith Hollows
DDH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm	-			
50 - 79 cm	9			
30 - 49 cm	/e.i			
20 - 29 cm	_			
10 - 19 cm	200			

Length of logs (m): (≥10 cm diameter, >50cm in

5 - 9 cm

<5 cm b

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
	14					
				1		
	1					
						+
	-	-				+

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² j	olot: S	heet _	of _	Survey Name	Plot Identifier	Recorders	
Date	12	09	2018	0470861- Bonshow	4	ACM , JW	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouche
	Kanyaran Grass Thomada triandra . 20 /	N	1.6	200		
	Cxolic potentians / XX 5, 10	2	0.1	37		
	Kidon word (4-1) Dicharder races 10:5 MK	2	1.0	20		
	Rock Form Chailantes sceberi 1000 MX, W. XX	N	0.1	240		
	WI (Mile) strong -or Orfice overs 20: 1/ 1/1/2	*	0.1	27		
	Rock Form Cheilantes sceberi 10° Mx, 10, XX W1 (Min) strong on Ortica overs 20; MX Grass 1 * Dicharthium satosum (Bluegrass)	N	0.2	202		
	Trifolium so L X	*	6.1	3		
	Opento avantiona III	*	0.1	4		
	Opento aucontione N1 - Engaceae (No) Ponto acconglica 9: 11/14 W2 Pintos Goodenia sp. 1 Cartague (H) Solero lacena birchii 2:14	N	0.	17		
	WZ Pods Godania sp. 1	N	0.1	2		
	Cartague (Schoolagna birchii 2:14	N	D.\	16		1
	Costus te Opuntia tomentosa /	*		X		
	W3 Wahlambergia granificala //	N	0.1	3		
	Callitres go L (groves)	*	0.1	- (
	Callitres op 1 (great)	N	0.1	1		
	Grass 2 W Sporobolus elongatus (Slander Rut's Tail	N	0.1	5		
	WS Euphorbia dallachyana	N	0.1	1		
	Pea flower Bossiaga scortectinii	N	0.1	3		
	Cyporacoas - Schoons sp. 1	N	1,0	1		*

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A


Page 1 of ____

Date: 12-09 -2018	Recorders: A(w \ I w	Plot ID: 5
Project No: <u>64 70861</u>	Project Name:	Sonohene Solar Farm	Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	IBRA region:	Nonlawar				
Datum:	IBRA sub-region:	Nondown	Neithan	Comebx		
Likely Vegetation Class:						Confidence H M L
Plant Community Type (PCT)	:				EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	804	0339451	6768659	239° SW	1	V
50 m	803	0339402	6768675	20105	~	/

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	29	30	3/	32	33	Average
	Photo	V	V	V	V	V	
Litter Cover (%)		82 -	90	85	20	98	
Bare Ground (%)		3	6	to 15	6	2	
Cryptogam Cover (%	6)						
Rock Cover (%)		15	4	#-	\$ 74	-	

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determinng PCT and Management Zone (optional)

Morphological Type	Landform Element	Landform Pattern	Microrelief
Lithology	Soil Surface Texture	Soil Colour	Soil Depth
Slope	Aspect	Site Drainage	Distance to nearest water and type

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)			
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)			
Fire damage			
Storm damage			
Weediness			
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Sheet B

Page 2 of ____

Plot ID: ____5__

DBH Class#	Tree Sten	ns Count #	Stems wit	h Hollows	
DDIT Class	Euc*	Non-Euc^	Euc*	Non-Euc^	
80 ⁺ cm					
50 - 79 cm	1				
30 - 49 cm	1(1)		1		
20 - 29 cm	M M M II		1		
10 - 19 cm	1]]				
5 - 9 cm	1,	T			
<5 cm ^b	✓	✓	n/a	n/a	

Recorders:

Date:

Count apply when the number of tree stems within a size class is \leq 10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
red/ Grey Gum?	38	34			11	
red/ Grey Gum?	20.	35			1	
as above?	23	36		5	1	
as above?	54	37			11	1
					+	
		5 0			12	
				/		
4			1			
-						

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

00 m ² p	olot: Sheet _ of _	Survey Name	Plot Identifier		Re	ecorders		
Date	12-09-2018	0470961- Banshiw	5	Ac	n 151	N		
GF Code	Top 3 native species in All other native and exo	each growth form group: Full tic species: Full species name	species name mandatory e where practicable	N, E or HTE	Cover	Abund	stratum	vouch
	Gum 1 (Eucalypte	a deallab Frample) (Ju	enite < 30cm byth Drews	N	10	120+6		
		ia sp. P. Leathat		N	15	2		
		ypress grazed down to 6 room		N	0.2	166		
				N	0.1	2		
	Opintra aurant	these	(1)/	*	0.1	4		
	Greise & (perensial)	Sporobolus alongatus	; 12	*N	0.1	12		
	Kidny ward & (Poly	Dichondra vegens	700, 20: //	N	0.1	222		
	Rock Form Charles	nthes sie ber	· ;7,2:10, 11.3;6	N	6.1	133		
	N2 Gooden	ice squ	1 11	N	Oil	3		
	Oxalis permans			N	0-1	2		
	Grass 2 Trapou	yon langilosmis	7 1000 /L/X	N	0.1	1000		
		200 36		N	0.1	4		
	WZ Und	Lovinined	1	*	al	2		
	Vine 1 Pand	lorea panderana	1	N	0.1	1		
	Shrub I Doc	lonum visiosa subsp. spal	white /	N	1.0	1		
	Raspalluluria a	disondens		N	X	X		
	Corymbyase				X			
			*					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'. Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

mation Required in Class:		for BAM I	Plot survey	Northern (Zone ID	
94 n Class:	IBRA regi	ion:	Nemdowar	Northern (amplex		
94 n Class:	T 7 9 7 10	To the sale		Northern (amplex		
n Class:	IBRA sub	-region:	Nondovar	Northern (complex		
					1		
ty Type (PCT):							Confidence H M L
					E	EC:	Confidence
	BAM	Nested	Plots Require	ement			
LP1	1	LP3		LP5	Plots	20m x 50m 20m x 20m	20×50 V
	LP2		LP4	midline	п	1m x 1m	
			Į.	olem J			
		LP1	LP1 LP3	LP1 LP3 LP4 LP4	LP1 LP3 LP5 midline	LP1 LP3 LP5 I - 1 L - 1	LP1

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1458	0338451	6767511	1930 SE	~	
50 m	1464	0338482	6767479	W 282		~

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1450	1460	1461	1462	1463	Average
	Photo	/	V	V	V		
Litter Cover (%)		60	30	80	80	95	69
Bare Ground (%)		35	74.0	18	20	S	29.6
Cryptogam Cover (%	6)	- 1	0	1	7	7	1.4
Rock Cover (%)		5	0	Z	0	0	1.4

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	Hillslope	Landform Pattern	ran Aus - but	Microrelief	
Lithology		Soil Surface Texture	Hosth Loof litter	Soil Colour	light brown	Soil Depth	,
Slope		Aspect	Total Time Inter	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence	
Clearing (incl. logging)	2	0	Stume prient	
Cultivation (incl. pasture)				
Soil Erosion				
Firewood CWD removal				
Grazing (identify native/stock)				
Fire damage				
Storm damage				
Weediness				
Other				

Sheet B Page 2 of ___

Date:	25-03-2019	Recorders:	ACM, TZ	Plot ID:\S
		The second secon		

Project Name: Bonshaw Solar Farm Zone ID: _ Project No: 0470861

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stem	s Count #	Stems w	rith Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80⁺ cm	1 = 2		11 = 2	
50 - 79 cm	HL 1 = 6	~ 4	JUN =5	
30 - 49 cm	*	<i>III</i> = 3		-
20 - 29 cm		Ht/ =6	\$	
10 - 19 cm	9	6, M, 5, 4, 4, 11 = 26	#	
5 - 9 cm		11,8; \$ 6 = 25	#	
<5 cm ^b		7,38,6,18,20=89	n/a	n/a
gth of logs	(m): (≥10 cm diameter, >50cm in		, 3, 4, 9, 10,5,6	= 66

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
Sibre Juneal Trophoris	83	1465	6338478	6767484	2 hollows	
O	49.5	146	0338477	484437	1	
ħ	8.7	1467	0338461	6767987	414100 / 1x 1501	
(\	52	1468	0338454	6767514	1 x 10 an	
11	51	1469	0338459	6767506	La Clora	
E-albas	72	1470	0338465	6767503	1	-
Enthers	54	14 71	033 8478	6363491	1	
*						

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 25/3 /2019	Bonshaw	15	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouche
	Eucalypho melani philoia		20	3	T	
	Eucalyptus albens		ح		T_	
	Callities glancophylla		13		S	ļ
	Callitris glancophylla Princha stricta		0.)	21		
	Chloris truncata	,	0.1	10		
			0.1	10		
	Avistina ramosa.		0 (1			
	Cymbopogon refractus		0:12	1.0		
i	Cyperus curacitis		0 = 1	5:0		
	Spuntia auracantiaca		0.1	4		
	Digitaria diffusa		0.1	20		
	Cymbopogon refraction Cyperus cyracilis Opuntia auracantiaca Digitaria diffusa Austrostija scabra		0.2	100		
	El Maria Mutans		0.1	10		
	Cheilanthes sieberi		0.1	7		
	Rostellularia adscendens		01	100		
	Sida corrugata		0.1	50		
	Vi Hadinia sulcata		0.1			
	Notelaza microcarpa		0.)	6		
,	Glycine clamaestina		0.1	26		
	Boerhevia dominii		ا م)	İ	
	Enneapoyon gracilis		5 ، ن	600		
	Enneapoyon gracilis Sclerolaena birchii		0.1	1		
	Entolasia stricta		0.2	50	} 	
	Dichondra sp. A		0.1	3		
	Solanum ferocissimum		6.1			
	Solanum ferocissimum Pandorea pandorana		0.1	1		
	Bothrichlan decipión		011	7		
	Aristida leptopoda		01	5		
	, ,					
				ļ		
<u>.</u>			:			
						<u>-</u> .
			: 			
_·						
			: 			
-						

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A Page 1 of ____


Plot ID: 16 Date: 25-03-2019 Recorders: ACM, TZ

Project No: 047 0861 Project Name: Rowloaw Solar Form Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	5k	IBRA region:	Nandewar				
Datum:	qu	IBRA sub-region:	Nandowar	Nathern	Comocks		
Likely Veg	etation Class:				0,,,,,,,,		Confidence H M L
Plant Com	munity Type (PCT):	594				EEC:	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1472	033 8659	636 75 69	LSSOW		V
50 m	1473	0338619	6767587	118° SE	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5		
Attribute	WP	WP 14177 1	1479	1479	1479 1480	1981	Average	
	Photo	V	V	V	J	V	T CONTRACTOR	
Litter Cover (%)		80	90	85	10	1	53.289	
Bare Ground (%)		20	S	3	90	89	43.620.6	
Cryptogam Cover (%	6)		D	5	0	0	1.2	
Rock Cover (%)		7	5	10	0	5	3.6	

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Crest	Landform Element	Hillerest	Landform Pattern	11:11	Microrelief	eppen woodland
Lithology	Sind - chiny min in	Soil Surface Texture		Soil Colour	factual process	Soil Depth	~
Slope	N cool	Aspect		Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	C	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R O	
Fire damage			
Storm damage			
Weediness		R	
Other			

Severity: 0 = no evidence; 1 = light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B Page 2 of ____

Date:	75-03-2019	Recorders:	ALM. TZ	Plot ID:
		The second secon		

Project Name: Bonshow Solar Farm Zone ID: Project No: 0430861

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree	Stems Count #	Stems w	vith Hollows
JBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm	/ = _		/ = 1	
30 - 49 cm	JH =5		1/ = 2	
20 - 29 cm	93 11 = 2	JHT 111,5, = 13		
.0 - 19 cm	\$	10,43 # #1/=28		
5 - 9 cm		2,2,/ =5		
<5 cm ^b		1/ =2	n/a	n/a
gth of logs	(m): (≥10 cm diameter, >50c	min 1,10,7,1,7,6,3	,13; 4; 15; 3, = 70	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Holos - D wombat ??

Count apply when the number of tree stems within a size class is \$10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

E doubata (Tumbledown Red fun

E other / watercom Eved broath book Silentered lonbook

Roocky crest

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
liber drawd transport	43	1474	03 86 26	67-67-587	1	
wealyphus Campbala	44	(47)	0338646	6767582	Į.	1
E Jealhota	Sa	1436	0338636	6767572	7_	1

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 25/2/19	Bonshaw	, 6	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Encamptus dealbata Encamptus melanopaloia Encamptus alberr Callitra glaucophylla Hypericum spi Ozothamous carsenoides		15		+	
	Enculyatus melanopaloia		3		1	
	Eucalyptus albens		3	1_		
	Callitin glaucophylla		20	ļ	T	
	Hyparican sp.		0.)	20		
	Ozothaninus capsinoides		0.2	200		
	Gymbopogon refractus Aristida ramosa		0.1	15		
	Aristida ramosa		0.3	230		
	Dichandra Sp. A		0.3	400	<u> </u>	
	Dichandra sp. A Roste II ularia adscendens		0.3	200		
	Colucia Clanda 156		0.3	200		
	Sido coxtugado		0.2	200		
	Glycine clandestina Sida corrugata Boerhavia dominii		0.1	40		
	Comments of the comments		011	40	ļ	
	CA John Streethe		0.1	100	 	
	Funta autarantique		0.3	200		
	Cyporus gracilic Opuntia auracantiaca Enneapogon gracilic Primelia stricta		0.2	80		
	D. Stricta		 	·	<u> </u>	
	Parçonsia encalyptifolia Vitadinia sulcata		0.1	10		
	V(Radinia Sulcata		0.1	5		
	Sologyne belliaides Cheildithes suberi Carrosa >p.	<u>·</u>	011	40		
	(heil xithes suderi		01	50		
<u>-</u>	Carvala sp.		0.1	1		ļ
			ļ	 		
				ļ <u>.</u>	ļ	ļ
				<u> </u>		
				ļ		
		<u></u>				ļ
			,,,			
				<u> </u>		
			I			
		ļ	<u> </u>			
, —						
		 	-	-		
			 	1		
		 	 	 		
			ļ	-	 	
		 		1	 	ļ <u>.</u>
		<u> </u>		1		ļ

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF - circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A

Page 1 of ____

Date: 2	5- 63 7019	Recorders:	ACM, TZ	Plot ID:
A STATE OF THE STA				The state of the s


Project No: 047086 Project

Project Name: Bonshuw Salar Farm Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandowar					
Datum:	94	IBRA sub-region:	Mandaur	North	Comoles			
Likely Veg	etation Class:				L			Confidence H M L
Plant Com	munity Type (PCT):	594 1	// 17			EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1485	0338763	6767760	770°W	V	V
50 m	1486	0338720	6767 787	124°5E	~	

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1487	14.88	1489	1496	1441	Average
	Photo		V	V	2	V	
Litter Cover (%)		50	70	30	10	40	30
Bare Ground (%)		30.	80	68	50	50	55.6
Cryptogam Cover (%	6)	25	0	0	0	5	2
Rock Cover (%)		70	D	2	40	10	14.4

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	Low Hill	Landform Pattern	Hillstone	Microrelief	giossy asen
Lithology		Soil Surface Texture		Soil Colour	highl armys bow	Soil Depth	
Slope	& 70°5	Aspect	MW	Site Drainage	el III Latiner	Distance to nearest water and type	v 5m

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	R	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	2	12 -0	
Fire damage			
Storm damage			
Weediness	1	R	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B

Page 2 of

Date: _	25-03-	2510	
Date.	70 00-	2019	

Recorders: ACM, TZ

Plot ID:

Project No: 047086

Project Name: Bonshow John Farm

Zone ID: _____

Plot 20m x 50m: Tree Classes a

DBH Class [#]	Tree Stems	Count #	Stems wit	h Hollows
JULI CIASS	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm			/	
50 - 79 cm			-	
30 - 49 cm				
20 - 29 cm				
10 - 19 cm		1		
5 - 9 cm			-	1
<5 cm ^b			n/a	n/a
gth of logs (m)	: (≥10 cm diameter, >50cm in	3,2,3,1,3,=12		

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Cull off traces = (logs) = 2,61,7, 2,5,3,2,2,1; > Not touching ground. Branches an

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
	/					
/	+ +					36
/			- A			
1	1					
						1
						-
	17-1					
	1					
/	7					
/						
/						
					1	

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 25/3/19	Bonshaw	17	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Evolvalus akinoides		0.1	20		
<u></u>	Calliting glauco phylla Boechavia dominii		6-1	.30	S	
	Boerhavia dominii		<0.1	60		
· · ·	Villadinia sulcata:		6.1	30		
	Solanum terocossimum		0.1	7		
	Solanum ferorissimum Dichondra Sport		ر ، ک	100		<u>, </u>
	Sida corragata Glycine clandestina		0.5	600		
	alycine clandestina		0.2	70	ı	1
	Sole cube he Minister		0.1	30		
	Alistian ramosa		0.1	40		
	Alistian ramosa		3	50		
	Oxalis perennals	·	ر د ه	16		
	Xanthiam spininsum		0.1	1		
	Charther Stebers		.0.1	5		
	Calotis lappulacen		0.1	5		
*	Ennaposan crachs		5	200		
	Enneapogen gracilis Opuntia aurantiaca		01	20		
	Chamaerys, drummondi.		0.1	20		
	(marin projeti)		0.1	20		
	Cyperus ornalis Pimelia stricta		0-1	5		
	Bothriochlea macra		<u> </u>	20		
	Rostellularia adscendens		O.1	30		
	Helictropium amplexicante		0:1	50		
	Scleria mackaviensis		i	2		
	Scotia machaviansis		0.1			
						ı
						
						·
			,			
					 	
	· · · · · · · · · · · · · · · · · · ·				<u> </u>	
			•			
	·					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Grassland

594 - DN9

EEC:

Sheet A Page 1 of ____


M L

Confidence

H M L

		Recorders	ACM, TZ	Plot ID: 1 8
Project	No: <u>047 0861</u>	Project Na	me: Booshow Solar Form	Zone ID:
	No. of the last of	quirements for BAM	7	
Zone:	56	IBRA region:	Nandayar	
Datum:	94	IBRA sub-region:	Nandower Northern Compted	
Likely Veg	etation Class:			Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Plant Community Type (PCT):

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1492	033 83 77	67 67 858	530 NE	V	2
50 m	1493	033 84 22	67 67 888	S5do 2 M	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1494 1496	1997	1948	1949	Average	
	Photo	V	V	v	7	V	
Litter Cover (%)		75	60	40	40	30	49
Bare Ground (%)		25	39	60	60	70	50.8
Cryptogam Cover (%	6)	٥	0	0	0	0	0
Rock Cover (%)		0	1	D	0	0	0.2

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1 mx 1 m plots centered at 5, 15, 25, 35 and 45 m along the midline of the 20 mx 50 m plot.LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	flat	Landform Element	Flat	Landform Pattern		Microrelief	grastard
Lithology		Soil Surface Texture		Soil Colour	124 Hd - Frenk	Soil Depth	-
Slope	164	Aspect	NE	Site Drainage	SW	Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	2	R.O .	
Fire damage			
Storm damage			
Weediness		2-6	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent(<3 yrs); NR = not recent(3 - 10 yr); O = old(>10 yrs)

Sheet B Page 2 of ____

Zone ID: __

Date:	25-03-2019	Recorders:	ACM, TZ	Plot ID:18

Project Name: Bonshaw Inlus Farm

Plot 20m x 50m: Tree Classes a

DBH Class [#]	Tree Stem	s Count #	Stems with Hollows		
JBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^	
80 ⁺ cm					
50 - 79 cm	Y				
30 - 49 cm					
20 - 29 cm					
.0 - 19 cm			/-		
5 - 9 cm	/ .				
<5 cm ^b	/		n/a	n/a	
th of logs (m)	: (≥10 cm diameter, >50cm in	12, 1, 3, 15, 1	2, 1 = 35		

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is \$10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
						-
1						
	-					-
	-/-					
/						

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 25/3/2019	Bonshaw	18	

Date	25/3/2019 Bonshaw 10	<u> </u>				
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	, Aristida ramosa		5	120		
	Aristida ramosa Cheilanthes sieberi		0.2	500		
	Dichinary Sp. A		0.1	200		
	Glycine clandestine		0,5	500		
	Myrryian SD		0.1	30		
	Dichendry Sp. A Glycine clandestine Myperian Sp Side corrugata Oxalis perannans Sclerolaena birchii Viffadinia sulcuta		0.1	30		,
	Oxalis perennang		ا ، م	100		
	Sclero laena birchii		011	200		
	Vittadinia sulcuta		0.1	70		
	Cunerus pracilis		0'1	70		
	Cyperus gracilis Einadia nutans		0.2	200		
	Soleonne bellicides		0.1	15		
	Solanum ferocissimum Bothriochlou macra		0.1	10		
	Rothrighlas macro		1.5	200		
-	Primelia stricta		24	2		
	Combaness vatradis		0.1	3		
	Cymbopogon refractus Paspaliolium olistans		0.1	2		
	1 1:1. 11 -called		25%			
-	Bothriconoa bilaba Panicum simila	<u></u>	· '	3	ļ -	
	P		0.1			
	Tanicum simile		5.7	2		
				 		
		 	 	-		
						
						
		<u> </u>		 	 	
				<u> </u>		
			ļ	-		
				<u> </u>		
				<u> </u>		
				ļ	ļ	-
		 		-		
		<u> </u>		 		
				ļ		
		<u> </u>				
		 				ļ
]			

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF - circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A

Page 1 of ____

Date:	25.03 - 2019	Recorders:	ACM, TZ	Plot ID:


Project No: 047 0861

Project Name: Bonshow Solar Form Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandryyar				
Datum:	94	IBRA sub-region:	Nondowar	Northern	Comolox		
Likely Vege	tation Class:						Confidence H M L
Plant Com	munity Type (PCT):	DNG			EEC:	X	Confidence (H) M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1500	033 83 26	67 68 072	155° SE	1	1
50 m	1501	033 83 59	150 83 43	2580W		

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1502	502 1503	1504	1505	1506	Average
	Photo	V	V	V	V	V	
Litter Cover (%)		30	15	40	80	40	1P
Bare Ground (%)		65	85	58	20	60	57.6
Cryptogam Cover (%	6)	0	D	0	0	D	0
Rock Cover (%)		0	٥	2	0	D	4.0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determinng PCT and Management Zone (optional)

Morphological Type	4012	Landform Element		Landform Pattern	Microrelief	
Lithology		Soil Surface Texture		Soil Colour	Soil Depth	
Slope	\$64	Aspect	SE	Site Drainage	Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	R = O	
Cultivation (incl. pasture)		1.7	
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R	
Fire damage			
Storm damage			
Weediness	1	R	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B

Page 2 of ____

Date: 25	03 - 2019	Recorders:	ACM, TZ		Plot ID:	19
Project No:	1380 740	Project Name:	Bonshaw S	nlor Form.	Zone ID:	

Plot 20m x 50m: Tree Classes ^a

DBH Class#	Tree Sten	ns Count #	Stems with	Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm				1
30 - 49 cm				/
20 - 29 cm	/			
10 - 19 cm				
5 - 9 cm				
<5 cm ^b			n/a	n/a
ngth of logs (m):	(≥10 cm diameter, >50cm in	1.5;	1; 2 = 45m	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
		81				
/						
			3			

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 25/3/19	Bonshaw	19	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouche
_	Sclerolaena birchii		20	6000		
	Aristida romosa		4	300		
	Aristida romosa Xanthain Spinulosum		0.1	2		ļ <u>.</u>
	Maireana microphylla Einarlia nutany Dichonitra sp. A Bothriochlon nu deipein		0.2	4		
	Einadia nutaris		(0.1	100		
	Dichondra Sp. A	<	101	5		
	Bothriochlon ma deipuis		<u> </u>	100		
	Glycine clandestina Chamae syce drummondii Cheilanthes sieberi		0,5	190		
	Chamae syce drummondii		201)	15		
	Cheilanthes sieberi		20.1	8		
	Roechavic domini		-0.1	1		
	Gomphrena celasoides Pumax drownii Korthamm lanatus unidendified grasses		-0.1	1		
	V Rumax brownii		0.1	1		
	Corrhamys lanatus		0.1			
	unidenditied grasses		5			
	, ,					
-			<u></u>			
						<u> </u>
			ļ			
				1-		
				1		
						,
			I			
			1			

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A Page 1 of ____


Date: 26-03-2019 Recorders: ACM, TZ Plot ID: 20

Project No: 0470861 Project Name: Bonshow Solar Form Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nundewar				
Datum:	94	IBRA sub-region:	Nember	Northern	Comolex		
Likely Veg	etation Class:	Cleard					Confidence H M L
Plant Com	munity Type (PCT):	01	and		EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1507	33 7983	67 67 855	3210 NW	V	V
50 m	1508	33 8003	67 67 809	330° NW		V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1509	190	1511	1512	1513	Average
	Photo	/	. V	V	V	V	
Litter Cover (%)		40	25	50	70	26	31
Bare Ground (%)		60	7.5	SD	80	80	69
Cryptogam Cover (%	6)	0	0	0	D	0	0
Rock Cover (%)		0	0	O	1	0	0.1

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Plain	Landform Pattern	Flood Plain	Microrelief	Deare soils Sparse agreed in
Lithology	Rub arganic Soils	Soil Surface Texture		Soil Colour	Durk brown	Soil Depth	
Slope	Flat	Aspect	ии	Site Drainage	-	Distance to nearest water and type	-

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	R-0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)			
Fire damage			
Storm damage			
Weediness	1	R-0	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B

Page 2 of ____

Date: _	26-03-2019	Recorders:	ALM. TZ	Plot ID:
E 60400				

Project No: 0470861 Project Name: Bonshow Salar Fram Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stems	Count #	Stems wit	h Hollows
DDH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm		/		
50 - 79 cm				
30 - 49 cm	E			
20 - 29 cm				
10 - 19 cm				
5 - 9 cm			1	
<5 cm ^b			n/a	n/a

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
		/				
		/				
		/				
1						
/						

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia, Angophora, Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

. 4

 400 m² plot: Sheet _ of _
 Survey Name
 Plot Identifier
 Recorders

 Date | 26/3 / 19 | // Sonshar
 20 / 72

	10/3/1/ 13°43/40 20	12				
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Aristida Leptopoda		02	13		
	Bothriochloa Decigaria		0.2	13		
	Rumex browning		=0-1	5		
	Rhynchosia minima		20.1	10		
	Marreann microphylle		10.1	\		
	Vantaium Spinesum		-001	L)-		
	Noteland microphylla	<u> </u>	50.1	7		
	Paspaladium distance		-01	1		
	Aristida Leptopoda Bothriochloa Blingment Rimmex brownii: Rhynchosia: minima Maireavia microphylle Yanthium spinosum Notelaed microphylla Paspaladium distans Tribulus: terrestin		20.1	 		
	Convalvatus erabescons		< 9.)	1		
	Carrier and Elegentia		2 2.1	<u> </u>		
	Also Maduma + Bocchora					
	- June					
		<u></u>				-
	· · · · · · · · · · · · · · · · · · ·					
			<u> </u>			
		_				
`						
					·	
	,	·				
				-		
- +						
	A 5 (11-4-)					
	see Growth Form definitions in Annandiv 1					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Black Sil NSO7a gran cover but dear, rut bare soil

Sheet A Page 1 of ____

Date: 26-03-2019 Recorders: ALM, TZ

_Plot ID: ________


Project No: 047084 Project Name: Bonshow Salve Form

Zone ID: _____

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandewar				
Datum:	94	IBRA sub-region:	Nandewar	NacHern Como	x y		
Likely Vege	etation Class:		-	-			Confidence H M L
Plant Com	munity Type (PCT):	594-Lau			EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1514	33 7855	6767 956	3210 NW	V	V
50 m	1520	3 3 7827	6767 997	1440 SE	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2 .	3	4	5	
Attribute	WP	WP 1515 1516 1517 1518	1519	Average			
	Photo /	V		V			
Litter Cover (%)		60	35	50	30	5	36
Bare Ground (%)		30	35	70	50	95	46
Cryptogam Cover (%	6)	20	36	30	30	60	34
Rock Cover (%)		1	1	0	[Ū	0	2.4

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Plain	Landform Pattern	Phin	Microrelief	open weadland /
Lithology		Soil Surface Texture		Soil Colour	Brown	Soil Depth	-
Slope	Level	Aspect	WW	Site Drainage	۵.	Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	2	D	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	2	R-D	
Fire damage			
Storm damage			
Weediness	1	R	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent(<3 yrs); NR = not recent(3 - 10 yr); O = old(>10 yrs)

Sheet B Page 2 of ___

Date: 26 - 03 - 2019 Recorders: ACM, TZ	Plot ID:	P71
---	----------	-----

Project No: 0470861

Project Name: Bonshow Salar Farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree St	ems Count #	Stems with	Hollows
	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm		-		
30 - 49 cm				
20 - 29 cm		1=1		
10 - 19 cm		FI= \\\ \\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\		
5 - 9 cm		6, 2, 1/1 /1 = 20		
<5 cm ^b		4,11, 6:7 2=17	n/a	n/a
gth of logs (m): (≥10 cm diameter, >50cm in	8, 2:1, 1, 14,6; 3	1,4 , 1, =40	

Collition glave

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
					- LO CIII	CIII
	-					1
			-			

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m² plot: Sheet _ Survey Name Plot Identifier Recorders Date 26/3/14 2)

	2 /3/19 DONSNEW L)					
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouch
•	Bothriochloa decipiens		0.5	15		
1	and some of the contract of the second		•			
	Sila corregata Boerhavia dominii		2.0	366		
	Cita coppugata		0.5			
	Boerhanea domini	· · · · · · · · · · · · · · · · · · ·	6.1	80		
	Binacla nultans Partulacca bicolor		(0)	T		
:						ļ
	Partulacca bicolor		201	20		
 -	Opuntia, aurantiaca Dichandra sp. A Cupressus semiglaura shrubs Rumex brownii Pimelia stricta		25.)	2		<u> </u>
	Dichandic sp. A		< ° \	,		
	Cupressus comiadanca was		3			
	Dunex brawni		20.1			
	Pinelia stricta		<0.1	4.		
	(Trees of Nedma		0.1			
	Glycine clandestina Portulacea oberacia			30		
	Tribulus terrestris	_	201			
	the during stressing		201	4		
	Vitadinia Sulcata: Sclerolaina birchii		2.01	ن		
	Scierolaena DWChil		<u> </u>	,		
	Rostellatoria aggendins		101	3		
	1054 (COLING) Adjusted wins		201			
i	- Alinhau ramosa	•	3	7:00		
 -i	Oxalis perentians		(4.)	4		
	Crymbopoyon refractus		<i>₹</i> (> `}	7		
	U G G	*				
	· · · · · · · · · · · · · · · · · · ·					
				L		
				ļ 		
					75	

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF - circle code if 'top 3'. Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ... 100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4×1.4 m, and $1\% = 2.0 \times 2.0$ m, $5\% = 4 \times 5$ m, $25\% = 10 \times 10$ m **Abundance:** 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

to ful grant dar 15 (som Cupyers mon)

gran 10


Sheet A Page 1 of ____

Project No: 0470861 Project Name: Bonshow Jolan Farm Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandavar				
Datum:	94	IBRA sub-region:	Nandowar	Northern	Complex		
Likely Vege	tation Class:		,				Confidence H M L
Plant Com	nunity Type (PCT):	594-Low			EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1522	3 3 8024	6768 238	298° NW	V	
50 m	15 23	337991	67 68277	134 0 SE	1	· ·

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1524 1575	1525	1526	1526 1527		Average
	Photo	V	1	- /	V	/	
Litter Cover (%)		60	70	25	15	7.0	38
Bare Ground (%)		40	30	75	AS.	60	62
Cryptogam Cover (%	6)	10	15	70	20	30	21
Rock Cover (%)		t	٥	0	0	7.0	4

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Crest	Landform Element	1 Flut	Landform Pattern	Low Hill	Microrelief	soil w/ proble
Lithology	5. thy -chayed soils	Soil Surface Texture		Soil Colour	light awaye-	Soil Depth	
Slope	Ven grath	Aspect	NW	Site Drainage		Distance to nearest water and type	-

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	ñ	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	11-0	
Fire damage			
Storm damage			
Weediness			
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B

Page 2 of ____

25000	A	100000000000000000000000000000000000000	A CONTRACTOR OF THE PARTY OF TH	121111111111111111111111111111111111111
Date: _	26-05-2019	Recorders:	ACM, TZ	Plot ID: 22

Project No: 047086/ Project Name: Bonshow Solar farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Sten	ns Count #	Stems with	Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm		y	/	
0 - 79 cm				
0 - 49 cm				
0 - 29 cm		+		
0 - 19 cm		A		
5 - 9 cm		// =3		
<5 cm ^b	/	1// =3	n/a	n/a
th of logs (m):	(≥10 cm diameter, >50cm in	11,2,8,2=23		-

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
						-
	+					
	100					

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 26 / 7/14	Bonshaw	22	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Coursessan serniglanca. Church		hou)	2		
	Rostellularia adscendens!		ا و ک	8		,
	Chamaesya drummondii Glycine clandistine Gida Erimon		201	8		
	abyeine clandistine		5.2	30		
	lian Expluses Pro		101	50		
	DV alike Vallage		1			
	In near ogen grains		1			
	Cymbo pogen refractus		601	3		
	En neapogon gracilis Cymbologan refractus Schrolaine Swichii		201	40		
	Pinelia stricta		201	16		
	Pincha stricta		201	20		
	L'à l'ofise lappulaux		201	1		
	Cátotiso lappulaca Barrhavia domini Portulacca Bicolor		201	12-		
	Portulacca Bicolor		201	50		
	FORMACCA OGRACEM		201	2		
	Heliotropium amplexicaule		201	2		
	Heliotropium amplexicaule Lonvulvulus alsinosius	,.,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	400	<u>S</u>		
	Desmodium varians			3		
	Opundia structa tommera		201	_2.		
		•				
					<u>-</u>	
					-	
						
					,	
					ļ	
				,		
				<u>-</u>		
<u> </u>						
			· .			

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A

Page 1 of

Date: 26 - 03 - 2019 Recorders: ALM, TZ Plot ID: 23


Project No: 0470861 Project Name: Barbhaw Solar Farm

Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandewar				
Datum:	94	IBRA sub-region:	Nandewar	Northern	Complex		
Likely Vege	tation Class:						Confidence H M L
Plant Community Type (PCT):						EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1531	033 8490	67 68 126	159° S	V	1
50 m	1532	033 8504	67 68 075	3310 NW	V	

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1533	1534	1535	1 536	1537	Average
	Photo	1	V	V	1	1	
Litter Cover (%)		90	30	80	30	08	70
Bare Ground (%)		10	70	20	36	20	30
Cryptogam Cover (%	6)	0	0	0	0	0	0
Rock Cover (%)		0	D	٥	.0	Ò	0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	Hill slope	Landform Pattern	Low Hill	Microrelief	graziland
Lithology	~	Soil Surface Texture		Soil Colour	Tight - orange - brown	Soil Depth	-
Slope	Gentle inclined	Aspect		Site Drainage	Direct market	Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R	
Fire damage			
Storm damage			
Weediness		A-0	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B

Page 2 of ____

Date:	26-03-2019	Recorders:	ACM TZ	Plot ID:
		The second secon		

Project No: 0410861 Project Name: Banshaw Salar Farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree St	ems Count #	Stems with	Hollows
JBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm	/	/		
50 - 79 cm	1			
30 - 49 cm				
20 - 29 cm		/ = 1		
.0 - 19 cm		// = 2		
5 - 9 cm		/ = 1		
<5 cm ^b		1 = 1	n/a	n/a
th of logs (m):	(≥10 cm diameter, >50cm	in 7,1,1 =4		

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is \leq 10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
		-				
	2/4					
						-

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

60

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

00 m² plc	ot: Sheet _ of _	Survey Name	Recorders					
Date	26/2/16	Bonshaw	23					
GF 7 Code A	Fop 3 native species in All other native and exot	each growth form group: Ful tic species: Full species nam	ll species name mandatory ne where practicable	N, E or HTE	Cover	Abund	stratum	vouche
	Carrens	alecres phil	smuly s	1	۵۰۷	2		
	pine)a	r shila			<0.1	3		
	Aroskala	yamosa yam refrae di vinyata p			1	A SE		
	Cymbopo	you retructi	AC 1		0.3	40		
	- Side co	errysta p			0.1	100		
	Cassima				10.1	1		
,	Helio A	raipum.			231	28		
	<u> </u>					1.		
	Gener	Juniae Juniae Sp. A. Vin decipsión Hay			201	12		
	' Sble	innie			201	2		
	Dich	andra SpA			107	9		
	Bothrior	Von deeignen	_		1	I		
	(hei) an	then !			(ایل می	7		
	Paraicun	peable of vivos	. :		200	2		
	Beerho	eni !			201	6		
	em i elembi	hable arrows	· A		20			
		, l						
<u> </u>								
		- 						
					· · · · · · ·			
			·					
+-								
	ho Son	<u> </u>				 		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ..., 100, 200, ..., 1000, ...

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

To but very control of the co

19-

Sheet A

Page 1 of ____

Date: 26-03-2014 Recorders: ACM, T2


Plot ID: 24

Project No: 0470861 Project Name: Boushow Salar Fair Zone ID: _____

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandewar			
Datum:	94	IBRA sub-region:	Nondewor	Northern	Camolex	
Likely Vege	tation Class:	9,4055				Confidence H M L
Plant Community Type (PCT):		596 DNG-	Low		EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1538	33 88 00	67 68 481	13005	V	V
50 m	1539	33 88 13	67 68 430	332° NW		

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1540	1541	1542	1593	1594	Average
	Photo	V	0	V	/	1	
Litter Cover (%)		60	80	80	85	60	73
Bare Ground (%)		40	10	20	15	40	15
Cryptogam Cover (%	%)	5	30	5	10	5	11
Rock Cover (%)		٥	81	Ò	0	0	0.2

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Plain	Landform Pattern	Plain	Microrelief	DAN YELDID
Lithology	Sub- clay	Soil Surface Texture		Soil Colour	light away -	Soil Depth	e
Slope	Losel	Aspect		Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	8	
Fire damage			
Storm damage			
Weediness	2	R-0	
Other			

Sheet B Page 2 of ____

Date: 26-03-2014	Recorders:	ACM, TZ	Plot ID:24
Project No: 0470861_	Project Name: _	Bonshaw Solar Farm	Zone ID:

Plot 20m x 50m: Tree Classes a

		Stems with Hollows		
Euc*	Non-Euc^	Euc*	Non-Euc^	
	1	1		
	-			
	((
		n/a	n/a	
	Euc	Euc* Non-Euc^		

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
į.						
						1
	,					
						1

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

40

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m² plot: Sheet _ of _ Survey Name Plot Identifier Recorders

Date 25/3/2019 Bonshaw 24

Date	26/3/2019 Bonshaw 24						
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher	
	Aristrae ramosa Bothiothloa recuprens Eragreshis brownin Cymbopogan velvactus Panieum simile		201	l]
	Bothiochlan ideapren		1.5	500]
	Erapteshin brownii		201	3_			
<u> </u>	Cymbs po cum / Nachos		O . 2	4			
-	Panieum simile		20.1	3			
	TO CHOMACO SPANS		015	200	10	50	\rangle
	Vittadinia sulcata		100	- E-	,	a Win	[7]
	Chem clanda Aina		Co.)	16		Street	3
	Cheilanth sieberi		0-1	30		,	
	Cheilantha sieberi		20.1	5			
<u> </u>	Solchogyne gracitis		201	12			
	Solchogyne gracitis Pomolia stroki unidutificash grasses		201	_3			
	unidustificable grasses		25				
	,						
			•				
							1
						1	1
							1
							۱.
	. , , , , , , , , , , , , , , , , , , ,						1
				!			1
						\	1
							1
							1
)	1
	· ·						1
						-	1
							1
					· · · · · · · · · · · · · · · · · · ·		1
							1
							1
		<u> </u>	<u> </u>		 		1
<u> </u>							1

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

total veg com 35%.

Lead 50%.

Lare 200

Sheet A

Page 1 of ____

Date: 26-03-2019 Recorders: ACM, TZ

Plot ID: 25


Project No: 0470 861 Project Name: Bonshaw Solar Farm

Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandower			
Datum:	94	IBRA sub-region:	Non de war	Nouthern	Compbx	
Likely Vege	tation Class:					Confidence H M L
Plant Comn	nunity Type (PCT):				EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1545	3 3 8687	67 68 607	5270M	V	V
50 m	1531	338635	6768598	67º NE	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1546	1543	1.548	1599	1550	Average
	Photo	V		1	1	V	
Litter Cover (%)		90	70	90	60	60	74
Bare Ground (%)		10	30	10	40	OF	26
Cryptogam Cover (%	%)	6	0	0	0	D	0
Rock Cover (%)	10	0	٥	9	D	9	0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot.

LP = Litter Plot (1m x 1m plot); WP = Waypoint;

gracing padock - Lengt

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Pluin	Landform Pattern	Plain	Microrelief	lond icoup
Lithology	51th clay	Soil Surface Texture		Soil Colour	light orange	Soil Depth	_
Slope	level	Aspect	W	Site Drainage	-	Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	2-0	
Fire damage			
Storm damage			
Weediness			
Other			

Sheet B Page 2 of ____

Date:	26-03-2019	Recorders:	ACM, TZ	Plot ID: 25

Project Name: Bonshaw Solar Farm Zone ID: Project No: __0470861__

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stems	Count #	Stems with	Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm		/		1
50 - 79 cm				1
30 - 49 cm				
20 - 29 cm	1	/	-/-	
10 - 19 cm				
5 - 9 cm				
<5 cm ^b	/		n/a	n/a
th of logs (m): (≥10 cm diameter, >50cm in	8,3 = 11		

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

RAM Attributes: Tree DRH & Hollows (Plot 20m v 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m²	plot: Sheet _ of _	Survey Name	Plot Identifier		Re	ecorders		
Date	26/2/2/19	Bonsham	25					
1	1	<u>-</u>		1		i		
GF Code	All other native and exot	each growth form group: Fuli ic species: Full species nam	e where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Bothwisel	ilva decip	ريت		0.7	200		
	Avishac	lepoto peda			(0-)	3		
	Rauchaus	•			<u> </u>	3		
	Eragrish	~ ^			2011	2.		
	Avastida	Palemon			Q.5	7		
	Wichon	ratemag (1 vi Sp A			o t	200	ı	
	. Milthd	inia sulco	lu-		201	1		
	Meliota	deservations			201	14		
	S de gren	<u> </u>			<u> </u>			
	Glyma :	\\			C 6.1	26		
	Downer	- Januaryan			01	1		
	Cymbe 11 res	on rowartus			0,1	15		
	Chelant	٧,	·		201	2.,		
	<u>Kumus</u>	Liable grows			2 4 1	72		
	um I chank	Liash Groth	3 		25			
		•						
								
	·							
	•							
			·		<u>_</u> _			
								
							<u> </u>	
								
							!	
						·		
		·	·····					
		•						
	<u> </u>							
	<u> </u>		ractivo Eravatia UTE: bia					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and $1\% = 2.0 \times 2.0$ m, $5\% = 4 \times 5$ m, $25\% = 10 \times 10$ m


Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

total page - 25%.

Sheet A Page 1 of ____

Date:	76-03-2019	Recorders	· ACM	5T.		_Plot ID: _	26
	1980£60 :0N		ame: Bonsh	uw Salav	Farm	_Zone ID: _	
Desktop I	nformation Requ	irements for BAM	Plot survey				
Zone:	56	IBRA region:	Non dewar				
Datum:	94	IBRA sub-region:	Nandoway	Northern	Comob x		
Likely Vege	etation Class:						Confidence H M L
Plant Comr	munity Type (PCT):				EEC:	- X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	155 2	3 3 9225	6768 408	ZSSOW	/	V
50 m	1558	33 91 77	67 68 412	870E		V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1553	1554	1555	1556	1557	Average
	Photo	V	V	V	7 3.000		
Litter Cover (%)		25	2.0	15	36	90	26
Bare Ground (%)		75	90	85	70	60	73.8
Cryptogam Cover (%	6)	10	5	0	2	2	3.8
Rock Cover (%)		-6		Ì	0	0	4.0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	Hull slope	Landform Pattern	Low Hill	Microrelief	grass -choud
Lithology	Silti-cky	Soil Surface Texture		Soil Colour	light arange.	Soil Depth	4
Slope	Gentle inchered	Aspect	- W	Site Drainage		Distance to nearest water and type	-

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	6	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal	3	R-0	
Grazing (identify native/stock)			
Fire damage			
Storm damage			
Weediness		D	
Other			

Severity: 0 = no evidence; 1 = light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B Page 2 of ____

Date: 26 - 03 - 17	Recorders:	ACM, TZ		Plot ID: 26
Project No: 0470861	Project Name:	Bonshaw	Solar Fora	Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stem	s Count #	Stems w	ith Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm				
30 - 49 cm				
20 - 29 cm				
10 - 19 cm	M = 5			
5 - 9 cm				
<5 cm ^b	/=1	1 = 2	n/a	n/a
ngth of logs	(m): (≥10 cm diameter, >50cm in	3, 2, 3, 1, 3, 2, 1,	1,3,2=21	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is \$10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
	*					
			p I			

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

 400 m² plot: Sheet _ of _
 Survey Name
 Plot Identifier
 Recorders

 Date | 36/3/2019
 Benshaw
 26

Date	20/5/2019 136NJAWO 26					•
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Compotersion semigranean		051	k f		
	Alinha jerichvensis		0.5	150		
	Enneapage quiti		1.0	7000		
	Convulvin ablasides		£ 12 1	es P		
	Brerhowia dominii		1.0	80		
	Cuporessus semiglausea. Alinhola jerichvensis Exmanyagen maints Convulvum abstracides Barhowia dominii Sida corrugata Procha stricta Cumbopogon refracteus Dichonara sp A		01	300		
	Procedin Stricta		1001	150		
	Cumbopogon refraction		12.1	10		**************************************
	Dichohava SpA		21	G 0		
_	C. Walan I		2 11	90 17 8		
	Hyperican		6.01	8		
	Bathrischen desepren		2.0	1		
	Bodhrischen Descepten Errarostis brown		201	1		
	Chemaesaa drummondii		201	3		
	Chemarsya drummondii Panicum Simila Chloris francara		601	1		
	Chloris transita		2.1	II.		
	Encalypotus dealbort (should)		0.1	ľ		
				•		
			 -			
			-			
			 			
				l I		
						•
			<u></u>			
				<u> </u>		
						<u> </u>
	·					
			<u> </u>			

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...


total cour 5-10% dead 10-25 rock oo 30il 50-75

Sheet A

Page 1 of ____

Date:	12-03-2019	Recorders	E: ACM TZ Plot ID:	27
Project N	138051	_ Project Na	ame: Ronshew Solar form Zone ID:	
Desktop Ir	nformation Requ	irements for BAM	Plot survey	
Zone:	56	IBRA region:	Mandowar	
Datum:	94	IBRA sub-region:	Mandawar Northern Complex	
Likely Vege	tation Class:	Grossland		Confidence H M L
Plant Comn	nunity Type (PCT):		200	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1560	33 83 44	6768 782	18405	V	V
50 m	1561	3 3 83 28	67 68 734	3011	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1562	1563	1564	1565	1566	Average
	Photo	1	V	1	V	V.	
Litter Cover (%)		O.F.	60	.50	70	80	66
Bare Ground (%)		30	240	40	25	7.0	31
Cryptogam Cover (%	6)	5	E	5	. 0	10	5
Rock Cover (%)		1	Ō	0	6	0	0.2

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slepr	Landform Element	Very Low All	Landform Pattern	Low Hill	Microrelief	grossland
Lithology	Clared soils	Soil Surface Texture		Soil Colour	trant orange-	Soil Depth	
Slope	Very godly to	Aspect	- 73	Site Drainage	_	Distance to nearest water and type	7

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R-0	
Fire damage			
Storm damage			
Weediness		R	
Other			

Severity: 0 = no evidence: 1 = light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B Page 2 of ____

_____Plot ID: ____2} Date: 27-03-2019 Recorders: ACM, TZ

Project No: 0410861 Project Name: Bonshow Solar Farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class [#]	Tree Stem	s Count #	Stems wi	th Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm		/	/	/
50 - 79 cm				
30 - 49 cm				
20 - 29 cm				
10 - 19 cm				
5 - 9 cm				1
<5 cm ^b			n/a	n/a
gth of logs (m): (≥10 cm diameter, >50cm in			

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is \$10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

RAM Attributes: Tree DRH & Hollows (Plot 20m v 50m

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
		13				
						-
				-		
						-

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m² plot: Sheet _ of _ Survey Name Plot Identifier Recorders

Date 27 22 23 27

Date	27 27 1					
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Avistian varmosa		(Lo	900		
	Sclerolauna birchii Heliotropium amplexicante		i i	4.0	<u> </u>	
	Sclerolaena brichit		01	60	<u> </u>	
	Heliotropium amplexicante		011	100		
,	Boerhavia dominii		200	3		
	Boerhavia dominii Cheilantha sieberi		50.3	30		
	Einadia nutans Thuncus usifatus Dichondra: repens		<0.1	50		
	Thunians usitatus		200	િ		
	Dichondra: repens		204	P		
	Side Covening.		(0)	ų.		
.[Sham is a		Co.)	40		
	Sida Corrigate Shjam Dichandra sp. A	•	٥٠)	30		
	Erageastis blankii Rumber brownii uni duni fiable grasso /serge Carreidii Ernadia bartata		600)	15		
	Rumbix blownii		20.7	5	-	
_	emigation distributions son		30			
	- Convince					
	Einadia hastata	·	<51	3		
	i.				<u> </u>	
		i				
		•				
	,					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Matrix very coned 50% of la sons 30%

Sheet A

Page 1 of ____

 Date:
 27-63-204
 Recorders:
 ACM, T3
 Plot ID:
 28

 Project No:
 047086
 Project Name:
 Bonshow
 Solar Fark
 Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	dewar			
Datum:	94	IBRA sub-region:	NC			
Likely Veget	tation Class:	DNG				Confidence H M L
Plant Comm	nunity Type (PCT):	516_0N9-No		EEC:	X	Confidence

BAM Nested Plots Requirement Plots LP1 LP3 LP5 midline Im x 1m Grey-crowng Lottler - colling

Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1569	33 86 58	67 68 836	18305	V	V
50 m	1570	3 3 86 43	67 68 787	3 SOLO N	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

N-7-50-11-15	LP	1	2	3	4	5	
Attribute	WP	1571	1572	1573	1574	1575	Average
	Photo	B	~	1	2	1	
Litter Cover (%)		20	50	23	30	46	33
Bare Ground (%)		90	40	75	70	60	65
Cryptogam Cover (%	6)	0	5	0	0	10	, i
Rock Cover (%)			0	D	Ō	0	0.17

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slop	Landform Element	Landform Pattern	Microrelief
Lithology		Soil Surface Texture	Soil Colour	Soil Depth
Slope		Aspect	Site Drainage	Distance to nearest water and type

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	12-0	
Fire damage			
Storm damage			
Weediness			
Other			

Severity: 0 = no evidence; 1 = light; 2 = moderate; 3 = severe -

Age: R = recent (<3 yrs), NR = not recent (3 - 10 yr), O = old (>10 yrs)

Sheet B Page 2 of ____

Date:	27-03-7019	Recorders:	ACM. TZ		Plot ID:	28
Project N	o: CH20861	Project Name:	Bonshaw	Solar Form	Zone ID:	

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stem	s Count #	Stems w	ith Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm				
30 - 49 cm				
20 - 29 cm				
10 - 19 cm				
5 - 9 cm		4		
<5 cm ^b	/		n/a	n/a
th of logs (m)	: (≥10 cm diameter, >50cm in		_	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
						-
						1

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m ² plot: Sheet	_ of	Survey Name	Plot Identifier	Recorders
Date 27/3/	2019	(280	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouche
	Helio Ropinini ampleixicante L'Dichandres Spara		25	5 83		
	Llelio Hopium ampleixicante		·3-5	7200		
	W. Dichandres Sp. An		0.)	7		
	Einadia: nutans Calotio la pulla Scleiola epa birchii		600	1 to 30		
·:	To Caloto lapular		201	اود		
	Scherolaena birchii		७५३	130		
	Marreana microphylla		<u> </u>	20		
			5.0	20		
	Marreana microphylla		₹ <u>81</u>	**************************************		
	William Sand		No. 1			
	Oxalis perennans Rumex blownii		101	3		
	Rumex blownii		4.8.2			
	Backavia domini		10			
	Brechavia adomini		<u> </u>	3		
	and the second s			~		
	<u> </u>					

_						
		-				
	·					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A Page 1 of ____

	Plot ID:	M, TZ	s:A	Recorder	27-03-2019	Date:
e ID:	Zone ID:	v Solar			No: 0470861	
			Nonda	IBRA region:	56	one:
Confidence H M L	olex	ucur Northam		IBRA sub-region:	etation Class:	Datum: likely Veg
Confidence	EEC:			596-DNG-LO	nmunity Type (PCT):	lant Com
Ra		quirement	d Plots R	BAM Nested		
	20m x 50m ✓	LP5		LP3	LP1	1==
J	■ 1m×1m √	midline	Į	P2		
n	1 1 20m x 20n	midline	Į.	 - 		

Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1575	33 90 52	67 68 750	17805	V	-
50 m	1577	33 9047	67 68 699	3470N	1	V

^{*}Magnetig bearing obtained with compas along the midline

	LP	1	2	3	4	5	
Attribute	WP	1578	1519	1280	1591	1285	Average
	Photo			2	V	V	
Litter Cover (%)		80	86	60	65	7.5	7-2
Bare Ground (%)		- 10	20	10	io	25	15
Cryptogam Cover (9	%)	15	10	70	10	5	12
Rock Cover (%)		10		30	25	i	13.4

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	HOU SLOPE	Landform Pattern	Low Hill	Microrelief	grassy
Lithology	I whereday	Soil Surface Texture		Soil Colour	hybr- owner	Soil Depth	-
Slope	very quito slope	Aspect	2470W	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R-6	
Fire damage			*
Storm damage			
Weediness		02	
Other			

Severity: 0 = no evidence: 1= light: 2 = moderate: 3 = severe

Age: R = recent(<3 yrs); NR = not recent(3 - 10 yr), O = old(>10 yrs)

Recorders:

Sheet B

Page 2 of ____

Plot ID: 29

BH Class#	Tree Stems C	Count #	Stems w	ith Hollows
DIT Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
0 - 79 cm				
0 - 49 cm				
0 - 29 cm				
.0 - 19 cm				
5 - 9 cm				
<5 cm ^b	/		n/a	n/a

Date:

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH . (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
				*		
	+					

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 27/3/14		24	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouche
	Ennea pagan gracilis Dichondra sp.A. Rieshavia dominii		۵۰۶ ۳	१७		
	Ennea nogon gracilis		*	005		
	i Dichondra sp.A.		ک، خ	100		
	Resarvaidominii		o 1	7 8		
	Wittadinia/ surleata		(6)	ಀೢ		
	Long to the state of the			11		
	Sida corrugata		ا ي	11.5		
	La Missione - Williams		0-14	0.5		
	Chamaesyce drummondi			ž _{ME}		
	Lumer brownii		19	્ઢ		
	Cheilanthes sieberi Timelta stricta		-2×1	169		
	imelia struka		3.2	15		
,	Cupressus sensiglanca Painicum 5 imile Bothriochloa Idecipums Calaty		201	1		
	Panicum Simile		01	30		
	Bothriochlaa Idei puns		10	1000		
	Calchi		201	15		
	(my Vte W (M) (L) A V (M) a 19MG		<01.	y.	i	
	Fragrostic brownii		201	4	,	
	Gumbopogan retractus		0.7	15		
	Eragrosfis brownii Chymbopogon refractus Chloris truncata		201	ı		
	Encalypting dealbata storub		<i>↑</i> 2	į.		
	imidentifiable pranto sedjer		- marker	* •		~
	, , ,		- , ;.			
						<u> </u>
						·
	· · · · · · · · · · · · · · · · · · ·					
	····	-				
	•					
	···			L		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

John 17 25 %.

Sheet A

Page 1 of ____

Wet - rain

Date:	105-50-42	Recorders	:_ A CM, T Z		Plot ID: _	30
Project N	10: 0470861	_ Project Na	me: Romhaw Sda	r farm	Zone ID: _	
Desktop li	nformation Requ	irements for BAM	Plot survey			
Zone:	56	IBRA region:	Non Lewar			
Datum:	94	IBRA sub-region:	Nandrusay Northern	Complet		
	tation Class		3,000	Core por		Confidence
Likely Vege	idion class.					H M L

BAM Nested Plots Requirement LP1 LP3 LP5 I - 1 20m x 50m I - 1 LP2 LP4 I midline 1m x 1m

Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1583	33 8535	67 69 602	25°NE	V	V
50 m	1584	33 85 65	67 69 043	7110 SW	/	/

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1585 1586	1586	CR21	1888	1589	Average
	Photo	~	V	V	V	V	
Litter Cover (%)		20	40	60	30	75	45
Bare Ground (%)		80	60	40	70	7.5	58
Cryptogam Cover (%	6)	0	0	O	0	6	0
Rock Cover (%)		0	Ō	0	O	5	0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Flat plain	Landform Pattern	Plain	Microrelief	dian parky
Lithology	Silly-chy	Soil Surface Texture		Soil Colour	light crange -	Soil Depth	-
Slope	Level	Aspect	-	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	2	6	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R-6	
Fire damage			
Storm damage			
Weediness		R	
Other			

Recorders:

Sheet B

Page 2 of ____

Plot ID: 30

BH Class#	Tree Stems	Count #	Stems with	Hollows
Bri Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
0 - 79 cm	*			
0 - 49 cm				
0 - 29 cm				
0 - 19 cm	- /			
5 - 9 cm			1	
<5 cm ^b			n/a	n/a

Date: ____

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
	-1					

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 27/5/19	Bunshew	30	

GF	Top 2 positive amoning in cools grounds forms when the Full province are mondated in		<u> </u>			
Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Arishia ramasa		0.5	۵.۵		
	Maireana interplant Maireana interplant Einavia inutans Heliatronoium amplexicaule Side corrugata! Dichondra spe. A Boerhavin dominii Tribulus terrestris		69	2 %		
	School amore beveliet		0.2	100		
	Zinalia l'nutans		Ø 1	90		
	Heliatropium amplexicaule.		0. 2	400		
_ > >_	Carnes	***	0.1	é o		
	Sida corrugata		201	W. 63		
	Dichondra Spe A		2001	20		<u> </u>
	Boerhavin dominii		0.1	40		
	Tribulus terrestris		01	60		
,	VITTAMINA SIMPLATA		201			
	6. Chloris truncata.		201	<i>P</i> 3		
	Convolvations erubescens		€30°1	30		
	Montania gracilis Rumer brownii Bothriochloa decipiens		201			
	Riumer brownii		601			
	Bothriochloa decipiens		02	549		
	•					
						•
	· · · · · · · · · · · · · · · · · · ·					
<u> </u>				!		
		<u> </u>				
		 -				
				'		
						<u> </u>
			·			
					<u>-</u>	
					:	
	<u> </u>					
	<u></u>			<u> </u>		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A

Page 1 of

Date: 27-03-2019 Recorders: ACM, T3


Plot ID: 31

Project No: 1947 0861 Project Name: Bonshow Solar Farm Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	53	IBRA region:	Nondeuriv				
Datum:	94	IBRA sub-region:	Nandowar	Northern	Camala		
Likely Veget	tation Class:						Confidence H M L
Plant Comm	nunity Type (PCT):	596_ Low			EEC:	χ	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1592	338402	67 69 323	19105		
50 m	1593	33 83 80	67 69 277	Noll	V	

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	WP 1594 1595 1596 159	1598	59B 159AA			
	Photo	V		V	V	V	
Litter Cover (%)		70	30	56	10	80	38
Bare Ground (%)		65	50	35	10	10	34
Cryptogam Cover (%	6)	5	10	2	60	-5	16.4
Rock Cover (%)		15	7.0	15	68	10	28

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	Hin Gost	Landform Pattern	Law Hill	Microrelief	boilder trad
Lithology	Silhe-closed sul	Soil Surface Texture	-	Soil Colour		Soil Depth	
Slope	Gentle Indired	Aspect	120° SE	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	2-0	
Fire damage			
Storm damage			
Weediness		R	
Other			

Sheet B Page 2 of ____

Date: 27-03-2019	Recorders:	ACM, TZ		Plot ID: 3
Project No: <u>0470861</u>	Project Name:	Bonshaw	Solar Form	Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class [#]	Tree Ster	ns Count #	Stems w	ith Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm	1		1	
30 - 49 cm				
20 - 29 cm		1		
10 - 19 cm				
5 - 9 cm				
<5 cm ^b			n/a	n/a
gth of logs (m): (≥10 cm diameter, >50cm in	2,		

a - Presence/Absence of trees in each class; b -regeneration is size class < 5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

RAM Attributes: Tree DRH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
Silver-leaved transport	54	1597			L5 cm	

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

400 m² p	lot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date	27/3/10	Bonshaw	31	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Notelaea microcarpa 3-4	m	3	Į į		
	Oblination to mentosalo m		105	1		
	Maironna microphylla Arishing raimosa		0.2	L.		
	Avinisa vamosa		6).5	25		
	Emely poutans		0.1	2000		
	Chamaesner drummondii		5.1	စီဝ		
	Obunta aurantiaca Sida corregata Selevalación biochti		0.1	3	_	
	Side corrupated		0.1	300		
	Selevalaona brichii		20:1	cy o		
	Boerhavia dominii		Lar	₹ ⊘		
	Heliotropinom amplexicante		201	(.SD)		
	Vittadinia sulcata		ا د د	7		
	Portulacca oleracca		< 0.1	2		
	Dichonara sp. A		201	0		
	Solanum		10)	ı		
;	12 Julaten Sierw		401	4.		
	Bodingolin dicipus		िक्			
	Bollineshin dicipus			100		
	Economic Somiclares (cashing)		(2 ¹ ,∞).	6		
	(heilandles sieberi Eureagen gracilis		200	8		
	EMMER goog en gracilis		<i>o</i> 1	20		
	• • •					
	'					
	:					
	· · · · · · · · · · · · · · · · · · ·					
1						
				 -		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF - circle code if 'top 3'. GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A

Page 1 of ____

	D47089	_ Project i	Name:	ponshaw Ja	av Farm	Zone ID:		
							Bubble in the	
Desktop Info	ormation Requ	irements for BA						
Zone:	56	IBRA region:	Nandow	av				
Datum:	94	IBRA sub-region:	N cindo	war Northern	Comerce			
ikely Vegetat			1.0.01	act Toberner	Chiper		Confidence	
		910011 - 0	leaved land				H M L	
Plant Commu	nity Type (PCT):				E	EC: X	Confidence	
		516 - DN9 - N	lad			C: X	H M L	
		BAM Nest	ed Plots Red	quirement				
ot 20m x 50m	- midline	LP3	LP4	de-		20m x 50m 20m x 20m 1m x 1m		
midline	Way Point	Easting	North			oto - Horizontal F	Photo - Vertical	
0 m	1500	33 85 18			1308	V		
50 m	1001	338517	63.60	320 3:	50°N	V	V	
	btained with compas	along the midline				4		
AM Attribute								
AM Attribute (1m x 1m plots)			2		4			
	LP	1	2	3		5		
Attribute	WP	1602	1603	1684	1605	1606	Average	
Attribute			1603	1684	1605	1606		
Attribute	WP Photo	1602	1603 V BS	80	1605	80 1606	76	
Attribute ter Cover (%) re Ground (%)	WP Photo	1602 70 30	1603 V RS : IS	90 10 1691	1605 1/ 165 25	1606 80 70	76	
Attribute ster Cover (%) are Ground (%) yptogam Cove	WP Photo	1602 70 30 0	1603 RS 1S O	90 10 1ear	1605 V 765 25 0	1606 80 70	20	
Attribute tter Cover (%) are Ground (%) ryptogam Cove ock Cover (%)	WP Photo	1602 76 30 0	1603 RS IS 0	90 10 10	1605 1/ 165 25 0	1606 % 80 70 0	76 20 0	
Attribute tter Cover (%) are Ground (%) ryptogam Cove ock Cover (%) ter cover is assesse	WP Photo er (%) ed as the average percular plot); WP = Waypo	1602 TO 30 0 entage ground cover of lint;	RS IS O	AO ID O o five 1m x 1m plots center	1605 765 25 0 0 deed at 5, 15, 25, 35 a	80 70 0	76 20 0	
Attribute tter Cover (%) are Ground (%) ryptogam Cove ock Cover (%) ter cover is assesse = Litter Plot (1m x	WP Photo or (%) ed as the average percular plot); WP =Waypo	1602 TO 30 0 entage ground cover of lint;	RS IS O	90 10 10	1605 765 25 0 0 deed at 5, 15, 25, 35 a	80 70 0	76 20 0	
Attribute tter Cover (%) are Ground (%) yptogam Cove ock Cover (%) ter cover is assesse = Litter Plot (1m x	WP Photo er (%) ed as the average perc 1m plot); WP =Waypo	1602 TO 30 0 entage ground cover of lint; graphy + site features	RS IS O itter recorded from	40 10 0 five 1m x 1m plots center	1605 165 25 0 0 0 0 0 0 0 0 0 0 0 0 0	1606 RO 70 0 and 45m along the midline	7.6 20 0 0 0 e of the 20m x 50m p	
Attribute tter Cover (%) are Ground (% ryptogam Cove ock Cover (%) ter cover is assesse = Litter Plot (1m x orphological Type	WP Photo er (%) ed as the average percular plot); WP =Waypo Physio	entage ground cover of lint; graphy + site features Landform Element	RS IS O itter recorded from	five 1m x 1m plots center	1605 165 25 0 0 0 0 0 0 0 0 0 0 0 0 0	RO 70 0 and 45m along the midline ne (optional) Microrelief	e of the 20m x 50m p	
Attribute Itter Cover (%) Are Ground (%) Typtogam Cover Ock Cover (%) Iter cover is assesses Litter Plot (1m x Prophological Type mology pe	WP Photo er (%) ed as the average percular plot); WP = Waypo Physio Cloyed Solb	entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	RS IS O O itter recorded from that may help in Hall back	five 1m x 1m plots center determining PCT and Landform Pattern Soil Colour	1605 25 0 0 0 0 0 0 0 0 0 0 0 0 0	and 45m along the midline ne (optional) Microrelief Soil Depth Distance to nearest water and type	76 20 0 0 0 e of the 20m x 50m p	
Attribute Inter Cover (%) Inter Ground (%) Inter Ground (%) Inter Cover (%) Inter Cover is assessed Inter Plot (1m x	WP Photo er (%) ed as the average percular plot); WP = Waypo Physio Cloyed Solls Gloyed Solls	entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	RS IS O O itter recorded from that may help in Hall back	five 1m x 1m plots center determining PCT and Landform Pattern Soil Colour	1605 165 25 0 0 0 0 0 0 0 0 0 0 0 0 0	and 45m along the midline ne (optional) Microrelief Soil Depth Distance to nearest water and type	e of the 20m x 50m p	
Attribute Inter Cover (%) Inter Ground (%) Inter Ground (%) Inter Cover (%) Inter Cover is assesses Inter Plot (1m x Inter Plot (1m	WP Photo er (%) ed as the average percular plot); WP = Waypo Physio Cloyed Solls Gloyed Solls	entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	RS IS O O itter recorded from that may help in Hall back	five 1m x 1m plots center determining PCT and Landform Pattern Soil Colour	1605 25 0 0 0 0 0 0 0 0 0 0 0 0 0	and 45m along the midline ne (optional) Microrelief Soil Depth Distance to nearest water and type	20 0 0 0 e of the 20m x 50m p	
Attribute Inter Cover (%) Inter Ground (%) Inter Ground (%) Inter Ground (%) Inter Cover (%) Inter Cover (%) Inter Cover (%) Inter Cover is assesses Inter Plot (1m x WP Photo er (%) ed as the average percurbance The plot); WP = Waypo Physio Cloyed Solls Sturbance	entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	RS IS O O itter recorded from that may help in Hall back	five 1m x 1m plots center determining PCT and Landform Pattern Soil Colour	1605 25 0 0 0 0 0 0 0 0 0 0 0 0 0	and 45m along the midline ne (optional) Microrelief Soil Depth Distance to nearest water and type	3 4055 Y		
Attribute Inter Cover (%) Inter Ground (%) Inter Ground (%) Inter Cover (%) Inter Cover is assessed Inter Plot (1m x Inter Plot (1m	WP Photo er (%) ed as the average percurbance Physio Cloved Solls Sturbance oval	entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	RS IS O Is that may help in Hill back	five 1m x 1m plots center determining PCT and Landform Pattern Soil Colour	1605 25 0 0 0 0 0 0 0 0 0 0 0 0 0	and 45m along the midline ne (optional) Microrelief Soil Depth Distance to nearest water and type	3 4022 A	
Attribute Itter Cover (%) are Ground (%) ryptogam Cove ock Cover (%) Iter cover is assesse = Litter Plot (1m x orphological Type hology	WP Photo er (%) ed as the average percurbance Physio Cloved Solls Sturbance oval	entage ground cover of lint; graphy + site features Landform Element Soil Surface Texture Aspect	RS IS O O itter recorded from that may help in Hall back	five 1m x 1m plots center determining PCT and Landform Pattern Soil Colour	1605 25 0 0 0 0 0 0 0 0 0 0 0 0 0	and 45m along the midline ne (optional) Microrelief Soil Depth Distance to nearest water and type	3 4055 Y	

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Weediness

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B Page 2 of ____

Date: 23-0	03-2019	Recorders:	ACM, TZ			Plot ID: 32
Project No:	2470861	Project Name: _	Bonshaw	Solar	Farm	Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Sten	ns Count #	Stems with	Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				/
50 - 79 cm	*			
30 - 49 cm				
20 - 29 cm	/			
10 - 19 cm				
5 - 9 cm				
<5 cm ^b			n/a	n/a
th of logs (m):	(≥10 cm diameter, >50cm in			

a - Presence/Absence of trees in each class; b -regeneration is size class < 5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

RAM Attributes: Tree DRH & Hollows (Plot 20m v 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
						-
						-
				4		
	100					
	7					
					-	-

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 27/2/19		32	

Date	27/3/19 52					
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Avistida romosa		40	1200		
	Maigreania micre pompla Sida rockugala Dorichmanea which backed		0.1	1		
	Side carrieda		40.1	16		
	Do in chimaner which backer		C01	6		
	Colorcan		200	17		
	Hed to I Vermin		201	17		
	Selvolaena mirehii		201	1		
	Colocer Hedrodren mitteling Selevalaena mitteling Lomandon hint ! Cheilantha		Q.	4-3		
	Cheilan ha		201			
	· /Lunus		201	13	•	
	Rollwiscom deigning		201	EP.		
	Ecitation		2 . \	į		
			- T-fr-	1		
	,					
						
					•	
	· · · · · · · · · · · · · · · · ·		· · · · · · · · · · · · · · · · · · ·			
			,			
-	· · · · · · · · · · · · · · · · · · ·	·				
				-		
<u> </u>	· · ·	·				·
			···			
				<u> </u>		
 						
				-		<u> </u>
-						
	· · · · · · · · · · · · · · · · · · ·			-		
	: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high			F – circle		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0×2.0 m, $5\% = 4 \times 5$ m, $25\% = 10 \times 10$ m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...


Sheet A Page 1 of ____

Plot ID: 33 Date: 27 -03 - 2019 Recorders: ACM. Project No: 047 0861 Project Name: Bombuy Solar Farm Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	26	IBRA region:	Newdowar					
Datum:	94	IBRA sub-region:	Nandowar Nort	noin	Complex			
Likely Vege	etation Class:	Cleared → gr						Confidence H M L
Plant Com	munity Type (PCT):					EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1607	33 83 61	67 69 671	326° NW	~	V
50 m	1613	33 83 58	67 69 7-22	273°W	V	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP L	1608 1609	1610	(61)	1612	Average	
	Photo	1	1	V	-		
Litter Cover (%)		15	10	5	5	2	7.4
Bare Ground (%)		80	65	90	AS	97	83.4
Cryptogam Cover (%	6)	8	0	0	0	Ō	0
Rock Cover (%)		5	2.5	5	10		4.2

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Plantar	Landform Element	Plain	Landform Pattern	Plain	Microrelief	
Lithology	sitty- topsoil	Soil Surface Texture		Soil Colour	fork brown	Soil Depth	
Slope	Leveled	Aspect		Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	2-0	
Fire damage			
Storm damage			
Weediness	1	R	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B

Page 2 of ____

Date: <u>17-03-7019</u>	Recorders:	ACM. TZ	Plot ID:33
Project No: 043 0661	Project Name:	Rada Sala Cara	Zone ID:

Plot 20m x 50m: Tree Classes a

OBH Class [#]	Tree Stems	Count #	Stems with	n Hollows
JBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm		/		
30 - 49 cm				
20 - 29 cm				
10 - 19 cm			/-	
5 - 9 cm				
<5 cm ^b	6		n/a	n/a

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DBH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
						-
	34					
						-
					-	

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

20 64

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

,00 III	plot: Sheet _ of _	Survey Name	Plot Identifier		Re	ecorders		
Date	27/3/79	Bonshaw	33					
GF Code	Top 3 native species in All other native and exo	each growth form group: Full tic species: Full species nam	I species name mandatory e where practicable	N, E or HTE	Cover	Ábund	stratum	vouche
	Car Muni	Ng 100 100 100	\$ (100	18		
	existe an	my percesses en	dear		05	المتاين	disea	r'ecceli
	Heliatrom	un	X - 12 - 224 2		401	65	-1	
	Consevulvu	my perusons ce	سرح		<u>4</u> .	3		
	Chamae Su	Le			۷ (6		
	Alinher Sa	-W 050.			<u>e</u> !.	3 .		
	Baywie chi	ion decipi	<u> </u>		2	3.5		
	New Joseph	was by warning			<u>t</u>	3		
	Schwalaen	non boldering			2	y-		
					, f			
						<u>.</u>		
								•
		_		-				
			-		,			
		<u> </u>	· ·					
		·						
			-,					
	•							
		· · · · · ·						
								
							· · · · ·	_
								•
	· ·							
	·	•	· . · · · · · · · · · · · · · · · · · ·			<u> </u>		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – ctrcle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

futal very court great 0.5 dead 40 rocks 20

Sheet A Page 1 of ____


Plot ID: 34 Date: 27-03-2019 Recorders: ACM, TZ

Project Name: Borshow Solar Farm Zone ID: Project No: _ 04+ 086| ____

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandowar					
Datum:	94	IBRA sub-region:	Nandaure	Nevthern	Comelex			
Likely Ve	getation Class:	Cleaned Land						Confidence H M L
Plant Con	nmunity Type (PCT):					EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1614	33 8569	67 69 788	2750W	V	V
50 m	1620	3 3 85 92	67 69 743	300 NM		

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

114	LP	1	2	3	4	5	
Attribute Litter Cover (%) Bare Ground (%) Cryptogam Cover (%)	WP	1615	1616	(617	1618	1619	Average
	Photo	V	V	~	V	~	
Litter Cover (%)		40	SO	5	25	10	
Bare Ground (%)		60	50	95	75	90	
Cryptogam Cover (9	%)	0	-0	0	0	0	
Rock Cover (%)		0	٥	D		0	

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Plain	Landform Pattern	Plain	Microrelief	Genred Port
Lithology	silly-cloyed seil	Soil Surface Texture		Soil Colour	light rorange	Soil Depth	
Slope	Level	Aspect	-	Site Drainage	н .	Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	O	
Cultivation (incl. pasture)	3	0	
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	O	
Fire damage			
Storm damage			
Weediness			
Other		The second second	

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Project No: _____ Project Name:

Sheet B

Page 2 of

Plot ID: 34

Zone ID:

OBH Class#	Tree Stems	Count #	Stems with Hollows			
2011 01033	Euc*	Non-Euc^	Euc*	Non-Euc^		
80 ⁺ cm				/		
50 - 79 cm						
30 - 49 cm		/				
20 - 29 cm						
10 - 19 cm						
5 - 9 cm						
<5 cm ^b			n/a	n/a		

Recorders:

Date: _____

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

BAM Attributes: Tree DRH & Hollows (Plot 20m x 50m)

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
				11		

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m ²	plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date	27/3/19		34	

	94 31 14			;		
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouch
	Schwichten dirpuis Betwickline dirpuis construction dispuis Action of the deal Metro Worming		0.3	200		
	Bothwichlan dispuis		0~ Z	100		
	cessoric arran lite		<u> </u>	_Q _O		
	doch		25	> కల		
	Mamphron Calosician		0.2	150		
.	Hetao hapmi		0.2	200		
	Conversion all reports	·	0.5	60		
	Dinanci Luck mis 1		04	100		
	Convert valor actions of reality	:	0.2	200		
			0.1	100		
	Nooman Every rostis brownii		200	20		
	to very rostra browni	,	201	2		
	Corrotaines		€0.1	-77		
	·					

				ļ'		
i						
						
	<u> </u>					
	•					
	-A					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Sheet A Page 1 of ____

Date: 27- 03-2019


Recorders: ACM, TZ Plot ID: 35

Project No: 0470861 Project Name: Bonshaw Solow Farm Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandawar				
Datum:	94	IBRA sub-region:	Nandowar	Northern	Complex		
Likely Vegetation Class:		PNG			Į sv		Confidence H M L
Plant Community Type (PCT):		516 - DNG -	Low		EEC:	X	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	14.21	33 86 71	67 69 805	262°W	V	V
50 m	1627	33 86 63	67 69 756	3660N		

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1622	12 1673	1624	1425	1626	Average
	Photo	V	V	V	V	V	
Litter Cover (%)		10	85	70	5		24.2
Bare Ground (%)		90	15	80	95	99	75.8
Cryptogam Cover (%	6)	0	2	0.	6	D	0.4
Rock Cover (%)		Ò	0	D	0	0	0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	flut	Landform Element	Plain	Landform Pattern	Plain	Microrelief	
Lithology	Clayed soil	Soil Surface Texture	-	Soil Colour	light overge-	Soil Depth	
Slope	Leveled	Aspect	_	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	6	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	R-0	
Fire damage			
Storm damage			
Weediness	111	R	
Other			

Severity: 0 = no evidence: 1 = light; 2 = moderate: 3 = severe

Age: R = recent (<3 yrs); NR = not recent (3 - 10 yr); O = old (>10 yrs)

Sheet B Page 2 of _

Date: _	27-03-2014	Recorders:	ACM, TZ	Plot ID:	35
					4

Project Name: Bonshaw Solar Farm Zone ID: Project No: 0470861

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stems C	Count #	Stems with Hollows		
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^	
80 ⁺ cm		/			
50 - 79 cm					
30 - 49 cm					
20 - 29 cm	/				
10 - 19 cm					
5 - 9 cm					
<5 cm ^b	/		n/a	n/a	
gth of logs (m	1): (≥10 cm diameter, >50cm in			+	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
						-
						1
			4			
		1				
	-					

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, Corymbia , Angophora , Lophostemon and Syncarpia)

[^] Non-Euc - Record of living native non eucalypt trees

 400 m² plot: Sheet _ of _
 Survey Name
 Plot Identifier
 Recorders

 Date 21/3/19
 Bonshaw
 35

Date	21/3/14 (5000) SS					
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Schero lamon box chet		7.0	150		
	Partulaces pleracen		201	18		
	Emalin mutons	3.33	Roll	10		
	Marrana microphylla	3 33		18		
	Bothrivehlen dur min	**************************************	<i>○</i>	200		
	Runex		201	(3)		
	Convulvation perceptoral		€0.1	\$		
	Side corrugale		2001 2001	2.9		
	Ennember 9		0.5	Za.		
	Chlorin trumenta		Co.)	2		
	Avistide varmosa		3	29		
	I amanta his ili			2.5		
	All other native and exotic species: Full species name where practicable Solera lasar by drift Perstalases blarases Line Am mators Ala is carra microphylla Bothribella decipaini Rumex Convalación evabraces Line paga Line paga Chlorin trumenta Avisticas a amosa Lomandon friedla gerrani Tili con identificada gerrani Tili direct		14			
	Mach		2.5			
	Evagate 12		200	3		
			<u></u>			
· · · · · · · · · · · · · · · · · · ·	,		·			
	-					
					<u></u>	
	·			1		
					·	
				· · · · · · · · · · · · · · · · · · ·		
						
1						
						
		· · · · · · · · · · · · · · · · · · ·	· ·			
	·				•	
	·					
				,		
	·					

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

For 30

Sheet A Page 1 of ____

Date: 28-03-2019

Recorders: ACM, TZ

Plot ID: 36

Project No: 0470 861

Project Name: Bonshow Solar Form

Zone ID: _

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandowar				
Datum:	94	IBRA sub-region:	Nemdowar	Northern	Canales		
Likely Veg	etation Class:						Confidence H M L
Plant Com	munity Type (PCT):					EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1659	33 9043	6768 195	3150 W	V	~
50 m	1630	3 3 90 15	67 68 237	14708E	~	

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1431 1432	1632	1633	1634	1635	Average
	Photo	-	~	1	~	V	
Litter Cover (%)		65	6 %	30	25	45	46.6
Bare Ground (%)		45	30	70	75	50	54
Cryptogam Cover (%	6)	15	0	5	0	S	5
Rock Cover (%)		٥	7	λ	- O	35	1.4

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Slope	Landform Element	14:11512	Landform Pattern	Low Ail	Microrelief	grass over
Lithology	Clayed sails	Soil Surface Texture	4	Soil Colour	light arouge brown	Soil Depth	
Slope	Gently inclined	Aspect	W	Site Drainage	11 500 10	Distance to nearest water and type	5 tom 5

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3.	R-0	
Fire damage			
Storm damage			
Weediness		D_	
Other			

Sheet B

Page 2 of ____

Date: 28 -03 - 2019	Recorders:	ACM, TZ	Plot ID: 36

Project No: 0470861 Project Name: Bonshow Solar Farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stem	ns Count #	Stems with	Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				/
50 - 79 cm				
30 - 49 cm				
20 - 29 cm	/		/	
10 - 19 cm				
5 - 9 cm				
<5 cm ^b			n/a	n/a
gth of logs (m):	(≥10 cm diameter, >50cm in	2,7,1,1,9,2	= 22	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Count apply when the number of tree stems within a size class is ≤ 10 . Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
						1
						1
	-					

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 28/3/10	Bonshow	36	

GF ode	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Bethrje chlou deingræns Aristide laumosa +? Enneapogon gracilis Sida Corrugada Cymbopogon refraetus Dichondra spid Rumey brownii Clym		1.5	600		
	Aristide recomosa +?		0.2	15		<u> </u>
	Enneapogon gracilis		4.0	600		
	Sida Corruyada		0.1	90		
	Cymbopogon refractus		01	30		
	Dichondra sp. A		0.1	100		
	Rumey brownii		201	ď		
	Ayen 100		Coil	90		
·	The state of the s		.	_		
	Erigrostia branini		01	24		
	Panicum smile !!		0.2	50		
	Cupromy glancophyna Boerhavia domini Meliofrepium amplexicane Prinelia girista Limiduntificile grams		401	4		
	Boerhavia dominii		20.1	14		
	Melintrepium ampleximus		20.1	¥		
	Pinelia With		601	1		
	1. I delieste granes		7	1		
	- (m) autority		20.)	,		
	. 1			 '-		
				 		
		-				
		-				
J.				 		
				-		
-				ļ		
				 	<u> </u>	
			 	ļ	<u> </u>	
	•					
			I	-		[

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

to a Vey 10 dead 40 roch 20 have 20

Sheet A

Page 1 of ____

Date: 28-03-2019

Recorders: ACM, T7

_____ Plot ID: _____ 37

Project No: 0470861

Project Name: Bonshaw Salar Farm

Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandewar			
Datum:	94	IBRA sub-region:	Nondoway	Northern	Comoker	
Likely Veg	etation Class:					Confidence H M
Plant Com	munity Type (PCT):				EEC:	Confidence H M

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1 636	33 85 64	545 83 73	17705		
50 m	1637	338579	61 68 321	327°NW	V	

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1638	1639	1640	16.41	1642	Average
	Photo	V	V	V	V	V	
Litter Cover (%)		70	90	80	60	60	72
Bare Ground (%)		30	1.0	20	40	40	1.8
Cryptogam Cover (%	%)	O	0	0	0.	0	0
Rock Cover (%)		0	0	0	D	0	0

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	= Flat	Landform Element	Place	Landform Pattern	Main	Microrelief	gracey doub
Lithology	Hoved soils	Soil Surface Texture	-	Soil Colour	Isolal micros	Soil Depth	
Slope	Very gentle intro	Aspect	WW	Site Drainage	NO 1 - 2 6-	Distance to nearest water and type	a som W

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	stones of felled trens resent
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	2-0	
Fire damage			
Storm damage			
Weediness		η	
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Recorders: ACM, TZ

Sheet B

n/a

16, 7, 1, 1, 7, 4, 11, 7, 1, 1, 9, 4, =64

Page 2 of

_Plot ID: _______

n/a

Plot 20m x 50m:	Tree Classes a			
DBH Class#		ms Count #	Stems w	ith Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm	×			
50 - 79 cm	-			
30 - 49 cm		*		
20 - 29 cm				
10 - 19 cm		1		
		1		

- Presence/Absence of trees in each class;	b -regeneration is size class <5 cm;	# Living trees only;
--	--------------------------------------	----------------------

Length of logs (m): (≥10 cm diameter, >50cm in

5 - 9 cm

<5 cm ^b

length)

Date: 28-03-2019

Count apply when the number of tree stems within a size class is ≤ 10 . Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
					1	
						-

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 28-/3/14	Bonshan	37	

	20 19/14					
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Sida corrugata !!		_0.1	50		
	Gyen () Gyneaposon gracilis Gynbapogon Vefractus Panicum simile		60.1	20		
	Guneapagan wantis	4	0.1	4		
	Cymbopogen repractus		0.)	20		
	Panicum simile		0)	11		
	136thris Chlaa (deipnin)		<i>"</i> 3	700		
	Excavasta browning		0.)	j1		
	Chailantles sieberi		C 6 ·]	3		
	Heliotropium ampliricaule		20.1	3		
	Dichindra sp. A 1		26.1			
	136+mir Chlar deriprins Ercyristin brownin Chailanthes sieber: Heliotropium amphricaule Dichindra sp. A. Convulvuluy alsinoides		600	7		
	Rumey brownii		60.5	8		
	1,		20.1	8		
,	Chamaesyce chrummondi					
	John noi fatus Solenogyne bellioides un i dentifiable promon Aristid- roman Oxak perennant Pimelio stilla		201	5 B		
	Solenogune bellioides		40)	7		
	in dentifiable grances		5	7		
ļ	Act, lide Vaccou		1	20	· ·	<u> </u>
	1) X h V ne Ye a name		201	1		
	Primalia stilla		4011	,		
	1 Time Im		<u> </u>	<u> </u>		
		,				
	,					
ļ						
	·					
<u></u>			ļ			
•			 			
<u> </u>			!			
			 			
	·					<u> </u>
 			 -			
				-	<u> </u>	
						<u> </u>

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4×1.4 m, and $1\% = 2.0 \times 2.0$ m, $5\% = 4 \times 5$ m, $25\% = 10 \times 10$ m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

dend incl trunk go rech oil 30

Sheet A

Page 1 of _

Date: 28 - 03 - 2019

Recorders: ACM, T2

Plot ID:

Project No: 0470 861

Project Name: Bonshaw Solar Farm

Zone ID: 36

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandruar				
Datum:	94	IBRA sub-region:	Nandeway	Northern	Comelex		
Likely Veg	getation Class:	had been no					Confidence
Plant Con	nmunity Type (PCT):	516 mod-bus			EEC:	×	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1643	33 8685	61 68 653	235°5W	V	V
50 m	1644	33 8640	61 68 645	80°€	V	/

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1645	1646	1647	1648	1649	Average
	Photo	-	V		V	V	
Litter Cover (%)		98	95	60	45	10	63.6
Bare Ground (%)		ı	5	40	77	80	36.4
Cryptogam Cover (%	6)	0	0	D	0	6	0
Rock Cover (%)		0	Ð	D	Ò	0	٥

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flok	Landform Element	Phin	Landform Pattern	Plain	Microrelief	would
Lithology	Sant of A	Soil Surface Texture		Soil Colour	dock bloken	Soil Depth	
Slope	Levelod	Aspect	_	Site Drainage	-	Distance to nearest water and type	- 30m V

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	stamps (with most
Cultivation (incl. pasture)			
Soil Erosion		1	
Firewood CWD removal			
Grazing (identify native/stock)	3	R	Cattle of sheep
Fire damage			
Storm damage			
Weediness		17	
Other			

Severity: 0 = no evidence; 1 = light; 2 = moderate; 3 = severe

Sheet B

Page 2 of ____

Date: 28-03-2019	Recorders:	ACM, TZ	Plot ID: 38
Project No: <u>0470861</u>	Project Name: _	Bonshaw Solar Farm	Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stems Co	ount#	Stems w	ith Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm				
30 - 49 cm	XI = 7			
20 - 29 cm	MX **** M			
10 - 19 cm	M(M) = 11			
5 - 9 cm	// = 2			
<5 cm ^b			n/a	n/a

- Presence/Absence of trees in each class:	h -regeneration is size class <5 cm:	# Living trees only:

Count apply when the number of tree stems within a size class is \leq 10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

f. albons

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
						+
						+
						-
						+

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 28/3/14	Bonsham	38	

	00) 114 90 90					
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
	Scleralaena birchii		601	12		
	Aristian ramosa		20.1	20		
	· Suealorptors alberrs		50%		7	
	Maileann microphylla		40.1	7		
	un identifiable orvaries		مورن			
	Ominhia aurantiaca		20.1	3		!
	Pinelia : stricta		20.1	1		
	Example blauni		(6)	2		
	Roshithatoria adscendens		<0.1	3		
	Aristian tamosa Encologotos albens Mai Icana microphylla miduli fiable granes Opuntia aurantiaca Pinelia stricta Evarostia brownis Roslitularia adscendens Tustic stripa scasta		40.1	4		
		· · · · · · · · · · · · · · · · · · ·		17		
			•			
						
	-		:			

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

dead / lither \$5 to 5% E. albens. !.

93000 y S

Sheet A

Page 1 of ____

Date: 28 -03 - 2019 Recorders: ACM, TZ Plot ID: ____

Project No: 0470861 Project Name: Ronshaw Solar Farm Zone ID: 39

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandouar				
Datum:	dt.	IBRA sub-region:	Nandouar	Naithern	Comobe		
Likely Vege	tation Class:						Confidence
Plant Com	munity Type (PCT):				EEC	:	Confidence

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1650	33 86 76	67 68 698	SCO NE	V	V
50 m	1656	6 648	67 68 719	225° 5W	V	V

^{*}Magnetig bearing obtained with compas along the midline _ 338+70

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1651	1652	1.653	1654	1655	Average
	Photo	1	V	V		V	
Litter Cover (%)		95	98	86	55	89	
Bare Ground (%)		5	1	15	40	7.0	
Cryptogam Cover (9	%)	0	D	0	7)	۵	
Rock Cover (%)		D	D	<	5	0	

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Flat	Landform Element	Plain	Landform Pattern	Plain	Microrelief	
Lithology	Clared soll	Soil Surface Texture	-	Soil Colour	Tight Blangs	Soil Depth	-
Slope	Loveled	Aspect	÷	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0.	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	2	R-O	
Fire damage			
Storm damage			
Weediness			
Other			

Seventy: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Sheet B

Page 2 of

Date: 70	-03 -2019	Recorders:	ACM T	7	Plot ID:
Date	103 60	Recorders.	MCM,	t	PIOUD;

Project No: _0470861_

Project Name: Bonshaw Solar Farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree Stems	s Count #	Stems with	Hollows
DOTT CIUSS	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				/
50 - 79 cm				
30 - 49 cm				
20 - 29 cm	/			
10 - 19 cm				
5 - 9 cm			/	
<5 cm ^b			n/a	n/a
th of logs (m): (≥10 cm diameter, >50cm in			

- Presence/Absence of trees in each class;	b -regeneration is size class <5 cm;	# Living trees only;
--	--------------------------------------	----------------------

day a small section of 210 cmp

Count apply when the number of tree stems within a size class is ≤10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus, ^ Non-Euc - Record of living native non eucalypt trees

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

400 m² p	olot: Sheet _ of _	Survey Name	Plot Identifier		R	ecorders		
Date	28/3/19		39					
GF Code		ach growth form group: Fui c species: Full species nam		N, E or HTE	Cover	Abund	stratum	voucher
	Arisfiaa ro	mugh	····		1-5	100		
	Side corr	birinii e bellicides a sp. A; marans	110		0.1	200		
	Scherolarya	birinii			0-3	200		
	Solenogun	e bellivides			201	40		
	Dichoran	a sp. A;	:		0.2	90		
	Eincolia	nutans			2011	⁵ 6		
	Marreans	microphyllo coprosser/ser a deispins uranfiaca 3cobra	1.		0.1	1		
	un i donti fio.	(oprosses / sea	syes :		5			
	Bothrio chlo	a decipin	·		0.2	200		
	Opunhi a	uranfiaca		,	6001	Z.		
	Austro Sipa	3 cabra			0.1	7		
	, 	·						
						ļ		
					ļ	 	 	
						ļ	 	
								
	· -							
						 		
			·			ļ		
						ļ		
					ļ	ļ		
	•			•	[,
								
					ļ	ļ		
						ļ		
						ļ		
	· · · · · · · · · · · · · · · · · · ·				-			
			·		 	 		
							 	
							-	
					<u> </u>	 	<u> </u>	
							 	
	e: see Growth Form definiti		l: native, E: exotic, HTE: hig			F - circle		

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover): Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

Total Voy 5 Dear 75 bare 20

Sheet A

Page 1 of ____


Date	70 02 2-10	Dagardana	Nova To	Plot ID: 40
Date:	28-03-2019	Recorders:	ACM, TZ	Plot ID:

Project No: 0470861 Project Name: Brokaw John Farm Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandewar			
Datum:	94	IBRA sub-region:	Nandowar	Northern	Comolex	
Likely Veg	etation Class:					Confidence H M L
Plant Com	munity Type (PCT):				EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1857	33 9305	67 68 405	40° NE	V	
50 m	1663	33 9350	67 68 429	20802M	V .	V

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	NP 1658 1659 1660	1961	1662	Average		
	Photo	V	V	V	V	V	
Litter Cover (%)		30	30	75	10	40	29
Bare Ground (%)		55	68	70	70	04	64
Cryptogam Cover (%	6)	0	0	0	0	0	0
Rock Cover (%)		5	5	<	10	Ò	€5

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Rolling Court	Landform Element	Halalage	Landform Pattern	14211	Microrelief	rach told
Lithology	Silly-clay	Soil Surface Texture		Soil Colour	light ovarge	Soil Depth	
Slope	Stop	Aspect	315° NW	Site Drainage	decensions	Distance to nearest water and type	Siso in N WW

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	RO	
Fire damage			
Storm damage			
Weediness			
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Sheet B Page 2 of ___

Date: _ 28-03-2019	Recorders:	ACM, TZ	Plot ID: 40
Project No: <u>0470861</u>	Project Name: _	Bonshaw Solar Form	Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class [#]	Tree Ster	ns Count #	Stems w	ith Hollows
DDN Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm	1/1		1	
30 - 49 cm	//		1	
20 - 29 cm				
10 - 19 cm				
5 - 9 cm				
<5 cm ^b			n/a	n/a
ngth of logs (m): (≥10 cm diameter, >50cm in	14,4,3,2,=7	3	4

a - Presence/Absence	of trees in each class:	b -regeneration is size class <5 cm:	# Living trees only:

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
Silver looked lembers	56	1664	3 3 93 35	67 68 429	1 10 and	
ll .	45	1665	33 93 14	67 68 405	Llocong	
	+					
						4

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 28 /3/14	Bonshew	40	

GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	vouche
	Cymbopogen refractus Enneapogen graniis Kristiau ramosa Opuntia aurantium Aristiau lystopoan Sida Corrugata IIII Dichondra Sp A Rostellularia		0.1	10	1	
	Enneapogan granitis		0.2	50		
	Klistika ramosa		0.5	100		
	opunta aurantian		0.5	30		
	Aristina hystopona		40.1	5		
	Side Corrugate 1		0.1	100		
	Dichondra Gn A		0.)	40		
	Rostellularia		201	3		
	Gyern II Chamae syce drumonini Porrsonsin emalyptifolia Bothvio, hlow decipeurs I Hyperium I rmelia glicta		201	20		
	Chamae syce drumondii		201	3		
	Porrsonsin encaliptifalia		2001	400		
	Bothrio, hlow decipions		Ø ·)	50		
	Huperian		601	8		
	Princlin dicta		40.1	4		
	Encotypin melanaphica		25%		1	
	Encotyphis milanophbia Ozothamnus cassinicides		601	1	-	
		7				
	Melichius arceolatus ' imiliantifiaste grames	,	20.1	1		
	Insight dead licely accorded		5	'		
	July Company of Marie					
				<u>-</u>		
						
						
	juil outside plat: 1/morchia Also Notelosa micro				'	
	Also Noteloca mirro					
						
	·					
						

GF Code: see Growth Form definitions in Appendix 1 N: native, E: exotic, HTE: high threat exotic GF – circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover): Note: 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m

Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

9/L Tesul og 60 Soil 32

Sheet A

Page 1 of

Date: 28-03-2019

Recorders: ACM.

Plot ID:

Project No: 0470861

Project Name: Bonshaw Solar form

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandawar			
Datum:	94	IBRA sub-region:	Mardemar	Narthern	Camplex	
Likely Ve	getation Class:				33.1(33)	Confidence H M L
Plant Con	nmunity Type (PCT):				EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1667	33 83 92	67 68 534	2590 W	V	~
50 m	1673	33 83 41	67 68 541	82° £	~	

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	2	3	4	5	
Attribute	WP	1668	1669	1670	1521	1672	Average
	Photo	V	1	V	V	~	
Litter Cover (%)		95	10	5	1	S	23.2
Bare Ground (%)		5	90	5	99	95	58.8
Cryptogam Cover (%	6)	0	0	60	()	0	12
Rock Cover (%)		۵	٥	90	D	0	18

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. lots of shape of cours south LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	Crest	Landform Element	Hill eyest	Landform Pattern	14:11 = 10.4	Microrelief	ración boold
Lithology	Clevery soils	Soil Surface Texture	4	Soil Colour	legt overge-	Soil Depth	-
Slope	Very yearly	Aspect	340° N	Site Drainage	-	Distance to nearest water and type	-

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	0	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	n0	
Fire damage	~		
Storm damage			
Weediness			
Other			

Severity: 0 = no evidence; 1= light; 2 = moderate; 3 = severe

Sheet B Page 2 of _

Date:	28 -03 -2019	Recorders:	ACM, TZ	Plot ID:2	11
					1

Project No: 0470861

Project Name: Bonshaw Solar Farm Zone ID:

Plot 20m x 50m: Tree Classes a

DBH Class#	Tree	Stems Count #	Stems wi	th Hollows
DBH Class	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm				
50 - 79 cm	1		1	
30 - 49 cm				
20 - 29 cm		W1		
10 - 19 cm	1	1		
5 - 9 cm		1		
<5 cm ^b			n/a	n/a
gth of logs	(m): (≥10 cm diameter, >50c	min $3, 2, = 5$		

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

Cultibres description

^ Non-Euc - Record of living native non eucalypt trees

Count apply when the number of tree stems within a size class is <10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20
Slag - Euc	53	PF31	338350	67 68246		2

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

	plot: Sheet of Survey Name Plot Identifier		Re	ecorders		
Date	28/3/14 Bonshaw 41					
GF Code	All other native and exotic species: Full species name where practicable	, E or HTE	Cover	Abund	stratum	vouch
	Coprarus geniglama 5-7 m par		10	73	T	
	Austro Aipa Scapra		01	40		
	Maircana microphylla 11		0.3	7		
	Copressus semiglance 5-7m put Any boot pa Scabra 1 Maircan prierophylla 11 Scherolaena birchii 11/1/4/11		0.3	100		
	Sida corrugata Meilansher Sieberi		40.1	50		
-	Meilanther Sieberi		1001	20		
	Glycen Oxaris per eman Ginalia predans		201	8		
	8 xarin per eman		601	15		
	Binalia mutans		601	15		
	Gamarrena		40.1	2		
	Gamajorena Blechavia demini Opundia anvantana Gynoden dactylon 1. Arichida romosa		401	30		
	Opendir auvanhan-		201	3		
	Gynodon dactinglan		20			1
	Arichda romósa		0.4	30		
						1
						<u></u>
						<u> </u>
				!		
·						
	· ·					
	31					
	:					
						
	I ad a change + con change. They like		1			
	E conversate here					
	Lots of sheep + con chang. They like to congregate here Aggreen to have been overseum with			 		
	Couch					
	Lew.vo					

GF Code: see Growth Form definitions in Appendix 1 **N:** native, **E:** exotic, HTE: high threat exotic **GF - circle code** if 'top 3'. **Cover:** 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); **Note:** 0.1% cover represents an area of approximately 63 x 63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m **Abundance:** 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

g/c very 200 deand/manure 40 tech 25% to 15

Sheet A

Page 1 of ____

Date: 28-03-2019

Recorders: ACM, TZ

Plot ID: 42

Project No: 0470861

Project Name: Bonshaw Solar Form

Zone ID:

Desktop Information Requirements for BAM Plot survey

Zone:	56	IBRA region:	Nandewar		
Datum:	94	IBRA sub-region:	Nandeway Northe	m Comolex	
Likely Veg	etation Class:				Confidence H M L
Plant Com	munity Type (PCT):			EEC:	Confidence H M L

BAM Nested Plots Requirement



Plot 20m x 50m - midline

Point on midline	Way Point	Easting	Northing	Orientation*	Photo - Horizontal	Photo - Vertical
0 m	1675	33 85 33	67 68 664	18°N	V	V
50 m	(681	33 85 34	67 68 715	16405		

^{*}Magnetig bearing obtained with compas along the midline

BAM Attribute (1m x 1m plots)

	LP	1	1 2 3 4		4	5		
Bare Ground (%) Cryptogam Cover (%	WP	1676	1437	1678	1679	1680	Average	
	Photo	ν	1	V	V	V		
Litter Cover (%)		40	30	5	90	60	55	
Bare Ground (%)		10	10	95	10	40	45	
Cryptogam Cover (%)		0	0	6	0	0	0	
Rock Cover (%)		ō	0	0	9	0	O	

Litter cover is assessed as the average percentage ground cover of litter recorded from five 1m x 1m plots centered at 5, 15, 25, 35 and 45m along the midline of the 20m x 50m plot. LP = Litter Plot (1m x 1m plot); WP = Waypoint;

Physiography + site features that may help in determining PCT and Management Zone (optional)

Morphological Type	±+14+	Landform Element	Plain	Landform Pattern	Plan	Microrelief	Grasts
Lithology	Clayer soils	Soil Surface Texture	100	Soil Colour	orange bean	Soil Depth	
Slope	Very Gentrical		N	Site Drainage		Distance to nearest water and type	

Plot Disturbance	Severity Code	Age Code	Observational Evidence
Clearing (incl. logging)	3	Ŏ.	
Cultivation (incl. pasture)			
Soil Erosion			
Firewood CWD removal			
Grazing (identify native/stock)	3	12-13	
Fire damage			
Storm damage			
Weediness			
Other			

Seventy: 0 = no evidence, 1= light; 2 = moderate; 3 = severe

Sheet B Page 2 of ____

Date: <u>28-03</u>	- 2019 Rec	orders: Acm	Plot ID:	
Project No: <u>0</u> 4	13861 Pro	ject Name: <u>Bo</u>	Zone ID:	
	: Tree Classes a	tems Count #		Stems with Hollows
DBH Class#	Euc*	Non-Euc^	Euc*	Non-Euc^
80 ⁺ cm	1		1	
50 - 79 cm	\$			

10 - 19 cm 5 - 9 cm <5 cm ^b n/a n/a Length of logs (m): (≥10 cm diameter, >50cm in

30 - 49 cm

20 - 29 cm

Count apply when the number of tree stems within a size class is \leq 10. Estimates can be used when >10 (eg 10, 200, 30, ..., 100, 200, 300). For a multi-stemmed tree, only the largest living stem is included in the count/estimate. Tree Stems must be living.

Species Name	DBH (cm)	WP	Easting	Northing	Hollow <20 cm	Hollow >20 cm
Silver-Loursed Ivon	18	1682	3385 39	67 68 671	3x 410.2	

a - Presence/Absence of trees in each class; b -regeneration is size class <5 cm; # Living trees only;

^{*} Euc - Record of living native eucalypt trees (includes all species of Eucalyptus,

[^] Non-Euc - Record of living native non eucalypt trees

36

-This document has not been endorsed or approved by Office of Environment and Heritage or Muddy Boots Environmental Training-

	ten.		
400 m ² plot: Sheet _ of _	Survey Name	Plot Identifier	Recorders
Date 28/3//9	Bunshaw	42	

Date	20/5/19 13083120 42	<u> </u>	•			
GF Code	Top 3 native species in each growth form group: Full species name mandatory All other native and exotic species: Full species name where practicable	N, E or HTE	Cover	Abund	stratum	voucher
<u> </u>	Cylina 1"		20-1	30		
	Glyan Stan corrugator Einadus naturas Euralypha melanophlora Jell Meireana microphyla		01	40		
	Ejnami notans		2011	20		
	theolyphy me anophlace file		20	1		
	Marcana microphyla		61	2		
	Aristide Hamisa		3	300		
	Heliotropis amplerocoula		(g.)	13		
	Dichondra Sp A Piniella strictà Rostellularia adscendens		< o 1	3		
	Dinalia «triolà		CO.1	1		
	Rostellularia adsundans	-	40.1	15		
			->			
	Boerhavia dominii		Co.1	15		
	Schwalaena birchii Tribulus Terrentini Portulaena oleracea Vittadinia sulcuisa		1.01	2		
	Sileralaena birchii		0.)	40		
	Tribulus dervatris		1.01	60		
	Partulara deracea		201	3		
	Vitalinia coloria		20.1			
	unidentifiable grusser		4	, A		
	uniones (free process		 7			
					 -	
						
	· · · · · · · · · · · · · · · · · · ·					
	· · · · · · · · · · · · · · · · · · ·					
		·				
	· · · · · · · · · · · · · · · · · · ·					

GF Code: see Growth Form definitions in Appendix 1

N: native, E: exotic, HTE: high threat exotic

GF - circle code if 'top 3'.

Cover: 0.1, 0.2, 0.3, ..., 1, 2, 3, ..., 10, 15, 20, 25, ...100% (foliage cover); Note: 0.1% cover represents an area of approximately 63×63 cm or a circle about 71 cm across, 0.5% cover represents an area of approximately 1.4 x 1.4 m, and 1% = 2.0 x 2.0 m, 5% = 4 x 5 m, 25% = 10 x 10 m Abundance: 1, 2, 3, ..., 10, 20, 30, ... 100, 200, ..., 1000, ...

g/ total vy 7 dean \$3 rock or

+ treeton's 0.2%

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

BAM Plot Data

 www.erm.com
 Version: 2.0
 Project No.: 0470861
 Client: GAIA Australia
 18 May 2020
 Page G3

							Compo	sition Structure					Function														
						Nati	ive Richness	(count no	spp)		Cov	er (Sum co	ver of spp ir	each grow	vth form gro	oup)	Large Tree (P/A)	НВТ	Litter Cover	Logs		Tree DBH (T	ree stems - ı	native species	s only) (P/A)		HTE
Plot	Zone	Easting	Northing	Bearing	Trees	Shrubs	Grasses	Forbs	Ferns	Other	Trees	Shrubs	Grasses	Forbs	Ferns	Other	>80 cm DBH	(P/A)	(average)	(m)	5-9cm DBH	10-19cm DBH	20-29cm DBH	30-49cm DBH	50-79cm DBH	<5cm DBH	(number)
1	56	337831	6767626	86 E	2	1	1	4	0	0	45.0	0.1	0.2	0.5	0.0	0.0	0	1	42	21.86	1	1	1	1	1	1	1
2	56	338406	6767613	344 N	6	3	3	2	0	3	80.5	0.3	0.3	0.2	0.0	0.3	1	1	75	49.7	1	1	1	0	1	1	1
3	56	338667	6767581	24 NE	3	3	0	2	1	0	45.2	0.4	0.0	0.2	0.1	0.0	0	1	43	37.1	1	1	1	1	1	1	1
4	56	338827	6768189	1 N	1	4	4	4	1	0	0.1	0.4	0.5	0.4	0.1	0.0	0	0	36.4	10	0	0	0	0	0	0	2
5	56	339451	6768659	239 W	3	3	2	3	1	1	35.2	0.3	0.2	0.3	0.1	0.1	0	1	75	33.5	1	1	1	1	1	1	1
6	56	339229	6768505	196 S	0	2	3	7	0	1	0.0	5.1	25.5	0.7	0.0	0.1	0	1	60	0	0	0	0	0	0	0	1
7	56	339300	6768650	236 SW	2	3	3	10	0	3	72.0	0.3	17.1	1.0	0.0	0.3	0	0	50	53	1	1	1	1	0	1	1
8	56	338603	6768514	252 W	4	6	7	7	0	1	68.5	1.9	5.6	2.0	0.0	0.1	0	1	57	54	1	1	1	1	0	1	1
9	56	338079	6768264	294 NW	1	3	9	10	1	1	35.1	0.3	2.7	1.0	0.1	0.1	0	1	24	16	1	1	1	1	0	1	1
10	56	338759	6768130	162 S	3	4	6	9	1	3	43.0	0.4	15.5	0.9	0.1	0.3	0	1	65	80	1	1	1	1	0	1	0
11	56	338572	6767919	247 SW	3	3	6	10	1	2	38.0	0.3	1.0	1.0	0.1	0.3	0	1	50	32	1	1	0	0	1	1	0
12	56	338521	6767776	207 SW	2	4	6	8	0	1	47.3	0.4	0.7	0.8	0.0	0.1	0	0	36	21	1	1	1	1	0	1	1
13	56	338198	6768336	81 E	0	2	4	5	0	1	0.0	0.2	0.9	0.5	0.0	5.0	0	0	7.4	0	0	0	0	0	0	0	1
14	56	338663	6768348	260 W	0	3	11	14	1	1	0.0	0.3	1.3	2.3	0.3	0.1	0	0	35	6	0	0	0	0	0	0	1
15	56	338451	6767511	143 SE	4	3	9	6	1	3	40.1	0.3	1.3	0.6	0.1	0.3	1	1	69	66	1	1	1	1	1	1	1
16	56	338659	6767569	255 W	4	2	4	6	1	3	41.0	0.4	0.8	1.1	0.1	0.5	0	1	53.2	70	1	1	1	1	1	1	1
17	56	338763	6767760	270 W	1	3	4	10	1	1	0.1	0.3	15.1	1.5	0.1	0.2	0	0	30	12	0	1	0	0	0	0	3
18	56	338377	6767858	53 NE	0	2	6	6	1	1	0.0	0.2	6.9	0.7	0.2	0.2	0	0	49	35	0	0	0	0	0	0	0
19	56	338320	6768072	155 SE	0	2	2	5	1	1	0.0	20.2	5.0	0.5	0.1	0.2	0	0	41	4.5	0	0	0	0	0	0	2
20	56	337983	6767855	321 NW	1	1	3	1	0	2	0.1	0.1	0.5	0.1	0.0	0.2	0	0	31	0	0	0	0	0	0	0	1
21	56	337855	6767956	321 NW	1	2	4	10	0	1	3.0	0.2	5.6	1.1	0.0	0.1	0	0	36	40	1	1	1	0	0	1	1
22	56	338024	6768238	298 NW	1	2	3	9	0	2	0.1	0.2	2.1	0.9	0.0	0.3	0	0	38	23	1	0	0	0	0	1	2
23	56	338490	6768126	159 S	1	2	4	4	1	1	0.2	0.2	2.4	0.4	0.1	0.1	0	0	26	4	1	1	1	0	0	1	1
24	56	338800	6768481	170 S	0	1	5	4	1	1	0.0	0.1	2.0	0.8	0.1	0.1	0	0	73	21	0	0	0	0	0	0	0
25	56	338687	6768607	257 W	0	0	5	5	1	1	0.0	0.0	1.3	0.5	0.1	0.1	0	0	74	11	0	0	0	0	0	0	2
26	56	339225	6768408	255 W	2	1	7	5	0	1	0.2	0.1	3.9	0.5	0.0	0.1	0	0	26	21	0	1	0	0	0	1	0
27	56	338344	6768782	184 S	0	1	3	5	1	1	0.0	0.1	40.4	0.5	0.1	0.1	0	0	66	0	0	0	0	0	0	0	1
28	56	338658	6768836	183 S	0	2	1	6	0	1	0.0	0.4	25.0	0.6	0.0	0.1	0	0	33	0	0	0	0	0	0	0	1
29	56	339052	6768750	178 S	2	1	7	8	1	1	0.2	0.1	16.3	8.0	0.1	0.1	0	0	72	43	0	0	0	0	0	1	0
30	56	338535	6769002	25 NW	0	2	3	7	0	3	0.0	0.3	1.1	0.7	0.0	0.3	0	0	45	0	0	0	0	0	0	0	0
31	56	338402	6769323	191 S	2	3	3	8	1	0	3.1	0.4	1.3	0.8	0.1	0.0	0	1	38	2	0	0	1	0	1	0	2
32	56	338518	6769371	173 S	0	2	4	2	1	1	0.0	0.2	40.3	0.2	0.1	0.1	0	0	76	0	0	0	0	0	0	0	1
33	56	338361	6769671	326 NW	0	1	2	2	0	1	0.0	0.1	0.2	0.2	0.0	0.1	0	0	7.4	0	0	0	0	0	0	0	2
34	56	338569	6769788	275 W	0	1	2	4	0	1	0.0	0.3	0.3	0.5	0.0	0.2	0	0	26	0	0	0	0	0	0	0	2
35	56	338671	6769805	262 W	0	2	6	4	0	1	0.0	5.0	3.0	0.4	0.0	0.1	0	0	24.2	0	0	0	0	0	0	0	0
36	56	339043	6768195	315 W	1	1	6	4	0	1	0.1	0.1	3.1	0.4	0.0	0.1	0	0	46.6	22	0	0	0	0	0	0	1
37	56	338564	6768372	177 S	0	1	6	7	1	1	0.0	0.1	4.4	0.7	0.1	0.1	0	0	72	64	0	1	0	0	0	1	1
38	56	338685	6768653	235 SW	1	3	2	1	0	0	50.0	0.3	0.2	0.1	0.0	0.0	0	0	63.6	42	0	1	1	1	1	0	1
39	56	338676	6768698	56 NE	0	2	3	4	0	0	0.0	0.4	1.8	0.5	0.0	0.0	0	0	81.6	0	0	0	0	0	0	0	1
40	56	339305	6768405	40 NE	1	3	5	4	0	2	25.0	0.3	0.8	0.4	0.0	0.2	0	1	29	23	0	0	0	1	1	0	1
41	56	338392	6768534	259 W	1	2	3	4	1	1	10.0	0.5	20.5	0.4	0.1	0.1	0	1	23.2	5	1	1	1	0	1	0	1
42	56	338533	6768664	18 N	1	3	1	7	0	1	20.0	0.3	3.0	0.7	0.0	0.1	1	1	55	0	0	0	0	0	0	0	2
Notes:											•																

Notes:

x = present

HTE = High Threat Exotic HBT = Hollow-bearing Tree P/A = Presence / Absence

DBH = Diameter at Breast Heght;

1 = Present; 0 = Absent

Logs = Length of logs with more than 10cm diameter

0470861_BDAR Table G.1: page 1 of 1

BAM Plot photographs



Plot 1: horizontal view from 0m midline (11 Sep 2018)



Plot 1: vertical view from 0m midline (11 Sep 2018)



Plot 2: horizontal view from 0m midline (11 Sep 2018)



Plot 2: vertical view from 0m midline (11 Sep 2018)



Plot 3: horizontal view from 0m midline (11 Sep 2018)



Plot 3: vertical view from 0m midline (11 Sep 2018)



Plot 4: horizontal view form 0m midline (12 Sep 2018)



Plot 4: vertical view from 0m midline (12 Sep 2018)



Plot 5: horizontal view from 0m midline (12 Sep 2018)



Plot 5: vertical view from 0m midline (12 Sep 2018)



Plot 6: horizontal view from 0m midline (11 Dec 2018)




Plot 6: vertical view form 0m midline (11 Dec 2018)



Plot 7: horizontal view from 0m midline (12 Dec 2018)



Plot 8: horizontal view from 0m midline (13 Dec 2018)



Plot 7: vertical view from 0m midline (12 Dec 2018)



Plot 8: vertical view from 0m midline (13 Dec 2018)



Plot 9: horizontal view from 0m midline (13 Dec 2018)



Plot 9: vertical view from 0m midline (13 Dec 2018)



Plot 10: horizontal view from 0m midline (13 Dec 2018) Plot 10: vertical view from 0m midline (13 Dec 2018)





Plot 11: horizontal view from 0m midline (13 Dec 2018) Plot 11: vertical view from 0m midline (13 Dec 2018)





Plot 12: horizontal view from 0m midline (13 Dec 2018)



Plot 12: vertical view from 0m midline (13 Dec 2018)



Plot 13: horizontal view from 0m midline (14 Dec 2018)



Plot 13: vertical view from 0m midline (14 Dec 2018)



Plot 14: horizontal view from 0m midline (14 Dec 2018) Plot 14: vertical view from 0m midline (14 Dec 2018)





Plot 15: horizontal view from 0m midline (25 Mar 2019)



Plot 15: vertical view from 0m midline (25 Mar 2019)



Plot 16: horizontal view from 0m midline (25 Mar 2019)



Plot 16: vertical view from 0m midline (25 Mar 2019)



Plot 17: horizontal view from 0m midline (25 Mar 2019)



Plot 17: vertical view from 0m midline (25 Mar 2019)



Plot 18: horizontal view from 0m midline (25 Mar 2019)



Plot 18: vertical view from 0m midline (25 Mar 2019)



Plot 19: horizontal view from 0m midline (25 Mar 2019)



Plot 19: vertical view from 0m midline (25 Mar 2019)



Plot 20: horizontal view from 0m midline (26 Mar 2019)



Plot 20:horizontal view from 0m midline (26 Mar 2019)



Plot 21: horizontal view from 0m midline (26 Mar 2019)



Plot 21: horizontal view from 0m midline (26 Mar 2019)



Plot 22: horizontal view from 0m midline (26 Mar 2019)



Plot 22: horizontal view from 0m midline (26 Mar 2019)



Plot 23: horizontal view from 0m midline (26 Mar 2019)



Plot 23: horizontal view from 0m midline (26 Mar 2019)



Plot 24: horizontal view from 0m midline (26 Mar 2019)



Plot 24: horizontal view from 0m midline (26 Mar 2019)



Plot 25: horizontal view from 0m midline (26 Mar 2019)



Plot 25: horizontal view from 0m midline (26 Mar 2019)



Plot 26: horizontal view from 0m midline (26 Mar 2019)



Plot 26: horizontal view from 0m midline (26 Mar 2019)



Plot 27: horizontal view from 0m midline (27 Mar 2019)



Plot 27: horizontal view from 0m midline (27 Mar 2019)



Plot 28: horizontal view from 0m midline (27 Mar 2019)



Plot 28: horizontal view from 0m midline (27 Mar 2019)



Plot 29: horizontal view from 0m midline (27 Mar 2019)



Plot 29: horizontal view from 0m midline (27 Mar 2019)



Plot 30: horizontal view from 0m midline (27 Mar 2019)



Plot 30: horizontal view from 0m midline (27 Mar 2019)



Plot 31: horizontal view from 0m midline (27 Mar 2019)



Plot 31: horizontal view from 0m midline (27 Mar 2019)



Plot 32: horizontal view from 0m midline (27 Mar 2019)



Plot 32: horizontal view from 0m midline (27 Mar 2019)



Plot 33: horizontal view from 0m midline (27 Mar 2019)



Plot 33: horizontal view from 0m midline (27 Mar 2019)



Plot 34: horizontal view from 0m midline (27 Mar 2019)



Plot 34: horizontal view from 0m midline (27 Mar 2019)



Plot 35: horizontal view from 0m midline (27 Mar 2019)



Plot 35: horizontal view from 0m midline (27 Mar 2019)



Plot 36: horizontal view from 0m midline (28 Mar 2019)



Plot 36: horizontal view from 0m midline (28 Mar 2019)



Plot 37: horizontal view from 0m midline (28 Mar 2019)



Plot 37: horizontal view from 0m midline (28 Mar 2019)



Plot 38: horizontal view from 0m midline (28 Mar 2019)



Plot 38: horizontal view from 0m midline (28 Mar 2019)



Plot 39: horizontal view from 0m midline (28 Mar 2019)



Plot 39: horizontal view from 0m midline (28 Mar 2019)



Plot 40: horizontal view from 0m midline (28 Mar 2019)



Plot 40: horizontal view from 0m midline (28 Mar 2019)



Plot 41: horizontal view from 0m midline (28 Mar 2019)



Plot 41: horizontal view from 0m midline (28 Mar 2019)



Plot 42: horizontal view from 0m midline (28 Mar 2019) Plot 42: horizontal view from 0m midline (28 Mar 2019)



GAIA BONSHAW SOLAR EIS Revised Biodiversity Development	Assessment Report	
ADDENDIV II	FLORA AND FAUNA SPECIES LIS	NTC
APPENDIX H	FLORA AND FAUNA SPECIES LIS	013

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

Table H.1 Bird Species Records

Family	Common Name	Scientific Name	BC Act	EPBC Act	B1	B2	В3	B4	B5	B6	B7	Incidentals
Acanthizidae	Weebill	Smicrornis brevirostris				1	2					I
Acanthizidae	White-throated Gerygone	Gerygone albogularis										II
Acanthizidae	Inland Thornbill	Acanthiza apicalis										I
Acanthizidae	Speckled Warbler^	Pyrrholaemus sagittatus										I
Acanthizidae	Yellow-rumped Thornbill	Acanthiza chrysorrhoa										I
Accipitridae	Wedge-tailed Eagle^	Aquila audax										I
Accipitridae	Whistling Kite	Haliastur sphenurus										I
Aegothelidae	Australian Owlet-nightjar	Aegotheles cristatus										I
Anatidae	Pacific Black-duck	Anas superciliosa								10		I
Artamidae	Australian Magpie	Cracticus tibicen			2	1		1		2	3	I
Artamidae	Grey Butcherbird	Cracticus torquatus			1	1	1	1			1	I
Artamidae	Pied Butcherbird	Pied Butcherbird								1		I
Artamidae	Pied Currawong	Strepera graculina			1		1	1				I
Cacatuidae	Galah	Eolophus roseicapillus						1		5	4	I
Cacatuidae	Little Corella	Cacatua sanguinea								30		
Cacatuidae	Sulphur-crested Coockatoo	Cacatua galerita								6		I
Campephagidae	Cicadabird	Coracina tenuirostris		Ma		1						I
Campephagidae	Black-faced Cuckooshrike	Coracina novaehollandiae										I
Campephagidae	White-winged Triller	Lalage sueurii										I
Charadriidae	Masked Lapwing	Vanellus miles										I

Family	Common Name	Scientific Name	BC Act	EPBC Act	B1	B2	В3	B4	B5	B6	B7	Incidentals
Climacteridae	Brown Treecreeper^ (eastern subspecies)	Climacteris picumnus victoriae	V									I
Columbidae	Crested Pigeon	Ocyphaps lophotes			2	2	1	1		1		I
Columbidae	Common Bronzewing	Phaps chalcoptera										I
Columbidae	Peaceful Dove	Geopelia striata										II
Corcoracidae	Apostlebird	Struthidea cinerea			6	2				3	2	I
Corcoracidae	White-winged Chough	Corcorax melanorhamphos			4							I
Corvidae	Australian Raven	Corvus coronoides			2							I
Corvidae	Torresian Crow	Corvus orru				3		1				I
Cuculidae	Channel-billed Cockoo	Scythrops novaehollandiae								1		I
Cuculidae	Eastern Koel	Eudynamys orientalis			1						1	
Estrildidae	Double-barred Finch	Taeniopygia bichenovii										I
Falconidae	Nankeen Kestrel	Falco cenghroides										I
Meliphagidae	Blue-faced Honeyeater	Entomyzon cyanotis						1				
Meliphagidae	Lewin's Honeyeaater	Meliphaga lewinii					1				1	
Meliphagidae	Little Friarbird	Philemon citreogularis			2	3	1		1		3	I
Meliphagidae	Noisy Miner	Manorina melanocephala			5	4	4	11		2	4	I
Meliphagidae	Striped Honeyeater	Plectorhyncha lanceolata					1					I
Meliphagidae	Brown Honeyeater	Lichmera indistincta										I
Monarchidae	Magpie-lark	Grallina cyanoleuca			2	2		1		9	1	I
Motacillidae	Australasian Pipit	Anthus novaeseelandiae						1			1	I
Nectariniidae	Mistletoebird	Dicaeum hirundinaceum										I

18 May 2020 Page H2 www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

Family	Common Name	Scientific Name	BC Act	EPBC Act	B1	B2	В3	B4	B5	В6	B7	Incidentals
Oriolidae	Olive-backed Oriole	Oriolus sagitattus										I
Pachycephalidae	Grey Shrike-thrush^	Colluricincla harmonica										I
Pachycephalidae	Rufus Whistler	Pachycephala rufiventris										I
Pardalotidae	Striated Pardalote	Pardalotus striatus			1			1		2	1	I
Petrocidae	Eastern Yellow Robin^	Eopsaltria australis										I
Petrocidae	Jacky Winter^	Microeca fascinans										I
Petrocidae	Red-capped Robin^	Petroica goodenovii										I
Pomatostomidae	Grey-crowned Babbler	Pomatostomus temporalis	V							1	4	
Psittacidae	Pale-headed Rosella	Platycercus adscitus			1	5					1	I
Psittacidae	Red-rumped Parrot	Psephotus haematonotus				10	4	1	1			I
Psittacidae	Red-winged Parrot	Aprosmictus erythropterus				6	1					
Psittacidae	Eastern Rosella	Platycercus eximius										I
Rhipiduridae	Grey Fantail	Rhipidura albiscapa										I
Rhipiduridae	Willie Wagtail	Rhipidura leucophrys			1	1		1			2	I

Notes: # = number of individuals; I - one record; II = two records; ^ - recorded adjacent to the Subject Land only; V - Vulnerable; Ma - Marine

18 May 2020 Page H3 www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

Table H.2 Reptile species records

Family	Common Name	Scientific Name	BC Act	EPBC Act	R1	R2	R3	R4	R5	R6	R7	R8	Incidentals
Agamidae	Eastern Bearded Dragon	Pogona barbata											I
Varanidae	Monitor Lizard	Varanus varius											1
Gekkonidae	Bynoe's Gecko	Heteronotia binoei											1
Scincidae	Tree Skink	Egernia striolata											I
Scincidae	Tree-base Litter-skink, Rainbow Skink	Lygisaurus foliorum											I
Gekkonidae	A Gecko	Gehyra sp.						1					
	Small brown gecko												1
	Skink												I
	Turtle												I

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page H4

Table H.3 Mammal species records

Family	Common Name	Scientific Name	BC Act	EPBC Act	Incidentals
Bovidae	European Cattle*	Bos taurus			Х
Bovidae	Sheep*	Ovis aries			х
Canidae	Dog*	Canis lupus familiaris			х
Leporidae	European Rabbit*	Oryctolagus cuniculus			scat, x
Macropodidae	Common Wallaroo	Macropus robustus			х
Macropodidae	Eastern Grey Kangaroo	Macropus giganteus			х
Macropodidae	Red-necked Wallaby	Macropus rufogriseus			х
Peramelidae	Northern Brown Bandicoot	Isoodon macrourus			х
Phalangeridae	Common Brushtail Possum	Trichosurus vulpecula			х
Suidae	Feral Pig*	Sus scrofa			Х
Fachyglossidae	Short-beaked Echidna	Tachyglossus aculeatus			Scat, diggings

^{*}Introduced species

Table H.4 Amphibian species list

Family	Common Name	Scientific Name	F1	F2	F3	Incidentals
Hyalidae	Desert Tree Frog	Litoria rubella	С			
Hylidae	Broad-palmed Frog	Litoria latopalmata				0
Hylidae	Peron's Tree Frog	Litoria peronii				0

Notes: C = heard calling; O = observed;

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page H5

Table H.5 **Microchiropteran Bat Species List**

Family	Common Name	Scientific Name	BC Act	EPBC Act	S2	S3	S4	S5	S6	S7	S8
Molossidae	Austronomus australis	White-striped Freetail-bat	-		D	D	D	D	D	D	D
Vespertilionidae	Chalinolobus gouldii	Chalinolobus gouldii Gould's Wattled Bat			D	D	D	D	D	D	D
Vespertilionidae	Chalinolobus morio	Chocolate Wattled Bat			D	D	D	D	D	D	D
Vespertilionidae	Chalinolobus picatus	Little Pied Bat	V		D	D	D	D	Р	D	D
Miniopteridae	Miniopterus orianae oceanensis	Eastern Bent-winged Bat	V		D	D	D	D	D	D	D
Molossidae	Ozimops planiceps / O. petersi	South-eastern Free-tailed Bat / Inland Free-tailed Bat	-	-	Р	Р	Р	Р	Р	Р	Р
Molossidae	Ozimops ridei	Ride's Free-tailed Bat			D	D	D	D	D	D	D
Emballonuridae	Saccolaimus flaviventris	Yellow-bellied Sheath-tailed Bat	V		D	D	D	D	D	D	D
Vespertilionidae	Scotorepens balstoni	Inland Broad-nosed Bat			D	D	D	D	D	D	Р
Vespertilionidae	Scotorepens greyii	Little Broadnosed Bat			17	15	46	210	26	10	27
Molossidae	Mormopterus eleryi *	Bristle-faced Free-tailed Bat	Е		5		11	25	1	12	
Vespertilionidae	Vespadelus troughtoni	Eastern Cave Bat	V		Р	Р	Р	Р	Р	Р	Р
Vespertilionidae	Vespadelus vulturnus	Little Forest Bat			D	D	D	D	D	D	D
Nyctophilus species	group formed by the three species belo	DW			D	D	D	D	D	D	
Vespertilionidae	Nyctophilus corbeni	Corben's Long-eared Bat	V	V							
Vespertilionidae	Nyctophilus geoffroyi	Lesser Long-eared Bat									
Vespertilionidae	Nyctophilus gouldi	Gould's Long-eared Bat									
Vespertilionidae/ Molosidae	Scotorepens greyii or Chalinolobus nigrogriseus ^v	Little Broad-nosed Bat or Hoary Wattled Bat	V		Р	Р	Р	Р	Р	Р	Р
Vestertilionidae /	Scotorepens greyii or Falsistrellus tasmaniensis ^v	Little Broad-nosed Bat or Eastern False Pipistrelle	V		Р	Р	Р	Р	Р	Р	

D = Definite call identification, i.e. at least one call attributed unequivocally to the species; P = Possible call identification, i.e. calls like those of the species were recorded, but were not reliably identified; V = Vulnerable; * reported as Setirostris eleryi by Balance! Environmental (see Appendix I)

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020 Page H6

GAIA BONSHAW SOLAR EIS

Revised Biodiversity Development Assessment Report

Table H.6 Flora Species List

 www.erm.com
 Version: 2.0
 Project No.: 0470861
 Client: GAIA Australia
 18 May 2020
 Page H7

Table H.7 Summary of Hollow Bearing Trees Observations

No	Latitude	Longitud	Scientific Name	Common Name	Туре	DBH	No Hollows	Size (cm)	Notes
1	-29.1927	151.3394	Eucalyptus moluccana	Grey Box	Т	-	1	15	-
2	-29.194	151.3394	Eucalyptus dealbata	Tumbledown Red Gum	S	-	1	20	-
3	-29.1943	151.3403	Eucalyptus moluccana	Grey Box	Т	-	1	5	-
4	-29.1952	151.3413	Angophora floribunda	Rough-barked Apple	Т	-	1	45	-
5	-29.1942	151.3419	Angophora floribunda	Rough-barked Apple	Т	-	1	15	-
6	-29.1968	151.3438	Eucalyptus moluccana	Grey Box	Т	-	1	30	-
7	-29.1972	151.3441	-	-	S	-	1	40	-
8	-29.1973	151.3444	-	-	S	-	1	40	-
9	-29.197	151.3446	-	-	S	-	1	-	Large top opening
10	-29.1982	151.3432	Eucalyptus melanophloia	Silver-leaved Ironbark	S	-	1	10	-
11	-29.2062	151.345	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	25	-
12	-29.2053	151.3453	-	-	S	-	1	25	-
13	-29.2098	151.3414	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	10	-
14	-29.2098	151.3415	-	-	S	-	1	35	-
15	-29.2056	151.3366	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	5	-
16	-29.2084	151.3373	-	-	S	-	1	10	-
17	-29.2095	151.3375	Eucalyptus moluccana	Grey Box	Т	-	1	5	-
18	-29.2097	151.3378	Eucalyptus moluccana	Grey Box	Т	-	1	10	-
19	-29.21	151.3331	Corymbia dolichocarpa	Long-fruited Bloodwood	Т	-	1	20	-
20	-29.2095	151.3336	-	-	S	-	1	10	-
21	-29.2094	151.3335	-	-	S	-	1	5	-
22	-29.2093	151.3301	-	-	S	-	1	-	Split at base to 1.75m
23	-29.2086	151.3311	-	-	S	-	1	10	-
24	-29.2058	151.3318	-	-	S	-	1	5	-
25	-29.2014	151.3478	-	-	S	-	1	35	-
26	-29.2022	151.3479	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	35	-
27	-29.2018	151.3485	Eucalyptus dealbata	Tumbledown Red Gum	Т	-	1	20	-
28	-29.2019	151.3485	Eucalyptus dealbata	Tumbledown Red Gum	Т	-	1	25	-
29	-29.2019	151.3454	-	-	S	-	1	25	-
30	-29.2068	151.3424	Eucalyptus moluccana	Grey Box	S	-	1	40	-
31	-29.2073	151.3423	Eucalyptus moluccana	Grey Box	S	-	1	15	-
32	-29.205	151.3368	-	-	S	-	1	15	-
33	-29.1968	151.3401	Eucalyptus moluccana	Grey Box	Т	-	1	15	-

No	Latitude	Longitud	Scientific Name	Common Name	Туре	DBH	No Hollows	Size (cm)	Notes
34	-29.2095	151.3319	Corymbia dolichocarpa	Long-fruited Bloodwood	Т	-	1	20	-
35	-29.194	151.3377	Eucalyptus moluccana	Grey Box	Т	-	3	20	Largest hollow 20cm D
36	-29.194	151.3381	Eucalyptus moluccana	Grey Box	Т	-	1	10	-
37	-29.1961	151.3387	Eucalyptus moluccana	Grey Box	S	-	1	30	-
38	-29.1977	151.3399	-	-	S	-	1	5	-
39	-29.1983	151.3399	Eucalyptus moluccana	Grey Box	S	80	1	30	-
40	-29.2005	151.3376	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	20	-
41	-29.2029	151.3369	Eucalyptus moluccana	Grey Box	T, S	-	1	10	-
42	-29.2022	151.336	Eucalyptus moluccana	Grey Box	Т	-	1	15	-
43	-29.201	151.3385	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	10	-
44	-29.2022	151.3378	Eucalyptus melanophloia	Silver-leaved Ironbark	T, S*	-	1	<10	-
45	-29.2095	151.332	Eucalyptus moluccana	Grey Box	Т	-	1	30	-
46	-29.2018	151.3426	-	-	S	-	1	15	-
47	-29.2011	151.3451	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	15	-
48	-29.2016	151.3383	-	-	S	-	1	15	-
49	-29.2056	151.3399	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	40	Hollow at approx 6m above ground level
50	-29.2068	151.3395	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	7	-
51	-29.2087	151.3384	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	10	-
52	-29.2107	151.3358	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	10	-
53	-29.21	151.3415	Corymbia dolichocarpa	Long-fruited Bloodwood	Т	-	1	30	-
54	-29.2077	151.3418	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	20	-
55	-29.2074	151.3411	-	-	Т	-	1	15	-
56	-29.2071	151.3409	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	35	-
57	-29.2062	151.3415	-	-	Т	-	1	10	-
58	-29.2063	151.3446	Eucalyptus dealbata	Tumbledown Red Gum	Т	-	1	15	-
59	-29.2	151.344	Eucalyptus melanophloia	Silver-leaved Ironbark	Т	-	1	15	-
60	-29.1996	151.3393	Eucalyptus moluccana	Grey Box	Т	-	1	40	-

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development	Assessment Panort
Revised biodiversity Development	ASSESSITIETE REPORT
APPENDIX I	MICROCHIROPTERAN BAT CALL REPORT

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020



Microbat Call Identification Report

Prepared for ("Client"):	ERM
Survey location/project name:	Bonshaw area (NSW)
Survey dates:	10-14 December 2018
Client project reference:	0470861
Job no.:	ERM-1901
Report date:	22 February 2019

DISCLAIMER:

© Copyright – Balance! Environmental, ABN 75 795 804 356. This document and its content are copyright and may not be copied, reproduced or distributed (in whole or part) without the prior written permission of Balance! Environmental other than by the Client for the purposes authorised by Balance! Environmental ("Intended Purpose"). To the extent that the Intended Purpose requires the disclosure of this document and/or its content to a third party, the Client must procure such agreements, acknowledgements and undertakings as may be necessary to ensure that the third party does not copy, reproduce, or distribute this document and its content other than for the Intended Purpose. This disclaimer does not limit any rights Balance! Environmental may have under the Copyright Act 1968 (Cth).

The Client acknowledges that the Final Report is intended for the sole use of the Client, and only to be used for the Intended Purpose. Any representation or recommendation contained in the Final Report is made only to the Client. Balance! Environmental will not be liable for any loss or damage whatsoever arising from the use and/or reliance on the Final Report by any third party.



Methods

Data received

Balance! Environmental received 8351 full-spectrum (WAV format) audio files, recorded using Song Meter SM2BAT detectors at seven sites (S2-S8) over four consecutive nights (10-13 December 2018). A file note included with the submitted data indicated that another site ("S1") was sampled but no data was obtained.

Call identification

Data were analysed in *Kaleidoscope Pro* (Version 5.1.7; Wildlife Acoustics, Maynard MA, USA). The first-pass analysis utilised the "Cluster Analysis" function, to scan all WAV files and automatically group recognised bat calls into clusters with similar call characteristics (based on zero-crossing analysis). Clusters were then assigned species identifications by manually reviewing call spectrograms and derived metrics and comparing them with those of regionally-relevant reference calls and/or with published call descriptions (Reinhold et al. 2001; Pennay *et al.* 2004). Consideration was also given to the probability of species' occurrence based on published distribution information (e.g. Churchill 2008; van Dyck *et al.* 2013) and on-line database records (e.g. http://www.ala.org.au).

Species identification was based largely on search-phase call sequences; however, where good-quality foraging sequences were available (*i.e.* a call sequence with contiguous search-phase, attack-phase and feeding-buzz components), those calls were used to provide additional evidence of some species' presence. The feeding buzzes of *Miniopterus* species (bent-winged bats) and some Molossids (free-tailed bats) are quite distinctive, compared with those of Vespertilionids (vesper bats) with which they often share search-phase characteristics (Corben 2010).

Reporting standard

The format and content of this report follows Australasian Bat Society standards for the interpretation and reporting of bat call data (Reardon 2003), available on-line at http://www.ausbats.org.au/.

Species nomenclature follows Jackson & Groves (2015), which uses several new genus/species names compared with common field guides (*e.g.* Churchill 2008; Van Dyck *et al.* 2013). New names used in this report include:

Ozimops ridei (Ride's Free-tailed Bat) - formerly Mormopterus ridei and M. 'species 2';

- O. petersi (Inland Free-tailed Bat) formerly M. petersi and M. 'species 3';
- O. planiceps (Southern free-tailed Bat) formerly M. planiceps and M. 'species 4';

Setirostris eleryi (Bristle-faced Free-tailed Bat) – formerly M. eleryi and M. 'species 6'; and

Miniopterus orianae (Large or Eastern Bent-winged Bat) – formerly M. schreibersii.



Results & Discussion

Cluster analysis output

The Cluster Analysis recognised 7705 bat calls and grouped them into 54 clusters; however, many clusters contained slight variants of the same species' calls. Three-quarters of the recognised calls were positively identified, with the other 25% unable to be allocated reliably to a single species. These were assigned to "unresolved species" groups. A breakdown of the numbers of calls attributed to species or unresolved groups for each site is provided at **Appendix 1**.

Species recorded

At least 12 and up to 16 species were recorded during the Bonshaw survey (see **Table 1**). Eleven call types were positively identified to individual species, while one call type was attributed to the genus *Nyctophilus*, within which species cannot be reliably differentiated using call characteristics. Three *Nyctophilus* species potentially occur in the study area: *N. corbeni; N. geoffroyi;* and *N. gouldi.* Half (3907) of the total recorded calls were reliably attributed to just three species: *Miniopterus orianae oceanensis* (Eastern Bent-winged Bat); *Ozimops ridei* (Eastern Free-tailed Bat); and *Saccolaimus flaviventris* (Yellow-bellied Sheath-tailed Bat).

Several call types that potentially represented multiple species were allocated to "unresolved" species groups. Where calls were allocated to one these "unresolved" groups, all group members are listed as "possibly present" in **Table 1** unless other, more typical calls form one or more members were also observed for the relevant site. Three of the "unresolved" species groups included species that were not otherwise found in the positively-identified call data. These include:

- Ozimops ridei / O. petersi / O. planiceps
 - O. ridei was positively identified where calls had typical Ozimops pulse-shape characteristics (predominantly flat search-phase pulses) and characteristic frequency (Fc) of 30-35 kHz
 - o Calls allocated to this "unresolved" group had similar pulse shapes but Fc27-30 kHz
 - o It is likely these were all low-frequency variants of *O. ridei*, but it is possible that *O. planiceps* and (remotely possible) *O. petersi* also occur in the study area
- Vespadelus vulturnus / V. troughtoni
 - V. vulturnus positively identified from numerous calls having typical steep, curvilinear pulses with hooked bodies (up-swept tail) and Fc~46-48 kHz
 - A number of calls with the same typical Vespadelus pulse-shape had Fc~48-50 kHz may have been form either of these species
- Vespadelus spp. / Chalinolobus morio
 - Calls were similar to above group but had mixed pulse shape and duration, varying from the shorter, hooked *Vespadelus* pulse-shape to longer, flatter pulse bodies with no tail or down-swept tail more typical of *C. morio*
 - o *C. morio* calls were reliably identified only where all pulses showed consistent shape with flatter characteristic section and obvious down-swept tail

Sample spectrograms of all call-types are shown at Appendix 1.



Threatened species

Several of the identified species are listed as threatened under State and Commonwealth nature conservation legislation (Environment Protection and Biodiversity Conservation Act 1999 [EPBC Act] – Cwlth; Biodiversity Conservation Act 2016 [BCA] – New South Wales; Nature Conservation Act 1992 [NCA] – Queensland). These include:

- Chalinolobus picatus (Little Pied Bat) Vulnerable, BCA
- Nyctophilus corbeni (Corben's Long-eared Bat) Vulnerable, EPBC, BCA, NCA
 - Note that this species was not positively identified, but may have been responsible for some of the calls allocated to the *Nyctophilus* genus
- Vespadelus troughtoni (Eastern Cave Bat) Vulnerable, BCA
 - Note that this species was not positively identified but may have been responsible for some of the unresolved calls allocated to two multi-species groups (see previous section)
- Miniopterus orianae oceanensis (Eastern Bent-winged Bat) Vulnerable, BCA
- Setirostris eleryi (Bristle-faced Free-tailed Bat) Endangered, BCA
- Saccolaimus flaviventris (Yellow-bellied Sheath-tailed Bat) Vulnerable, BCA

Table 1. Microbat species recorded during the ERM Bonshaw survey, 10-14 December 2018.

- ♦ = 'definite' at least one call was attributed unequivocally to the species
- \Box = 'possible' calls like those of the species were recorded, but were not reliably identified

Site code:	S2	S3	S4	S5	S6	S7	S8
Chalinolobus gouldii	•	*	•	*	*	*	*
Chalinolobus morio	*	*	*	*	*	*	+
Chalinolobus picatus	*	*	♦	*		*	*
Nyctophilus species	*	*	*	*	*	*	
Scotorepens balstoni	*	*	•	*	*	*	
Scotorepens greyii	*	*	*	*	*	*	+
Vespadelus vulturnus	*	*	♦	*	*	*	*
Vespadelus troughtoni							
Miniopterus orianae oceanensis	*	*	*	*	*	*	*
Austronomus australis	*	*	*	*	*	*	+
Ozimops ridei	*	*	•	*	*	*	+
Ozimops planiceps / O. petersi							
Setirostris eleryi	*	*	*	*	*	*	*
Saccolaimus flaviventris	+	*	+	*	+	*	♦



References

- Churchill, S. (2008). Australian Bats. Jacana Books, Allen & Unwin; Sydney.
- Corben, C. (2010). Feeding Buzzes. Australasian Bat Society Newsletter 35, 40-44.
- Jackson, S. and Groves, C. (2015). *Taxonomy of Australian Mammals*. CSIRO Publishing, Melbourne.
- Pennay, M., Law, B. and Reinhold, L. (2004). *Bat Calls of New South Wales*. Department of Environment and Conservation, Hurstville.
- Reardon, T. (2003). Standards in bat detector based surveys. *Australasian Bat Society Newsletter* **20**, 41-43.
- Reinhold, L., Law, B., Ford, G. and Pennay, M. (2001). *Key to the bat calls of south-east Queensland and north-east New South Wales*. Department of Natural Resources and Mines, Brisbane.
- van Dyck, S., Gynther, I. and Baker, A. (ed.) (2013). *Field Companion to the Mammals of Australia*. New Holland; Sydney.



Glossary

Technical terms used in this report are described in the following table.

Approach phase The part of a bat *call* emitted as the bat starts to home in on a detected

prey item; a transitional series of *pulses* between the *search phase* and *feeding buzz*, that become progressively steeper and shorter in

duration.

Call Refers to a single bat call, made up of a series of individual sound

pulses in one or more phases (search, approach, feeding buzz).

CF (=Constant Frequency)

A type of *pulse* in which the dominant component consists of a more-

or-less 'pure tone' of sound at a Constant Frequency; with *shape* appearing flat on the sonogram. Often also contains a brief FM component at the beginning and/or end of the CF component (viz. FM-

CF-FM).

Characteristic frequency (Fc)

The frequency of the flattest part of a *pulse*; usually the lowest

frequency reached in the qCF component of a pulse. This is often the

primary diagnostic feature for species identification.

Duration The time period from the beginning of a *pulse* to the end of the pulse.

Feeding buzz The terminal part of a call, following the approach phase, emitted as

the bat catches a prey item; a distinctive, rapid series of very steep,

very short-duration pulses.

FM (=Frequency Modulated) A type of *pulse* in which there is substantial change in frequency from

beginning to end; shape ranges from almost vertical and linear through

varying degrees of curvature.

FC range Refers to the range of frequencies occupied by the *characteristic*

frequency section of pulses within a call or set of calls.

Frequency sweep or "band-width" The range of frequencies through which a *pulse* sweeps from

beginning to end; Maximum frequency (Fmax) - minimum frequency

(Fmin).

Knee The transitional part of a *pulse* between the initial (usually steeper)

frequency sweep and the *characteristic frequency* section (usually flatter); time to knee (Tk) and frequency of knee (Fk) can be diagnostic

for some species.

Pulse An individual pulse of sound within a bat call; the shape, duration and

characteristic frequency of a pulse are the key diagnostic features used

to differentiate species.

Pulse body The part of the *pulse* between the *knee* and *tail* and containing the

characteristic frequency section.

Pulse shape The general appearance of a *pulse* on the sonogram, described using

relative terms related to features such as slope and degree of

curvature. See also CF, qCF and FM.

qCF (=quasi Constant Frequency) A type of *pulse* in which there is very little change in frequency from

beginning to end; shape appears to be almost flat. Some pulses also contain an FM component at the beginning and/or end of the qCF

component (viz. FM-qCF).

Search phase The part of a bat *call* generally required for reliable species diagnosis.

A consistent series of *pulses* emitted by a bat that is searching for prey or and/or navigating through its habitat. Search phase pulses generally have longer duration, flatter slope and more consistent shape than

approach phase and feeding buzz pulses.

Sequence Literally, a sequence of *pulses* that may be from one or more bats; but

generally refers to a call or part (e.g. phase) of a call.

Tail The final component of a pulse, following the characteristic frequency

section; may consist of a short or long sweep of frequencies either

upward or downward from the Fc; or may be absent.




Appendix 1 Bats recorded during the Bonshaw survey: number of calls allocated to each species or unresolved group.


Site code:	S2	S3	S4	S5	S6	S7	S8	Species totals
Positively identified calls								
Chalinolobus gouldii	46	44	33	267	43	9	13	455
Chalinolobus morio	1	3	1	3	1	3	3	15
Chalinolobus picatus	10	12	3	54		5	7	91
Nyctophilus species	5	2	1	11	5	13		37
Scotorepens balstoni	2	1	2	15	12	2		34
Scotorepens greyii	19	6	32	273	17	8	18	373
Vespadelus vulturnus	11	8	15	62	2	7	3	108
Miniopterus orianae oceanensis		150	10	390	19	52	67	757
Austronomus australis	24	38	13	16	62	26	14	193
Ozimops ridei	351	227	98	440	216	163	158	1653
Setirostris eleryi	70	40	79	270	31	48	26	564
Saccolaimus flaviventris		491	104	377	79	30	73	1497
Unresolved calls								
C. gouldii/O. ridei	181	106	64	257	97	60	60	825
C. gouldii/S. balstoni	16	8	20	111	42	2	6	205
S. balstoni/O. ridei	26	11	4	44	9	4		98
S. greyii/C. picatus	1	5	4	31	3	5		49
S. greyii/S. eleryi		4	7	44	5	6		66
Ozimops species	59	31	15	31	25	31	34	226
V. vulturnus/M. o. oceanensis	32	27	10	75	5	13	3	165
V. vulturnus/V. troughtoni		10	117	66	10	10	5	229
V. vulturnus/V. troughtoni/C. morio	6	22	16	12	4	3	2	65
Total calls from site:	1283	1246	648	2849	687	500	492	7705




Appendix 2 Representative call sequences from the Bonshaw survey, 10-14 December 2018. *Kaleidoscope* spectrograms with zero-crossing overlay (white); time between pulses removed











GAIA BONSHAW SOLAR EIS Revised Biodiversity Developm	ent Assessment Report
APPENDIX J	BAM REPORTS – VEGETATION INTEGRITY ASSESSMENT

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

0470861 GAIA Bonshaw Solar BDAR_Version 4.docx



BAM Credit Summary Report

Proposal Details

Assessment Id Proposal Name BAM data last updated *

00015157/BAAS18113/19/00015159 **Bonshaw Solar Farm** 26/11/2019

Assessor Name Report Created BAM Data version *

Matt Jenkins 17/03/2020 22

Date Finalised Assessor Number **BAM Case Status**

To be finalised BAAS18029 Open

Assessment Type Assessment Revision

Part 4 Developments (General)

Ecosystem credits for plant communities types (PCT), ecological communities & threatened species habitat

Zone	Vegetation zone name	Vegetation integrity loss / gain	Area (ha)	Constant	Species sensitivity to gain class (for BRW)	Biodiversity risk weighting	Potential SAII	Ecosystem credits	
Grey Bo	Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion								
1	516_Very_Low	20.7	3.0	0.25	High Sensitivity to Potential Gain	2.00		31	
2	516_Derived_Mo derate	24.7	2.8	0.25	High Sensitivity to Potential Gain	2.00		34	

Assessment Id Proposal Name Page 1 of 5

^{*} Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.



BAM Credit Summary Report

3	516_Disturbed_Gr assland	15.8	7.4	0.25	High Sensitivity to Potential Gain	2.00		0
							Subtotal	65
_	barked Apple - White C nd Bioregion	ypress Pine - B	lakely's Red	d Gum ri	parian open forest / woodland of the Na	andewar Bioreg	ion and New Engla	and
12	544_Low	35.0	0.8	0.25	High Sensitivity to Potential Gain	2.00	TRUE	15
							Subtotal	15
Silver-le	eaved Ironbark - White	Cypress Pine s	hrubby ope	n forest	of Brigalow Belt South Bioregion and N	andewar Bioreg	jion	
4	594_Moderate	61.3	4.1	0.25	High Sensitivity to Potential Gain	1.75		110
5	594_Low	33.3	9.5	0.25	High Sensitivity to Potential Gain	1.75		139
6	594_Disturbed_Gr assland	11.8	49.8	0.25	High Sensitivity to Potential Gain	1.75		0
							Subtotal	249
Tumble	down Red Gum - White	Cypress Pine	Silver-leav	ed Ironb	park shrubby woodland mainly in the no	rthern Nandew	ar Bioregion	
7	596_Moderate	62.6	11.2	0.25	High Sensitivity to Potential Gain	1.50		262
8	596_Low	38.2	0.3	0.25	High Sensitivity to Potential Gain	1.50		4
9	596_Very_Low	23.6	0.3	0.25	High Sensitivity to Potential Gain	1.50		3
10	596_Derived_Low	8.0	9.0	0.25	High Sensitivity to Potential Gain	1.50		0



BAM Credit Summary Report

6_Disturbed_Gr 5.6 sland	50.6	0.25	High Sensitivity to Potential Gain	1.50		0
					Subtotal	269
					Total	598

Species credits for threatened species

Vegetation zone name	Habitat condition (HC)	Area (ha) / individual (HL)	Constant	Biodiversity risk weighting	Potential SAII	Species credits
Ninox connivens / Bar	king Owl (Fauna)					
516_Very_Low	20.7	3	0.25	2	False	31
594_Moderate	61.3	4.1	0.25	2	False	126
594_Low	33.3	9.54	0.25	2	False	159
596_Moderate	62.6	11.15	0.25	2	False	349
544_Low	35.0	0.83	0.25	2	False	15
					Subtotal	680
Setirostris eleryi / Bris	tle-faced Free-tailed Bat (I	auna)				
516_Very_Low	20.7	3	0.25	2	False	31
516_Disturbed_Grasslard	15.8	7.39	0.25	2	False	58
594_Moderate	61.3	4.1	0.25	2	False	126
594_Low	33.3	9.54	0.25	2	False	159



BAM Credit Summary Report

594_Disturbed_Grasslan d	11.8	49.81	0.25	2	False	293
596_Moderate	62.6	11.15	0.25	2	False	349
596_Low	38.2	0.25	0.25	2	False	5
596_Very_Low	23.6	0.3	0.25	2	False	4
596_Disturbed_Grasslan	5.6	50.34	0.25	2	False	140
544_Low	35.0	0.83	0.25	2	False	15
					Subtotal	1180
Tyto novaehollandiae /	' Masked Owl (Fauna)					
516_Very_Low	20.7	3	0.25	2	False	31
594_Moderate	61.3	4.1	0.25	2	False	126
594_Low	33.3	9.54	0.25	2	False	159
596_Moderate	62.6	11.15	0.25	2	False	349
544_Low	35.0	0.83	0.25	2	False	15
					Subtotal	680
Vespadelus troughtoni	/ Eastern Cave Bat (Fauna)				
516_Very_Low	20.7	3	0.25	3	True	46
594_Moderate	61.3	4.1	0.25	3	True	189
594_Low	33.3	9.54	0.25	3	True	238
596_Moderate	62.6	11.15	0.25	3	True	523

Assessment Id 00015157/BAAS18113/19/00015159 Proposal Name

Page 4 of 5



BAM Credit Summary Report

596_Low	38.2	0.25	0.25	3	True	7
596_Very_Low	23.6	0.3	0.25	3	True	5
544_Low	35.0	0.83	0.25	3	True	22
					Subtotal	1030

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development A	Assessment Report	
APPENDIX K	BAM CALCULATOR CREDIT REPORT	

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020



Proposal Details

Assessment Id	Proposal Name	BAM data last updated *
00015157/BAAS18113/19/00015159	Bonshaw Solar Farm	26/11/2019
Assessor Name Matt Jenkins	Assessor Number BAAS18029	BAM Data version * 22
Proponent Names	Report Created 17/03/2020	BAM Case Status
GAIA Australia Assessment Revision	Assessment Type	Open Date Finalised
1	Part 4 Developments (General)	To be finalised

Potential Serious and Irreversible Impacts

* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

Name of threatened ecological community	Listing status	Name of Plant Community Type/ID
White Box Yellow Box Blakely's Red Gum Woodland		544-Rough-barked Apple - White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion

Species	
Vespadelus troughtoni / Eastern Cave Bat	
Vespadelus troughtoni / Eastern Cave Bat	
Vespadelus troughtoni / Eastern Cave Bat	



Vespadelus troughtoni / Eastern Cave Bat

Additional Information for Approval

PCTs With Customized Benchmarks
No Changes

Predicted Threatened Species Not On Site

Name

Calyptorhynchus lathami / Glossy Black-Cockatoo

Haliaeetus leucogaster / White-bellied Sea-Eagle

Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	Number of credits to be retired
516-Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	Not a TEC	13.2	65.00
594-Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	Not a TEC	63.5	249.00



596-Tumbledown Red Gum - White Cypress Pine - Silver- leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	Not a TEC	71.1	269.00
544-Rough-barked Apple - White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion	White Box Yellow Box Blakely's Red Gum Woodland	0.8	15.00

516-Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion

ke-for-like credit retirement options			
Class	Trading group	НВТ	IBRA region
Western Slopes Grassy Woodlands This includes PCT's: 201, 202, 266, 267, 274, 275, 276, 277, 278, 280, 282, 283, 286, 301, 337, 383, 426, 433, 437, 441, 444, 483, 509, 516, 589, 590, 593, 599, 847, 955, 1303, 1304, 1315, 1329, 1383, 1695	Western Slopes Grassy Woodlands - ≥ 70% - <90% cleared group (including Tier 4 or higher).	Yes	Nandewar Northern Complex, Binghi Plateau, Inverell Basalts, Northern Basalts, Severn River Volcanics, Stanthorpe Plateau and Tenterfield Plateau. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.



544-Rough-barked Apple -White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion

	Like-for-like credit retirement options					
's st	Name of offset trading group	Trading group	НВТ	IBRA region		
r	White Box Yellow Box Blakely's Red Gum Woodland This includes PCT's: 2, 74, 75, 83, 250, 266, 267, 268, 270, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 286, 298, 302, 312, 341, 342, 347, 350, 352, 356, 367, 381, 382, 395, 403, 421, 433, 434, 435, 436, 437, 451, 483, 484, 488, 492, 496, 506, 508, 509, 510, 511, 528, 538, 544, 563, 567, 571, 589, 590, 597, 599, 618, 619, 622, 633, 654, 702, 703, 704, 705, 710, 711, 796, 797, 799, 840, 847, 851, 921, 1099, 1103, 1303, 1304, 1307, 1324, 1329, 1330, 1331, 1332, 1333, 1334, 1383, 1401, 1512, 1601, 1606, 1608, 1611, 1691, 1693, 1695, 1698	_	Yes	Nandewar Northern Complex, Binghi Plateau, Inverell Basalts, Northern Basalts, Severn River Volcanics, Stanthorpe Plateau and Tenterfield Plateau. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.		



594-Silver-leaved Ironbark -
White Cypress Pine shrubby
open forest of Brigalow Belt
South Bioregion and
Nandewar Bioregion

Like-for-like credit retirement options					
Class	Trading group	НВТ	IBRA region		
North-west Slopes Dry Sclerophyll Woodlands This includes PCT's: 228, 429, 435, 517, 527, 529, 564, 588, 594, 595, 597, 598, 856, 1165, 1306, 1308, 1317, 1387, 1586, 1607	North-west Slopes Dry Sclerophyll Woodlands - ≥ 50% - < 70% cleared group (including Tier 6 or higher).	Yes	Nandewar Northern Complex, Binghi Plateau, Inverell Basalts, Northern Basalts, Severn River Volcanics, Stanthorpe Plateau and Tenterfield Plateau. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.		

596-Tumbledown Red Gum -White Cypress Pine - Silverleaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion

Like-for-like credit retirement options

Class Trading group HBT IBRA region	
-------------------------------------	--



North-west Slopes Dry Sclerophyll	North-west Slopes Dry	Yes	Nandewar Northern Complex, Binghi
Woodlands	Sclerophyll Woodlands - <		Plateau, Inverell Basalts, Northern
This includes PCT's:	50% cleared group (including		Basalts, Severn River Volcanics,
228, 380, 381, 382, 384, 385, 386, 389,	Tier 7 or higher).		Stanthorpe Plateau and Tenterfield
390, 391, 393, 394, 412, 413, 418, 429,			Plateau.
432, 435, 453, 506, 517, 527, 529, 543,			or
549, 555, 562, 563, 564, 573, 587, 588,			Any IBRA subregion that is within 100
591, 594, 595, 596, 597, 598, 856, 1165,			kilometers of the outer edge of the
1306, 1308, 1317, 1387, 1560, 1586,			impacted site.
1587, 1605, 1606, 1607, 1611, 1613			·

Species Credit Summary

Species	Area	Credits
Ninox connivens / Barking Owl	28.6	680.00
Setirostris eleryi / Bristle-faced Free-tailed Bat	136.7	1180.00
Tyto novaehollandiae / Masked Owl	28.6	680.00
Vespadelus troughtoni / Eastern Cave Bat	29.2	1030.00

Ninox connivens/ Barking Owl 516_Very_Low	Like-for-like credit retirement options		
	Spp	IBRA region	



	Ninox connivens/Barking Owl	Any in NSW
544_Low	Like-for-like credit retirement options	
	Spp	IBRA region
	Ninox connivens/Barking Owl	Any in NSW
594_Low	Like-for-like credit retirement options	
594_LOW	Like for the create retirement options	
	Spp	IBRA region
	Spp Ninox connivens/Barking Owl	IBRA region Any in NSW
	Spp Ninox connivens/Barking Owl	Any in NSW
594_Moderate		
594_Moderate	Ninox connivens/Barking Owl	



	596_Moderate	Like-for-like credit retirement options	
		Spp	IBRA region
		Ninox connivens/Barking Owl	Any in NSW
Setirostris eleryi/	516_Disturbed_Grass	Like-for-like credit retirement options	
Bristle-faced Free-tailed Bat	land	Spp	IBRA region
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
	516_Very_Low	Like-for-like credit retirement options	
		Spp	IBRA region
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
	544_Low	Like-for-like credit retirement options	
		Spp	IBRA region

Assessment Id 00015157/BAAS18113/19/00015159

Proposal Name

Page 8 of 14



	Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
	Like-for-like credit retirement options	
land	Spp	IBRA region
	Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
594_Low	Like-for-like credit retirement options Spp Setirostris eleryi/Bristle-faced Free-tailed Bat	IBRA region Any in NSW
594_Moderate	Like-for-like credit retirement options	
594_Moderate	Like-for-like credit retirement options Spp	IBRA region



596_Disturbed_Grass	Like-for-like credit retirement options	
	Spp	IBRA region
	Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
596_Low	Like-for-like credit retirement options	
	Spp	IBRA region
	Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
596_Moderate	Like-for-like credit retirement options	
	Spp	IBRA region
	Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
596_Very_Low	Like-for-like credit retirement options	



		Spp	IBRA region
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW
Tyto novaehollandiae/	516_Very_Low	Like-for-like credit retirement options	
Masked Owl		Spp	IBRA region
		Tyto novaehollandiae/Masked Owl	Any in NSW
	544_Low	Like-for-like credit retirement options	
		Spp	IBRA region
		Tyto novaehollandiae/Masked Owl	Any in NSW
	594_Low	Like-for-like credit retirement options	
		Spp	IBRA region
		Tyto novaehollandiae/Masked Owl	Any in NSW



Tyto novaehollandiae/ Masked Owl	594_Low				
	594_Moderate	rate Like-for-like credit retirement options			
		Spp	IBRA region		
		Tyto novaehollandiae/Masked Owl	Any in NSW		
	596_Moderate	Like-for-like credit retirement options			
		Spp	IBRA region		
		Tyto novaehollandiae/Masked Owl	Any in NSW		
Vespadelus troughtoni/	516_Very_Low	Like-for-like credit retirement options			
Eastern Cave Bat		Spp	IBRA region		
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW		



Vespadelus troughtoni/	544_Low	Like-for-like credit retirement options	
Eastern Cave Bat		Spp	IBRA region
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW
	594_Low	Like-for-like credit retirement options	
		Spp	IBRA region
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW
	594_Moderate	Like-for-like credit retirement options	
		Spp	IBRA region
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW
	596_Low	Like-for-like credit retirement options	
		Spp	IBRA region



596_Moderate Like-for-like credit retirement options Spp IBRA regio	
Spp IBRA regio	
	ion
Vespadelus troughtoni/Eastern Cave Bat Any in NSV	SW
596_Very_Low Like-for-like credit retirement options	
Spp IBRA regio	ion
Vespadelus troughtoni/Eastern Cave Bat Any in NSV	SW

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development Assessment Report

APPENDIX L PERSONNEL

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

Personnel involved in preparation of this BDAR and their contribution is summarised in Table L1 below.

Table L.1 Summary of Personnel and Duties

Name	Position	Tasks and Responsibilities
Joanne Woodhouse	Principal Ecologist and Project Manager	 Project Management including liaison with client, BCD and other stakeholders Technical review Bird survey and BAM plot support (Spring 2018)
Matt Jenkins North Coast Ecology	Project Ecologist Accredited BAM Assessor	 Field ecologist: Summer 2020 survey Additional field work design and BAM plots Report writing and BAM Calculator report to support Response to Submission (RTS) and revised BDAR
Dr Adriana Corona Mothe	Field Ecologist Accredited BAM Assessor	 Field ecologist: fauna survey and BAM plots (2018-2019 survey periods) Field work design and BAM plots Report writing and BAM Calculator reports to support EIS submission
Tom Cotter	Biodiversity and GIS Consultant	 Field Ecologist: fauna and BAM plots (summer 2018). Vegetation mapping, species polygon Draft figures
Dr Toivo Zoete	Senior Environmental Consultant	Flora assessment in the 20m x 20m sub-plot of BAM plots (Autumn 2019).
Georgina Race	GIS Consultant	Vegetation mapping and area calculationsGIS Support
Viet Nguyen	GIS Consultant	GIS SupportVegetation area calculationsFigure preparation
Danielle Robinson	GIS Consultant	GIS SupportField figure preparationField map layers preparation

GAIA BONSHAW SOLAR EIS Revised Biodiversity Development	Assessment Report
APPENDIX M	UPDATED SPECIES POLYGONS AND CREDIT REPORTS JUNE 2020

www.erm.com Version: 2.0 Project No.: 0470861 Client: GAIA Australia 18 May 2020

The following updated species polygons and credit report have been provided as requested by the Department of Planning, Industry and Environment (DPIE) in response the Response to Submissions (RTS) provided in May 2020.

Provide an addendum to the BDAR, addressing:

- The species polygons for the Masked Owl and Barking Owl must be expanded to include each of the recorded hollow bearing trees (green dots) within the development site as identified in figure 5.3 of the BDAR, because hollow bearing trees are an important part of the breeding and foraging habitat for these owls. The expanded species polygons do not need to include extensive areas of disturbed grassland.
- The Bristle-faced Free-tailed bat species polygon currently covers the entire development site, including the mapped disturbed grassland areas. We note the precautionary approach of including large areas of disturbed grassland for potential foraging habitat. However, due to the large area of disturbed grassland, we would also accept a reduced species polygon, that aims to capture the more viable bat habitat components, such as hollow bearing trees, similar to that described above for the owls.

Based on these amendments as detailed below, the proposed Bonshaw Solar Farm development in its current footprint requires a total offset of 598 ecosystem credits and 2,244 species credits.

Retirement of these credits will be carried out in accordance with the Biodiversity Conservation Regulation 2017 and the NSW Biodiversity Offsets Scheme.

Bristle-faced Free-tailed Bat

In accordance with Table 1 of the OEH (2018b) 'Species Credit' threatened bats and their habitats guide, and based on advice from DPIE, the revised species polygon for the Bristle-faced Free-tailed Bat has been prepared and includes the following features:

- All woodland PCT on the Subject Land where the Subject Land is within 500m of a river, creek or riparian area; and
- A 50m buffer to all hollow-bearing trees recorded within the Subject Land.

Disturbed grasslands have been excluded from the updated species polygons. The updated species polygon for the Bristle-faced Free-tailed Bat is provided in **Figure L-1**.

No amendments have been made to the Eastern Cave Bat species polygon (refer to Figure 5.3 in the main document).

Barking Owl

The TBDC provides that where a breeding site has been identified in accordance with the BAM the species polygon should be established by providing a circular buffer with a 100m radius around the nest tree. This includes suitable habitat for male roosts, feeding/grooming perches and fledgling requirements. It does not account for foraging habitat. The shape of the buffer can be modified where evidence indicates an alternative shape would better meet the species needs in the context of the assessment site.

As no targeted surveys (e.g. call playback and identification of nest trees) were completed for this dual credit species, breeding habitat has been assumed to be present by identifying suitable habitat. As requested by DPIE, a species polygon has been mapped for this species by including all mapped hollow-bearing trees that contained hollows of a suitable size (i.e. >20cm wide) and height above the ground (i.e. >4m) in accordance with breeding habitat constraints provided in BAM-C. The updated species polygon for the Barking Owl is provided in **Figure L-2**.

As a dual credit species, potential foraging habitat is addressed within the ecosystem credit requirement for PCTs associated with this species. Conservation management of these PCTs will

increase the quality and suitability of forage habitat for this species through improvement of structure and function components and likely associated increases in prey items (i.e. from habitat enhancement and feral predator control). Therefore, the species polygons have been revised to include potential breeding sites only as required in the BAM.

Given the approach described above, the species polygon for Barking Owl is considered to represent a conservative overestimate of actual/potential breeding habitat in that all hollow-bearing trees of a suitable size (i.e. >20cm wide) have been included rather than a single confirmed nest tree.

Masked Owl

The TBDC provides that where a breeding site has been identified in accordance with the BAM the species polygon should be established by providing a circular buffer with a 100m radius around the nest tree. This includes suitable habitat for male roosts, feeding/grooming perches and fledgling requirements. It does not account for foraging habitat. The shape of the buffer can be modified where evidence indicates an alternative shape would better meet the species needs in the context of the assessment site.

As no targeted surveys (e.g. call playback and identification of nest trees) were completed for this dual credit species, breeding habitat has been assumed to be present by identifying suitable habitat. Although no breeding habitat constraints are explicitly identified in BAM-C for this species, the Approved Recovery Plan for Large Forest Owls (DEC 2006) states that the Masked Owl nests in hollows greater than 40cm wide and greater than 100cm deep, with hollow entrances at least 3m above ground, in trees of at least 90cm diameter at breast height. The TBDC also provides that large tree hollows used.

As requested by DPIE, a species polygon has been mapped for this species by including all mapped hollow-bearing trees that contained hollows of a suitable size (i.e. >40cm wide). The updated species polygon for the Masked Owl is provided in **Figure L-3**.

As a dual credit species, potential foraging habitat is addressed within the ecosystem credit requirement for PCTs associated with this species. Conservation management of these PCTs will increase the quality and suitability of forage habitat for this species through improvement of structure and function components and likely associated increases in prey items (i.e. from habitat enhancement and feral predator control). Therefore, the species polygons have been revised to include potential breeding sites only as required in the BAM.

Given the approach described above, the species polygon for Masked Owl is considered to represent a conservative overestimate of actual/potential breeding habitat in that all hollow-bearing trees of a suitable size (i.e. >40cm wide) have been included and particularly given that this species is generally known to roost and breed in more forested habitats in NSW. The Approved Recovery Plan for Large Forest Owls (DEC 2006) also provides that the Masked Owl prefers unlogged or lightly logged forest with high densities of old hollow trees and avoids young regrowth.


Revised Credit Obligations


Based on the updated mapping of the threatened species polygons, impacts to habitat of Threatened Species requiring offsets now include:


- Habitat for the Bristle-faced Free-tailed Bat requiring 827 species credits;
- Foraging habitat for Eastern Cave Bat requiring 1,030 species credits;
- Potential (assumed) breeding habitat for Masked requiring 104 species credits; and
- Potential (assumed) breeding habitat for Barking Owl requiring 283 species credits.

A total of 598 ecosystem credits are also required to offset the impacts of the project.

The proposed Bonshaw Solar Farm development in its current footprint requires a total offset of 598 ecosystem credits and 2,244 species credits. Retirement of these credits will be carried out in accordance with the Biodiversity Conservation Regulation 2017 and the NSW Biodiversity Offsets Scheme.









Proposal Details

Assessment Id	Proposal Name	BAM data last updated *
00015157/BAAS17014/19/00015159	Bonshaw Solar Farm	04/06/2020
Assessor Name Matt Jenkins	Assessor Number BAAS18029	BAM Data version * 27
Proponent Names GAIA Australia, GAIA Australia Pty Ltd	Report Created 17/06/2020	BAM Case Status Finalised
Assessment Revision 1	Assessment Type Part 4 Developments (General)	Date Finalised 17/06/2020

Potential Serious and Irreversible Impacts

* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

Woodland Community forest / woodland of the Nandewar Bioregion and New England Tableland	Name of threatened ecological community	Listing status	Name of Plant Community Type/ID	
Bioregion	,	Community	544-Rough-barked Apple - White Cypress Pine - Blakely's Red Gum riparian oper forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion	1

Species	
Vespadelus troughtoni / Eastern Cave Bat	
Vespadelus troughtoni / Eastern Cave Bat	
Vespadelus troughtoni / Eastern Cave Bat	



Vespadelus troughtoni / Eastern Cave Bat

Additional Information for Approval

PCTs With Customized Benchmarks
No Changes

Predicted Threatened Species Not On Site

Name

Calyptorhynchus lathami / Glossy Black-Cockatoo

Haliaeetus leucogaster / White-bellied Sea-Eagle

Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	Number of credits to be retired
516-Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	Not a TEC	13.2	65.00
594-Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	Not a TEC	63.5	249.00



596-Tumbledown Red Gum - White Cypress Pine - Silver- leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	Not a TEC	71.3	269.00
544-Rough-barked Apple - White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion	White Box Yellow Box Blakely's Red Gum Woodland	0.8	15.00

516-Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion

Like-for-like credit retirement options			
Class	Trading group	НВТ	IBRA region
Western Slopes Grassy Woodlands This includes PCT's: 201, 202, 266, 267, 274, 275, 276, 277, 278, 280, 282, 283, 286, 301, 337, 383, 426, 433, 437, 441, 444, 483, 509, 516, 589, 590, 593, 599, 847, 955, 1303, 1304, 1315, 1329, 1383, 1695	Western Slopes Grassy Woodlands >=70% and <90%	Yes	Nandewar Northern Complex, Binghi Plateau, Inverell Basalts, Northern Basalts, Severn River Volcanics, Stanthorpe Plateau and Tenterfield Plateau. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.



544-Rough-barked Apple -White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion

	Like-for-like credit retirement options			
's st	Name of offset trading group	Trading group	НВТ	IBRA region
r	White Box Yellow Box Blakely's Red Gum Woodland This includes PCT's: 2, 74, 75, 83, 250, 266, 267, 268, 270, 274, 275, 276, 277, 278, 279, 280, 281, 282, 283, 284, 286, 298, 302, 312, 341, 342, 347, 350, 352, 356, 367, 381, 382, 395, 403, 421, 433, 434, 435, 436, 437, 451, 483, 484, 488, 492, 496, 506, 508, 509, 510, 511, 528, 538, 544, 563, 567, 571, 589, 590, 597, 599, 618, 619, 622, 633, 654, 702, 703, 704, 705, 710, 711, 796, 797, 799, 840, 847, 851, 921, 1099, 1103, 1303, 1304, 1307, 1324, 1329, 1330, 1331, 1332, 1333, 1334, 1383, 1401, 1512, 1601, 1606, 1608, 1611, 1691, 1693, 1695, 1698		Yes	Nandewar Northern Complex, Binghi Plateau, Inverell Basalts, Northern Basalts, Severn River Volcanics, Stanthorpe Plateau and Tenterfield Plateau. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.

Bonshaw Solar Farm



Like-for-like credit retirement options					
Class	Trading group	НВТ	IBRA region		
North-west Slopes Dry Sclerophyll Woodlands This includes PCT's: 228, 429, 435, 517, 527, 529, 564, 588, 594, 595, 597, 598, 856, 1165, 1306, 1308, 1317, 1387, 1586, 1607	North-west Slopes Dry Sclerophyll Woodlands >=50% and <70%	Yes	Nandewar Northern Complex, Binghi Plateau, Inverell Basalts, Northern Basalts, Severn River Volcanics, Stanthorpe Plateau and Tenterfield Plateau. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.		

596-Tumbledown Red Gum -White Cypress Pine - Silverleaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion

Like-for-like credit retirement options



North-west Slopes Dry Scleroph	yll North-west Slopes Dry	Yes	Nandewar Northern Complex, Binghi
Woodlands	Sclerophyll Woodlands <50%		Plateau, Inverell Basalts, Northern
This includes PCT's:			Basalts, Severn River Volcanics,
228, 380, 381, 382, 384, 385, 386	5, 389,		Stanthorpe Plateau and Tenterfield
390, 391, 393, 394, 412, 413, 418	3, 429,		Plateau.
432, 435, 453, 506, 517, 527, 529), 543,		or
549, 555, 562, 563, 564, 573, 587	7, 588,		Any IBRA subregion that is within 100
591, 594, 595, 596, 597, 598, 856	5, 1165,		kilometers of the outer edge of the
1306, 1308, 1317, 1387, 1560, 15	86,		impacted site.
1587, 1605, 1606, 1607, 1611, 16	513		·

Species Credit Summary

Species	Area	Credits
Ninox connivens / Barking Owl	32.9	283.00
Setirostris eleryi / Bristle-faced Free-tailed Bat	56.2	827.00
Tyto novaehollandiae / Masked Owl	7.1	104.00
Vespadelus troughtoni / Eastern Cave Bat	29.2	1030.00

	516_Derived_Moder	Like-for-like credit retirement options		
Barking Owl	ate	Spp	IBRA region	



	Ninox connivens/Barking Owl	Any in NSW
F1C Distructs and Cura		
land	Spp Like-for-like credit retirement options	IBRA region
	Ninox connivens/Barking Owl	Any in NSW
516_Very_Low	Like-for-like credit retirement options	
	Spp	IBRA region
	Ninox connivens/Barking Owl	Any in NSW
544_Low	Like-for-like credit retirement options	
544_Low	Like-for-like credit retirement options Spp	IBRA region



594_Disturbed_Grass land	Like-for-like credit retirement options	
	Spp	IBRA region
	Ninox connivens/Barking Owl	Any in NSW
594_Low	Like-for-like credit retirement options	
	Spp	IBRA region
	Ninox connivens/Barking Owl	Any in NSW
594_Moderate	Like-for-like credit retirement options	
	Spp	IBRA region
	Ninox connivens/Barking Owl	Any in NSW



	Spp	IBRA region	
	Ninox connivens/Barking Owl	Any in NSW	
596_Disturbed_Gra	ass Like-for-like credit retirement options	Like-for-like credit retirement options	
land	Spp	IBRA region	
	Ninox connivens/Barking Owl	Any in NSW	
596 Moderate	Like-for-like credit retirement options		
596_Moderate	Like-for-like credit retirement options Spp	IBRA region	
596_Moderate	Like-for-like credit retirement options Spp Ninox connivens/Barking Owl	IBRA region Any in NSW	
596_Moderate	Spp	-	
596_Moderate 596_Very_Low	Spp	-	
	Spp Ninox connivens/Barking Owl	-	



Setirostris eleryi/ Bristle-faced Free-tailed Bat	516_Derived_Moder ate	Like-for-like credit retirement options		
		Spp	IBRA region	
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW	
			·	
	516_Disturbed_Grass land	Like-for-like credit retirement options		
		Spp	IBRA region	
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW	
516_Very_Low				
	Like-for-like credit retirement options			
		Spp	IBRA region	
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW	



Setirostris eleryi/ Bristle-faced Free-tailed	516_Very_Low						
Bat	544_Low	Like-for-like credit retirement options					
		Spp	IBRA region				
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW				
		Like-for-like credit retirement options					
	land	Spp	IBRA region				
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW				
	594 Low	Like for like gradit ratirement entions					
	594_Low	Like-for-like credit retirement options	IDDA vanian				
		Spp	IBRA region				
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW				
	594_Moderate	Like-for-like credit retirement options					



Setirostris eleryi/Bristle-faced Free-tailed Bat Any in NSW
596_Derived_Low Like-for-like credit retirement options
Spp IBRA region
Setirostris eleryi/Bristle-faced Free-tailed Bat Any in NSW
·
596_Disturbed_Grass land Like-for-like credit retirement options IBRA region
Setirostris eleryi/Bristle-faced Free-tailed Bat Any in NSW
596 Low Like-for-like credit retirement entions
596_Low Like-for-like credit retirement options Spp IBRA region



	596_Moderate	Like-for-like credit retirement options						
		Spp	IBRA region					
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW					
	596_Very_Low							
		Like-for-like credit retirement options						
		Spp	IBRA region					
		Setirostris eleryi/Bristle-faced Free-tailed Bat	Any in NSW					
Tyto novaehollandiae/	516_Derived_Moder	Like-for-like credit retirement options						
Masked Owl	ate	Spp	IBRA region					
		Tyto novaehollandiae/Masked Owl	Any in NSW					



Tyto novaehollandiae/ Masked Owl	516_Disturbed_Grass land	Like-for-like credit retirement options					
		Spp	IBRA region				
		Tyto novaehollandiae/Masked Owl	Any in NSW				
	516_Very_Low	Like-for-like credit retirement options					
		Spp	IBRA region				
		Tyto novaehollandiae/Masked Owl	Any in NSW				
	594_Disturbed_Grass land	Like-for-like credit retirement options					
		Spp	IBRA region				
		Tyto novaehollandiae/Masked Owl	Any in NSW				
	594_Low	Like-for-like credit retirement options					
		Spp	IBRA region				

Assessment Id 00015157/BAAS17014/19/00015159

Proposal Name

Page 14 of 18



	Tyto novaehollandiae/Masked Owl	Any in NSW
596_Derived_Low	Like-for-like credit retirement options	
	Spp	IBRA region
	Tyto novaehollandiae/Masked Owl	Any in NSW
596_Disturbed_Grassland	S Like-for-like credit retirement options Spp Tyto novaehollandiae/Masked Owl	IBRA region Any in NSW
596_Moderate	Like-for-like credit retirement options	
596_Moderate	Like-for-like credit retirement options Spp	IBRA region



Vespadelus troughtoni/ Eastern Cave Bat	516_Very_Low	Like-for-like credit retirement options					
		Spp	IBRA region				
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW				
	544_Low	Like-for-like credit retirement options					
		Spp	IBRA region				
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW				
	594_Low	Like-for-like credit retirement options					
		Spp	IBRA region				
		Vespadelus troughtoni/Eastern Cave Bat	Any in NSW				
	594_Moderate	Like-for-like credit retirement options					



	Spp	IBRA region
	Vespadelus troughtoni/Eastern Cave Bat	Any in NSW
596_Low	Like-for-like credit retirement options	
	Spp	IBRA region
	Vespadelus troughtoni/Eastern Cave Bat	Any in NSW
596 Moderate	Like-for-like credit retirement options	
596_Moderate	Like-for-like credit retirement options	IRDA ragion
596_Moderate	Spp	IBRA region
596_Moderate		IBRA region Any in NSW
596_Moderate	Spp	
596_Moderate	Spp	
	Vespadelus troughtoni/Eastern Cave Bat	





Assessment Id Payment data version Assessment Revision Report created

00015157/BAAS17014/19/000151 63 1 17/06/2020

59

Assessor Name Assessor Number Proposal Name BAM Case Status

Matt Jenkins BAAS18029 Bonshaw Solar Farm Finalised

Assessment Type Date Finalised

PCT list Part 4 Developments (General) 17/06/2020

Price calculated	PCT common name	Credits
Yes	516 - Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	65
Yes	594 - Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	249
Yes	596 - Tumbledown Red Gum - White Cypress Pine - Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	269
Yes	544 - Rough-barked Apple - White Cypress Pine - Blakely's Red Gum riparian open forest / woodland of the Nandewar Bioregion and New England Tableland Bioregion	15

Species list

Price calculated	Species	Credits
Yes	Ninox connivens (Barking Owl)	283
Yes	Tyto novaehollandiae (Masked Owl)	104
Yes	Vespadelus troughtoni (Eastern Cave Bat)	1030
Yes	Setirostris eleryi (Bristle-faced Free-tailed Bat)	827

Assessment Id Proposal Name Page 1 of 8



Assessment Id Proposal Name Page 2 of 8



Ecosystem credits for plant communities types (PCT), ecological communities & threatened species habitat

IBRA sub region	PCT common name	Threat status	Offset trading group	Risk premiu m	Administ rative cost	Methodology adjustment factor	Price per credit	No. of ecosystem credits	Final credits price
Nandewar Northern Complex	516 - Grey Box grassy woodland or open forest of the Nandewar Bioregion and New England Tableland Bioregion	No	Western Slopes Grassy Woodlands >=70% and <90%	34.31%	\$800.02	2.0046	\$ 27,662.63	65	\$ 1,798,070.77
Nandewar Northern Complex	594 - Silver-leaved Ironbark - White Cypress Pine shrubby open forest of Brigalow Belt South Bioregion and Nandewar Bioregion	No	North-west Slopes Dry Sclerophyll Woodlands >=50% and <70%	34.31%	\$80.00	2.3073	\$1,694.69	249	\$421,978.15
Nandewar Northern Complex	596 - Tumbledown Red Gum - White Cypress Pine - Silver-leaved Ironbark shrubby woodland mainly in the northern Nandewar Bioregion	No	North-west Slopes Dry Sclerophyll Woodlands <50%	23.65%	\$80.00	2.0204	\$1,698.34	269	\$456,852.33

Assessment Id Proposal Name Page 3 of 8



Assessment Id Proposal Name Page 4 of 8



Nandewar	544 - Rough-barked Apple - White	No	Western Slopes	34.31%	\$80.00	2.3617	\$907.80	15	\$13,617.00
Northern	Cypress Pine - Blakely's Red Gum		Grassy						
Complex	riparian open forest / woodland of the		Woodlands						
	Nandewar Bioregion and New England		>=50% and						
	Tableland Bioregion		<70%						

Subtotal (excl. GST) **\$2,690,518.25**

GST **\$269,051.82**

Total ecosystem credits (incl. GST) \$2,959,570.08

Species credits for threatened species

Species profile ID	Species	Threat status	Price per credit	Risk premium	Administrative cost	No. of species credits	Final credits price
10561	Ninox connivens (Barking Owl)		\$173.02	34.3100%	\$80.00	283	\$88,404.43
10820	Tyto novaehollandiae (Masked Owl)		\$447.26	34.3100%	\$80.00	104	\$70,794.35
10829	Vespadelus troughtoni (Eastern Cave Bat)		\$741.31	34.3100%	\$80.00	1030	\$1,107,923.06
10923	Setirostris eleryi (Bristle-faced Freetailed Bat)		\$636.69	34.3100%	\$80.00	827	\$773,359.41

Subtotal (excl. GST)

\$2,040,481.25



Assessment Id Proposal Name Page 6 of 8



510(0)(0)(0)(0)(0)(0)(0)		
	GST	\$204,048.12
Total species credits (inc	. GST)	\$2,244,529.38
	Grand total	\$5,204,099,46



Assessment Id Proposal Name Page 8 of 8

ERM has over 160 offices across the following countries and territories worldwide

Argentina The Netherlands Australia New Zealand Belgium Norway Brazil Panama Canada Peru Chile Poland China Portugal Colombia Puerto Rico France Romania Germany Russia Hong Kong Singapore India South Africa Indonesia South Korea Ireland Spain Italy Sweden Japan Switzerland Taiwan Kazakhstan Kenya Thailand Malaysia UAE Mexico UK Mozambique US Myanmar Vietnam

ERM's Newcastle Office

Level 1 45 Watt Street Newcastle NSW 2300

PO Box 803

Newcastle NSW 2300

T: +61 2 4903 5500

www.erm.com



ERM	Bonshaw Solar Farm (SSD 9438) Response to RFI – May 2020 Page 18 of 8
ATTACHMENT D	EVIDENCE OF WATER SUPPLIER ABILITY

From: Trident Water tridentwaterrad@gmail.com &

Subject: Re: Regarding drinking water delivery for 200 construction workers

Date: 20 May 2020 at 11:46 am

To: Shawn HawnKi LEE shawn.lee@gaiaau.com

Good Morning Shawn,

Thank you very much for your email.

Please excuse the delay in getting back to you. Attached is a presentation of Trident Waters capability to service the Bonshaw Energy Park.

Trident Water is approved to operate by NSW Health. All our trucks adhere to strict guidelines regarding quality assurance for water transport and storage. With the advent of Covid-19 this now has become more relevant than ever.

Since it inception in 2019 we have successfully and safely delivered over 500 kilolitres to our clients.

We are proud to be council approved locally in the Northern Rivers, comprising of the following shires Inverell, Tenterfield, Richmond Valley and Rous City Council which incorporates three council areas. They are Lismore City, Ballina Shire and the Byron Shire.

Trident Water can offer assurances that no matter what adverse supply or climate conditions are presented. Our supply network is able to meet all your water logistical needs over the duration of both phases of the Bonshaw Solar Farm - Construction and operational.

The most relevant to the Bonshaw Solar farm are the Inverell and Tenterfield Shires. Being approved for both shires we are in a unique position to service the BonShaw Solar Farm within 2 hours from Inverell and 1.5 hours from Tenterfield. As our trucks are fully laden with water these times may vary slightly due to road conditions.

<u>Fleet</u>

Mercedes Unimog 4x4 Heavy duty crane truck. 5 kilolitre tank. Specialist remote access utility vehicle, 1.35 ton crane with a 5.1 metre reach. Water supply, dust suppression, fire mitigation

Hino 500 series: Mine Spec for construction.

15 kilolitre tank. Civil Construction specialist water supply, dust suppression, fire mitigation.

Both trucks are fitted with CB radios and reverse cameras and 100% redundancy via a spare pump should there be any mechanical issues. We can always get your supply to you.

We are currently expanding our fleet to include an 8x8 - 18kl Water truck and a Unimog trailer fitted with a 7kl tank.

Expanding our total capacity to 45 kilolitres in Q4 of this financial year.

I hope this has clarified a few questions as to the capability of Trident Water to supply the Bonshaw Solar Farm.

We are proud to be local and are here to help in any way we can Shawn. It would be an honour to be involved in a project of this magnitude. Please feel free to get in contact.

Best Regards

Roman Guerra Trident Water 0412580564 tridentwaterrad@gmail.com From: Trident Water tridentwaterrad@gmail.com

Subject: Re: Regarding drinking water delivery for 200 construction workers

Date: 23 January 2020 at 9:35 am

To: Shawn HawnKi LEE shawn.lee@gaiaau.com

Good Morning Shawn,

Thank you for your for choosing to getting in contact with a local business regarding Water supply to the proposed GAIA Solar farm site in Bonshaw, NSW.

We would be more than happy to liaise with you regarding this project and provide you with presentation as requested.

In order to do so we will need some more detailed information regarding the camp requirements.

Is the quote just intended just for drinking supply?

Just off the top of my head for a 200 man construction site, consumption rate would be average 2.5 (winter) to 5 (Summer) litres a day \times 200 men=500 - 1000 litres. consumed on site per day.

A year and half program 548 days construction time x 1kl per day = 548kl consumed for the entire project.

This is just a rough estimate of consumption as your construction is due to start in July 2020 (winter) and go through two summer seasons til completion. During the summer season the consumption rates may double and may halve during the winter months

We would need to factor in other variables for your site such as:

Camp kitchen requirements- Catering crew

Steward services for cleaning of offices, amenities, mess, laundry and accommodation blocks.

On site dust suppression for the crew to work safely whist installing the farm- what is the area size of the whole site.

Is it close to any residential areas as this will play a major role for your sites water usage due to council requirements.

Shawn would it be possible for us to schedule a time to discuss your requirements. It would enable us give you the best and most detailed presentation factoring all the above, terms and payment schedules.

We look forward to having a discussion and getting a proposal together for you.

Best Regards

Roman Guerra Trident Water 0412580564

On Wed, 22 Jan 2020 at 11:08, Shawn HawnKi LEE (shawn.lee@gaiaau.com) wrote:

Hi,

This is Shawn Lee from GAIA Australia Pty. Ltd., which is preparing for a Sola Farm business in Bonshaw area. Site location is next to the Transgrid Dumaresq Substation.

It will be 100MW Solar PV electricity generating facility, and now we are expecting to start the construction from coming July this year.

Accordingly, then we need to have enough amount of drinking water for about 200 workers(peak time) under construction.

In average, there will be 150 workers on site during construction. It will take 1.5 years for construction from July, 2020.

Please let us know about your quote and some details, such as payment plan, kinds of services, delivery time and so on, anything we should know before order(or contract). Please contact me via email or phone call for any issues.

Best regards,

ERM	Bonshaw Solar Farm (SSD 9438) Response to RFI – May 2020 Page 19 of 8
'	
ATTACHMENT E	UPDATED ABORIGINAL CULTURAL HERITAGE

ASSESSMENT



GAIA Australia

Bonshaw Solar Farm

Cultural Heritage Assessment

19 June 2020

Project No.: 0464261



Document details	
Document title	Bonshaw Solar Farm
Document subtitle	Cultural Heritage Assessment
Project No.	0464261
Date	19 June 2020
Version	3.0
Author	Katherine Deverson
Client Name	GAIA Australia

Document	history					
				ERM approval to issue		
Version	Revision	Author	Reviewed by	Name	Date	Comments
Draft	00	Katherine Deverson	Erin Finnegan	Paul Douglass	11 April 2019	Approved for issue to Client
Draft	01	Katherine Deverson	Erin Finnegan	Paul Douglass	18 April 2019	Approved for issue to Client
Final	02	Katherine Deverson	Amanda Antcliff	Paul Douglass	12 July 2019	Approved for issue to Client
Final	03	Stephanie Moore	Erin Finnegan	David Dique	13 February 2020	Response to Submission
Final	04	Lachlan Giles	Michael Rookwood	David Dique	19 June 2020	Addressing AHIMS

 www.erm.com
 Version: 3.0
 Project No.: 0464261
 Client: GAIA Australia
 19 June 2020

Signature Page

19 June 2020

Bonshaw Solar Farm

Cultural Heritage Assessment

Stephanie Moore Heritage Consultant Erin Finnegan

Principal Heritage Consultant

Amanda Antcliff Project Manager David Dique Partner

Environmental Resources Management Australia Pty Ltd Level 15 309 Kent Street Sydney, 2000 NSW

© Copyright 2020 by ERM Worldwide Group Ltd and / or its affiliates ("ERM"). All rights reserved. No part of this work may be reproduced or transmitted in any form, or by any means, without the prior written permission of ERM

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia 19 June 2020

CONTENTS

1.	INTR	INTRODUCTION1					
	1.1	Objectiv	Overview Objectives Description of the Proposed Development				
	1.3	•	·				
		1.3.1 1.3.2	Location Proposed Development				
	1.4		ship				
	1.5		Structure				
2.	RELE	EVANT LI	EGISLATION	6			
3.			COMMUNITY CONSULTATION				
0.	3.1		1: Notification of Project Proposal and Registration of Interest				
	3.2	•	Stage 2: Presentation of Information about the Proposed Project				
	3.3	_	Stage 3: Gathering Information about Cultural Significance				
	3.4	Stage 4	1: Review of Draft Cultural Heritage Assessment Report				
4.	ENVI	RONMEN	NTAL BACKGROUND	11			
	4.1	Environ	nmental Context	11			
		4.1.1	The Northern Highlands Bioregion	11			
		4.1.2	Topography and Landforms				
		4.1.3	Flora and Fauna				
		4.1.4	Land Use and Disturbance				
5.	ARC	ARCHAEOLOGICAL BACKGROUND					
	5.1						
	5.2 5.3	•	legional Archaeological Backgroundocal Archaeological Context				
	0.0		5.3.1 Dumaresq to Lismore Transmission Line Aboriginal and Historic Heritage				
		5.5.1	Assessment				
		5.3.2	OEH AHIMS Register				
	5.4	Implicat	tions for the Project Area – Predictive Model	21			
		5.4.1	Predictive Model	22			
6.	HIST	ORICAL	CONTEXT	23			
	6.1	Historic	cal Overview	23			
		6.1.1	Township of Bonshaw	23			
		6.1.2	The Project Area	25			
	6.2	Historic	cal Heritage Database Searches	25			
		6.2.1	Commonwealth Heritage List				
		6.2.2	State Heritage Register				
	6.3	6.2.3	Local Environmental Planal Heritage Predictive Model				
7.			HODOLOGY AND RESULTS				
٠.	7.1						
	7.1	, 3,					
		7.2.1	Description of the Project Area				
		7.2.2	Survey Coverage				
		7.2.3	Survey Results	33			
8.	SIGN	IIFICANC	E ASSESSMENT	53			
	8.1	Assess	ment of Aboriginal Cultural Significance	53			
		8.1.1	Social Significance	54			
		8.1.2	Historic Significance	54			

		8.1.3	Scientific Significance	5/	
		8.1.4	Aesthetic Significance		
	8.2	-	al Heritage Statement of Significance		
	6.2 8.3	-	ent of Historic Heritage Significance		
•					
9.	IMPAG	JI ASSES	SSMENT	65	
10.	CONC	LUSIONS	S AND RECOMMENDATIONS	70	
	10.1	Conclusion	ons	70	
	10.2	Recomm	Recommendations		
		10.2.1	Aboriginal Heritage	70	
		10.2.2	Historic Heritage		
REFE	RENC	ES		78	
	ENDIX A		ORIGINAL STAKEHOLDER CONSULTATION LOG		
	ENDIX E		RVEY METHODOLOGY SENT TO RAPS AUGUST 2018		
APPE	ENDIX (C AH	IMS EXTENSIVE SEARCH RESULTS		
l ist c	of Table) C			
Table			islation	6	
Table		, ,	red Aboriginal Parties (RAPs)		
Table		-	rar Bioregion Attributes		
Table			Heritage Site Types Definitions		
Table			Extensive Search Details		
Table	-		Registered Sites		
Table			ns of Archaeological Potential		
Table			Surface Visibility Rating		
Table			rvey Attendees		
Table			m Summary		
Table			Coverage		
Table			al Cultural Heritage Field Survey Results		
Table	_	•	ry Significance of Aboriginal Heritage Sites		
Table			ry Impact Assessment of Aboriginal Heritage Sites		
Table		, ,			
Table	. 10.1	Manage	Therit weasures for identified Aboriginal Flentage Sites	7 0	
	of Figur				
Figure		-	Area and Location		
Figure		•	Footprint		
Figure	e 5.1	•	Tribal Boundaries in Aboriginal Australia, PA indicated with blue circle (Tin		
Figure	e 5.2	Referen	ce Map from Site Recording Card #11-3-0083 (OzArk 2011)	19	
Figure			Search Results		
Figure	e 6.1	1909 Pa	rish of Bowman map, PA is indicated in red (NSWLRS 1909)	25	
Figure			Units		
Figure		•	lecorded Heritage Sites		
Figure		•	nmatic Presentation of the Significance Assessment Process (OEH 2011).		
Figure		_	e Sites Impacted by Project		

Acronyms and Abbreviations

AHD Australian Height Datum

AHIMS Aboriginal Heritage Information Management System

BP Before Present

BSS Battery Storage System

Burra The Australia ICOMOS Charter for Places of Cultural

Charter Significance

CHA Cultural Heritage Assessment
CHL Commonwealth Heritage List
CIV Capital Investment Value

CSIRO Commonwealth Scientific and Industrial Research

Organisation

DECCW Department of Environment, Climate Change and Water

DP&E Department of Planning and Environment

EIS Environmental Impact Statement

EP&A Act Environmental Planning and Assessment Act 1979 (NSW)
EPBC Act Environment Protection and Biodiversity Conservation Act

1999

ERM Environmental Resources Management Australia Pty Ltd

ESD Ecologically Sustainable Development

GAIA GAIA Australia

IHO Interim Heritage Orders

LALC Local Aboriginal Land Council
LEP Local Environmental Plan
LGA Local Government Area
LLS Local Land Services

MW Megawatts
MWh Megawatt hour

NNTT National Native Title Tribunal

NP&W Act National Parks and Wildlife Act 1974

NPWS NSW National Parks and Wildlife Service

NTS Corp Native Title Services corporation
OEH Office of Environment and Heritage

PA Project Area

PAD Potential Archaeological Deposit

PV Photovoltaic

QA Quality Assurance

RAP Registered Aboriginal Party

SEPP State Environmental Planning Policy

SHI State Heritage Inventory
SHR State Heritage Register

SSD State Significant Development

SU Survey Unit

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia 19 June 2020 Page iii

1. INTRODUCTION

1.1 Overview

ERM was commissioned to undertake a heritage assessment to support an Environmental Impact Statement (EIS) being prepared for the proposed Bonshaw Solar Farm (the 'Project'), located off the Bruxner Highway, Bonshaw NSW.

This work has been conducted to conform to the Secretary's Environmental Assessment Requirements (SEARs) for the Project. In line with the SEARs, this Cultural Heritage Assessment (CHA) has been prepared in accordance with the Code of Practice for Archaeological Investigations of Aboriginal Objects in New South Wales (Code of Practice) (DECCW 2010a), the Aboriginal Cultural Heritage Consultation requirements for Proponents (DECCW 2010b), The Australia ICOMOS Charter for Places of Cultural Significance, The Burra Charter, 2013 (Burra Charter), and the NSW Heritage Manual.

1.2 Objectives

This CHA assesses the potential impacts of the Project on Aboriginal and historic cultural heritage, and prepares strategies to manage risks to identified heritage values during construction and operation.

This report documents:

- the consultation process undertaken with Aboriginal communities for the Project and their involvement in the Project;
- the landscape and natural resources of the Project Area (PA) in order to establish background parameters;
- a synthesis of local and regional Aboriginal archaeological research to develop a contextual basis for predictive models;
- a review of relevant heritage databases including Australian Heritage Database, the NSW Office
 of Environment and Heritage (OEH) Aboriginal Heritage Information Management System
 (AHIMS) database; the NSW State Heritage Inventory, and Schedule 5 of the Inverell Local
 Environmental Plan 2013;
- a review of the PA's non-Aboriginal history to gain an understanding and appreciation of past land uses and associated historical ground disturbance;
- a predictive model for Aboriginal and historic site types and location relevant to the PA;
- the archaeological methodology implemented during the study;
- the cultural and archaeological sensitivity of landforms that may be subject to impacts;
- the field survey results;
- the significance of any located Aboriginal objects and places;
- a description of the Project and whether or not it has the potential to result in impacts to Aboriginal cultural heritage and historic heritage items; and
- provision of management and mitigation measures based on the results of the investigation.

1.3 Description of the Proposed Development

1.3.1 Location

The proposed Bonshaw Solar Farm is located in the Inverell Local Government Area (LGA) approximately 16 km south of Bonshaw and 66 km north of Inverell. The PA, as shown in *Figure 1.1*, is approximately 352 ha with elevations ranging from 329 m to 500 m Australian Height Datum (AHD) and is located on Lot 2, DP1039185 (the 'Site').

The landholder has been consulted and provided in principle support to be finalised with a landholder agreement. Lot subdivision is not proposed.

The PA is partially bounded on its northern boundary by the Bruxner Highway. The eastern boundary is partially bounded by Glenrock Road. An unsealed road extends perpendicularly from Bruxner Highway towards the southern end of the PA then bends westerly towards the existing 330kV TransGrid Dumaresq Substation located to the south-western from the Subject Land.

The PA consists of open space, and creek and bush areas. Aerial imagery suggests that a significant proportion of the land across the PA had previously been cleared with portions historically being used for agriculture purposes.

1.3.2 Proposed Development

GAIA is proposing to develop a large scale solar photovoltaic (PV) generation facility and associated infrastructure with a capacity of 500 MW within the PA.

The Project will involve the construction, operation and maintenance of a solar PV generation facility and associated infrastructure, supplying electricity to the national electricity grid. The development footprint is approximately 700 ha within the PA. The proposed development footprint is shown in *Figure 1.2*.

The Project would include:

- a network of PV modules in a fixed or tracking arrangement;
- a site office (two proposed options);
- a access tracks from Bruxner Highway;
- underground or overhead cabling for connection between arrays and inverters and transformers;
- parking and internal access tracks;
- perimeter security fencing;
- battery storage; and
- two grid connection options to the 330kV TransGrid Dumaresq Substation, on the boundary of the PA.


The location within the PA of the PV modules, cabling, battery storage and substation and switchyard infrastructure is subject to further detailed design during the preparation of the EIS.


1.4 Authorship

Katherine Deverson (ERM Heritage Consultant) authored the report and Erin Finnegan (Principal Heritage Consultant) undertook a technical review. Amanda Antcliff (ERM Project Manager) and Paul Douglass (ERM Partner) undertook quality assurance (QA) reviews of the report.

1.5 Report Structure

Chapter 1	Introduction;
Chapter 2	Legislative framework and statutory requirements;
Chapter 3	Aboriginal community consultation process undertaken for the CHA;
Chapter 4	Environmental and landscape background relating to the PA;
Chapter 5	Archaeological context of the PA, including known and potential heritage sites within and in near vicinity to the PA;
Chapter 5	Aboriginal heritage predictive model;
Chapter 6	Historical background of the PA;
Chapter 7	Survey methodology and results;
Chapter 8	Significance assessment of heritage sites located within the PA;
Chapter 9	Impact assessment; and
Chapter 10	Conclusions and Recommendations.
Appendix A	Aboriginal Stakeholder Consultation Log
Appendix B	Survey Methodology
Appendix C	AHIMS Database Search Results





2. RELEVANT LEGISLATION

Together with best practice principles outlined in documents such as *The Burra Charter* (Australia ICOMOS 2013a), the following legislative and regulatory context forms the basis of the framework within which the historic and Aboriginal heritage values of the PA must be considered.

Aboriginal cultural heritage in NSW is protected by the *National Parks and Wildlife Act 1974* (NP&W Act). Land managers are required to consider the effects of their activities, or proposed development, on the environment under several pieces of legislation, principally the *Environmental Planning and Assessment Act* 1979 (EP&A Act). Cultural heritage, which includes Aboriginal and historical heritage, is subsumed within the definition of "environment". In certain circumstances, Commonwealth legislation protecting heritage may also apply to heritage places in NSW. The key legislation applying to the Project is summarised below in *Table 2.1*.

Table 2.1 Key Legislation

State Legislation

Environmental Planning and Assessment Act 1979 (NSW)

The principal NSW planning legislation is the *Environmental Planning and Assessment Act 1979* (EP&A Act). The EP&A Act provides a system of environmental planning and assessment administered by the NSW Department of Planning and Environment (DP&E). The EP&A Act establishes when and how a development or activity is to be assessed and who is the relevant approval or determining authority.

Section 4.36 (2) of the EP&A Act states that "a State environmental planning policy may declare any development, or any class or description of development, to be State significant development".

Part 2 Clause 8 of the State Environmental Planning Policy (State and Regional Development) 2011 (State and Regional Development SEPP) states that:

- 1. Development is declared to be State significant development for the purposes of the Act if:
 - a. the development on the land concerned is, by the operation of an environmental planning instrument, not permissible without development consent under Part 4 of the Act, and
 - b. the development is specified in Schedule 1 or 2.

Schedule 1 and 2 of the State and Regional Development SEPP contains an extensive list of developments that are considered State Significant Development (SSD). Schedule 1 Cl 20 identifies the following as SSD:

Electricity generating works and heat or co-generation

Development for the purpose of electricity generating works or heat or their co-generation (using any energy source, including gas, coal, biofuel, distillate, waste, hydro, wave, **solar** or wind power) that:

- a. has a capital investment value of more than \$30 million, or
- b. has a capital investment value of more than \$10 million and is located in an environmentally sensitive area of State significance.

The project is a development for the purpose of electricity generation using a solar energy source with a capital investment value (CIV) of more than \$30 million. The Site is not located in an environmentally sensitive area of State Significance (as defined in the State and Regional Development SEPP).

Solar Energy Systems are considered permissible with consent through clause 34 (7) of State Environmental Planning Policy (Infrastructure) 2007.

The project meets both the requirements of clause 8 of the State and Regional Development SEPP as it is not permissible without development consent and is development specified in Schedule 1. Therefore, the project is SSD for the purposes of the EP&A Act.

State Legislation

National Parks and Wildlife Act 1974 (NSW)

The NSW National Parks and Wildlife Services (NPWS) provide guidelines for Aboriginal heritage assessment, including those conducted under the EP&A Act. Where Aboriginal heritage assessment is conducted under the Integrated Development Approval process, a more detailed set of NPWS guidelines applies.

All Aboriginal objects within the State of New South Wales are protected under Part 6, and particularly Section 90, of the NPW Act.

Under Section 5 of the Act, "Aboriginal Object" means any deposit, object or material evidence (not being a handicraft made for sale) relating to the Indigenous habitation of the area that comprises New South Wales, being habitation before or concurrent with (or both) the occupation of that area by persons of non-Aboriginal extraction, and includes Aboriginal remains.

Sites of traditional significance that do not necessarily contain archaeological materials may be gazetted as 'Aboriginal places' and are protected under Section 84 of the Act. This protection applies to all sites, regardless of their significance or land tenure. Under Section 90, a person who, without first obtaining the consent of the Director-General, knowingly destroys, defaces or damages, or knowingly causes or permits the destruction or defacement of or damage to, an Aboriginal object or Aboriginal place is guilty of an offence.

Heritage Act 1977 (NSW)

The NSW Heritage Act 1977 establishes the NSW Heritage Council and the State Heritage Register (SHR). The aim of the Act is to conserve the heritage of New South Wales. The aim of heritage management is not to prevent change and development, but to ensure that the heritage significance of recognised heritage items is not harmed by changes.

The SHR is a separate listing to the State Heritage Inventory and includes items which are accorded SHR listing through gazettal in the NSW Government Gazette. Nominated items are considered by the NSW Heritage Council which then makes a recommendation to the Minister for Heritage. The Minister is empowered to place Interim Heritage Orders (IHO) on an item of potential State significance on the basis of advice received from the Heritage Council:

- a. An item is important in the course, or pattern, of NSW's cultural or natural history.
- **b.** An item has strong or special association with the life or works of a person, or group of persons of importance in NSW's cultural or natural history.
- c. An item is important in demonstrating aesthetic characteristics and/or a high degree of creative or technical achievement in NSW.
- **d.** An item has strong or special association with a particular community or cultural group in NSW for social, cultural or spiritual reasons.
- **e.** An item has potential to yield information that will contribute to an understanding of NSW's cultural or natural history.
- f. An item possesses uncommon, rare, or endangered, aspects of NSW's cultural or natural history.
- **g.** An item is important in demonstrating the principal characteristics of a class of NSW's a) cultural or natural places: or b) cultural or natural environments.

Refer to Section 6.2 of this CHA for results of SHR search relating to the PA.

Commonwealth Legislation

Environment Protection And Biodiversity Conservation Act 1999 (Commonwealth)

The Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act) is the Australian Government's central piece of environmental legislation. It provides a legal framework to protect and manage nationally and internationally important flora, fauna, ecological communities and heritage places — defined in the Act as matters of national environmental significance. The EPBC Act focuses on the protection of matters of national environmental significance, with the states and territories having responsibility for matters of state and local significance.

The Commonwealth Heritage List (CHL) is established under the EPBC Act and comprises places on Commonwealth land or owned by Commonwealth Agencies that are determined to have "significant" heritage value to Australia. The Act also establishes the National Heritage List, comprising places considered to be of "outstanding" heritage value to Australia.

Client: GAIA Australia

State Legislation

Refer to Section 6.2 of this CHA for results of CHL search relating to the PA.

Aboriginal and Torres Strait Islander Heritage Protection Act 1984 (Commonwealth)

The Aboriginal and Torres Strait Islander Heritage Protection Act 1984 protects areas and/or objects which are of significance to Aboriginal people and which are under threat of destruction. The Act can, in certain circumstances override state and territory provisions, or it can be implemented in circumstances where state or territory provisions are lacking or are not enforced. A significant area or object is defined as one that is of particular importance to Aboriginal people according to Aboriginal tradition. The Act must be invoked by or on behalf of an Aboriginal or Torres Strait Islander or organisation.

3. ABORIGINAL COMMUNITY CONSULTATION

This chapter contains details of the Aboriginal community consultation undertaken regarding the Aboriginal cultural heritage of the PA. In accordance with the guideline Aboriginal Cultural Heritage Consultation Requirements for Proponents 2010 (DECCW, 2010b), consultation with Aboriginal people is an essential part of the heritage assessment process to:

- determine potential harm on Aboriginal cultural heritage from proposed activities; and
- inform decision making for any management and mitigation measures where it is determined that harm cannot be avoided.

The guideline also sets out four stages of consultation requirements. Fulfilment of these requirements is outlined below.

3.1 Stage 1: Notification of Project Proposal and Registration of Interest

On behalf of the Proponent, ERM has actively sought to identify stakeholder groups or people wishing to be consulted about the Project and has invited them to register their interest as follows.

To identify relevant stakeholders, letters about the Project (dated 3 July 2018) were supplied to the following bodies:

- Moombahlene Local Aboriginal Land Council (LALC);
- Local Land Services (LLS) South-East Region;
- National Native Title Tribunal (NNTT);
- Native Title Services Corporation (NTS Corp);
- NSW OEH Regional Operations Coffs Harbour Branch;
- Office of the Registrar, Aboriginal Land Rights Act (1983);
- Tenterfield Shire Council; and
- Inverell Shire Council.

The results from the letters indicated two Aboriginal parties with potential interest in the PA. A Project notification letter was sent to each of the identified parties on 24 July 2018. Correspondence from the LLS has suggested that the Site is located within the Toomelah LALC boundaries. Our review of the boundaries placed the Site within Moombahlene LALC. ERM contact the Toomelah LALC to confirm that it is not in Toomelah region, a map with the known LALC boundaries and PA location was generated and forward to the LALC for their review and comment. No response was received following this correspondence.

A local press advertisement requesting Aboriginal party participation was placed in the Inverell Times and Tenterfield Star on 3 July 2018, the advertisements were published on 6 July 2018 and 11 July 2018 respectively (refer to Appendix A). The response period for Aboriginal stakeholders to register an interest in the Project was open for two consecutive weeks.

Project No.: 0464261 Client: GAIA Australia

3.2 Stage 2: Presentation of Information about the Proposed Project

The Aboriginal parties that registered an interest in being consulted are presented in Table 3.1.

Table 3.1 Registered Aboriginal Parties (RAPs)

Organisation al Cultural Sites Ser

Aboriginal Cultural Sites Services
Edgerton Kwiambal
Moombahlene LALC

A study methodology was sent to each of the registered parties (dated 9 August 2018). The letter included:

- an outline of proposed works;
- the proposed methodology and dates for pedestrian survey;
- a request for Aboriginal parties to identify any particular areas of interest within the PA to survey;
 and
- an invitation to attend a site survey between 11 to 13 September 2018.

No comments were received in relation to the proposed methodology. All three groups agreed to attend the site survey.

3.3 Stage 3: Gathering Information about Cultural Significance

All Registered Aboriginal Parties (RAPs) were invited to participate in the field survey on Tuesday 11, Wednesday 12, and Thursday 13 September 2018. Emails and telephone calls were made to each party to confirm attendance and clarify any potential questions or issues about the methodology, and to determine meeting locations and times for each survey day (refer to *Appendix A*).

Information was also sought regarding cultural knowledge of the PA. All groups expressed that the PA holds or possesses potential cultural sensitivity, through historic family connection to and knowledge of the area.

ERM recognise and value the input of all Aboriginal stakeholders in the consultation process. All RAPs were invited to discuss the Project and the results of the field survey during and following the site survey. The aim of this was to ensure that management actions and consultation commitments are developed in full consultation and are based on the recognised cultural heritage values of the sites.

3.4 Stage 4: Review of Draft Cultural Heritage Assessment Report

A draft copy of the CHA report was provided to all Aboriginal parties who registered an interest in the project on 9 May 2019, for the purposes of receiving written or oral general comments and more specific comments on the cultural significance of the PA and the identified sites and the recommended management and mitigation measures.

Feedback was received from two of the RAPs (also refer to *Appendix A*):

- Edgerton Kwiambal endorsed the draft ACHA.
- Aboriginal Cultural Sites Services provided the comments below that highlight the cultural importance of the area:

Yes comment on site I recommend that an AHIP BE ON the sites that they will construct the solar panels .Remove to a safe spot, or find another area for farmers the area is a high occupancy area that was used my Ancestors.

4. ENVIRONMENTAL BACKGROUND

4.1 Environmental Context

Interactions between people and their surroundings are of integral importance in both the initial formation and the subsequent preservation of the archaeological record. The nature and availability of resources, including water, flora, fauna, and stone materials had (and continues to have) a significant influence over the way in which people utilise the landscape.

Alterations to the natural environment also impact upon the preservation and integrity of cultural materials within that environment. Current vegetation and erosional regimes also affect the visibility and detectability of archaeological evidence. For these reasons, it is essential to consider environmental factors as a component in any cultural heritage assessment.

4.1.1 The Northern Highlands Bioregion

Bioregions and sub-bioregions are large, geographically distinct areas of land with common characteristics such as geology, landform patterns, climate, ecological features and plant and animal communities. The Interim Biogeographic Regionalisation for Australia (IBRA) provides a regional and national planning framework for the systematic development of a comprehensive, adequate and representative National Reserve System. Bioregions delineate salient environmental characteristics which can highlight patterns in Aboriginal site patterning.

The PA is located within the Nandewar Bioregion, which is located inland from the coastal regions and bordered by the North Coast, New England Tablelands and Brigalow Belt South. The region includes parts of NSW, and extends into Queensland. The total area of this bioregion is 2,700,313 hectares (equating to 2.59% of NSW) (NSW NPWS 2003).

Table 4.1 Nandewar Bioregion Attributes

	<u> </u>
Characteristic	Description
	The bioregion overlies part of the New England Fold Belt. The New England Fold Belt is the youngest structural feature in NSW and is separated from the Lachlan Fold Belt by the Sydney-Bowen Basin that is filled with Mesozoic sediments. The oldest rocks in the sequence are Devonian sedimentary and volcanic rocks, formed in an island arc environment. The youngest are Triassic sandstones and shales deposited by rivers on the edge of the Gunnedah Basin, about 250 million years ago, at a time when New England was being lifted by intrusions of granite. Major volcanic eruptions occurred in two phases: in the lava field flood basalts of the Inverell area (34-32 and 22-19 millions of years ago), and in a central volcano similar to that in the Nandewar Ranges (21-17 million years ago). The maximum preserved thickness of the flows is 800m in the variety of lavas present. Only the core of the Nandewar volcano remains as exposed plugs and dykes. Flows from the New England centres buried river gravels and lake sediments that are now being exposed and contain deposits of tin, sapphires and diamonds. A narrow strip of ultrabasic rocks, including serpentinites that are derived from a deep ocean floor, marks the suture where a former island arc complex was linked to the Australian mainland. These rocks pass through Woodsreef and Tamworth where they are associated with limestones in which karst landscapes are formed. The composition of these rocks is so unusual that they always have distinct soils and vegetation

Characteristic	Description
Landforms	The Nandewar Bioregion is formed on Palaeozoic sedimentary rocks on the western edge of the New England Tablelands and includes the Tertiary basalts of Inverell and Kaputar. The hilly landscapes are warmer but drier than the tablelands and carry vegetation communities more typical of the western slopes, with some tableland species.
Soils	The bioregion is characterised by clay or loam soils, but siliceous soils derived from acid volcanic rocks are also found. On the sedimentary rocks, shallow stony soils occur on ridges passing to texture contrast soils on almost all slopes. These change in colour from red brown subsoils on upper slopes to yellow subsoils on lower slopes. They support diverse vegetation communities that are also affected by altitude. The granites develop gritty shallow profiles between outcrops and tors on the crests, grading to harsh texture contrast soils with yellow clay subsoils that are prone to gully development on the lower slopes. Basalt areas on Kaputar have frequent rock outcrops interspersed with shallow, stony, brown loams. Black earths are found on lower slopes and in valleys. In the Inverell area the basalts develop black earth profiles that thicken downslope and, where the underlying sands and gravels are exposed, the coarse sandy soils may develop podsol pans and support different vegetation. Alluvial loams and clays with moderate to high fertility are found in the valleys. Dark, alkaline, pedal clays develop on limestone, and the serpentinites have shallow stony profiles with concentrations of elements that are toxic to many plants.
Vegetation	The vegetation of the Nandewar Bioregion is influenced primarily by geology and the influence of altitude on temperature and rainfall. The bioregion is characterised by box woodlands that occur on clay or loam soils, typically at low to mid elevation in agriculturally productive areas. The principal dominants of these box woodlands are white box (Eucalyptus albens), yellow box (Eucalyptus melliodora), Blakely's red gum (Eucalyptus blakelyi) and grey box (Eucalyptus mollucana). Bimbil box (Eucalyptus populnea subsp. bimbil), fuzzy box (Eucalyptus conica) and western grey box (Eucalyptus microcarpa) also occur, particularly in the western half of the bioregion.

Source: Taken directly from https://www.environment.nsw.gov.au/bioregions/NandewarBioregion.htm

4.1.2 Topography and Landforms

The topographical setting incorporates a variety of landforms which have been classified in this assessment according to the definitions set out in Speight (2009). Speight (2009) states that a landscape can be classified by its landform pattern, and then further classified by individual landform elements. The wider landform pattern, for instance, could be one of flood plains or hills. These landscapes can be further categorised into individual landform elements within the wider landform pattern, for instance cliff, foot slopes or valley flats.

According to these definitions, topographic mapping conducted for the Project indicates that the PA is comprised of several landforms that include gentle slopes, upper flats, crests and open depressions. These landforms would have provided suitable areas for hunter gatherers to undertake a range of subsistence activities (for example camping, hunting and tool making). Certain landforms such as flat or gently sloping raised areas near a water source or ridge flats may have been more frequently occupied than others such as areas away from water, where land use may have been more transient in nature.

4.1.3 Flora and Fauna

Prior to European initiated land clearance, the PA would have supported a range of flora and fauna that would have been be utilised by Aboriginal people for subsistence purposes. A broad range of plants would have been available such as white box (Eucalyptus albens), yellow box (Eucalyptus melliodora), Blakely's red gum (Eucalyptus blakelyi) and grey box (Eucalyptus mollucana). Bimbil box (Eucalyptus populnea subsp. bimbil), fuzzy box (Eucalyptus conica) and western grey box (Eucalyptus microcarpa). "Riparian forests of river oak (Casuarina cunninghamiana), sometimes with river red gum (Eucalyptus camaldulensis), occur along the major watercourses, with Blakely's red gum and rough-barked apple (Angophora floribunda) forming the common association along minor drainage lines. Forest and woodlands of northern smooth-barked apple (Angophora leiocarpa) and dirty gum (Eucalyptus chloroclada) are associated with sandstone parent material on the north-western edge of the bioregion" (NSW NPWS 2003).

There is a high diversity of fauna within the bioregion including, woodland birds, diurnal birds such as regent honeyeaters, lorikeets, and little corellas, nocturnal birds such as several owls and bush stone curlews, frogs, turtles, lizards, snakes, koalas, squirrel gliders, greater gliders, and various bats.

4.1.4 Land Use and Disturbance

The PA generally consists of a cleared pastoral landscape primarily utilised for sheep and cattle grazing. Some areas of the upper soil horizon within the PA have been affected by ploughing. Ploughing generally disturbs the upper 300mm of soil horizons (i.e. 300mm is the maximum plough depth), however, depending on ploughing method, it is possible that up to 600mm could be disturbed. Therefore, deeper soil deposits may retain *in situ* Aboriginal archaeological sites. For this reason, the plough zone is not considered as an area of significant disturbance, however, any artefacts located within the top 300-600mm of the soil horizon may be considered as not to be *in situ*, depending on ploughing methods applied in that area in the past.

Other land disturbances to the PA include the construction of roads, land clearance and fencing. Other ground disturbances have also occurred within the PA such as soil erosion and bioturbation (bioturbation is the reworking of soils and sediments by animals and plants). These post depositional processes have likely adversely affected the archaeological record in the PA.

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia

5. ARCHAEOLOGICAL BACKGROUND

The preliminary archaeological and cultural landscape context discussed in this report has been established through a review of documentation relating to regional ethnographic accounts, information from the AHIMS database, and previously conducted archaeological projects and reports. It is noted that there are several limitations to the use of this existing information such as:

- Aboriginal people involved in previous studies may not have disclosed relevant cultural knowledge and the cultural significance of certain areas due to sensitivities in Aboriginal politics;
- no responsibility can be taken by ERM for errors or omissions in primary and secondary source material cited in this report; and
- the AHIMS search results presented below are based on previous archaeological work and is therefore limited to specific locations and field conditions (visibility, time constraints, etc.) and therefore may not necessarily be a true reflection of the archaeological record.

5.1 Ethno-History

Ethnographic information relating to the Aboriginal occupation of the PA has been obtained from sources containing knowledge that has been passed down from Aboriginal people, as well as documentation written by early European settlers and government officials during the mid to late 18th century (Barwick 1984). Human occupation of south east Australia dates from at least 20,000 years BP. Evidence of early Aboriginal occupation of the State has been provided by sites such as the Burrill Lake rock shelter (Lampert 1971), Cloggs Cage (Flood 1980) and New Guinea 2 (Ossa 1995). The major Aboriginal groups thought to traditionally occupy the wider Nandewar Bioregion were the Anaiwan, the Kamilaroi, the Weraerai, and the Kwaimbul (NSW NPWS 2003).

ERM recognises that Indigenous people are the primary source of information on their heritage and culture; the Anaiwan, Kamilaroi, Weraerai, and the Kwaimbul people are the Aboriginal traditional owners of the wider area. This brief history of the region has been compiled through desktop research, regional studies and historical documents. It should be recognised that most of the available ethnographic information relating to the early Aboriginal occupation of the area comes predominantly from historical documentation written by early European settlers and government officials during the mid to late 19th century. However, the Aboriginal ties to the land and cultural significance of the broader region is recognised within this CHA.

The broader area around the PA forms part of the New England Ranges encompassing eastern and western river systems. Walker describes the region as being dominated by undulating uplands, extending as a tableland belt of varying width, with gently rolling country with shallow valleys terminating in the eastern escarpment east of Walcha (1977, pp.11). Australian Aboriginal people occupied the land according to a complex system of spatial organisation and landscape use (Clark 1990, pp.11-14). Individual groups were intimately familiar with their own geographical regions and the seasonal availability of resources within it. Sutton (1989, pp.7-8) writes that Aboriginal people of the area utilised the surrounding natural resources and landscape formations as a part of their subsistence procurement strategies in addition to more traditional hunting methods; for example, they used the trees to trap macropods via nets strung between them and then herded the animals into the nets and they used traps, nets and baskets in the swamps to catch fish, cravfish, hunt water birds and collect their eggs (Rosen 2009). Indigenous people were also able to supplement their diet by digging for yams, roots and foraging for other edible plants, berries and seeds. Grasslands were maintained by burning or 'fire stick farming' to regenerate local plants and vegetation as well as attract large game into the area for hunting (Rosen 2009). Of the flowering and fruiting plants available as food a large number of them grew in spring and summer including geebung, wild parsnip, Apple Berry, spreading brachyloma and Honey Pots (Rosen 2009). A number of these species are encouraged to grow and flower by fire, including the Xanthorrhoea species, and the Imperata cylindrical (Rosen 2009).

The nature of their subsistence practices required Indigenous groups to be seasonally mobile, moving through the landscape and taking advantage of seasonal fluctuations in available resources. As cited in Hudson (2006), researchers McBryde and Belshaw state that Aboriginal populations on the Tablelands moved to warmer climates (eastern coastal and western slopes) during the colder winter months and occupied rock shelters and overhangs likely adapting their diet and subsistence practices to suit their environment as they moved through the landscape. Many plant resources present in the area are known to have been used by Aboriginal people prior to and after European occupation (Rosen 2009). A number of edible plants were available all year round although spring and summer typically would have been the most productive seasons with a proliferation of species including saw-sedge, bearded heath, Australian Bluebell, ferns and the *Styphelia triflora* (Rosen 2009).

Tools used by Aboriginal people of the Northern Tablelands region included spears, clubs, waddies, boomerangs, shields, paddy-melon sticks and fire sticks made of wood; stone was also utilised to create tools including knives, scrapers, ground-edge axes and even composite technologies like hafted stone chisels (McBryde 1974). Spears were both hand-thrown and thrown with the aid of woomeras (spear throwers). Stone axes were sharpened by pulling or rubbing the axe head through cracks or gaps in other stone outcrops or on bedrock material such as granite or sandstone (Rosen 2009).

Traditional or tribal boundaries were not delineated by treaties or surveyor demarcation but were often defined by the people who occupied them and their culture traditions including linguistic associations, social relations and kinship systems, and spiritual links to the land. Through this commonality and shared culture they may have also controlled and regulated who accessed and passed through their territory. Thus, traditional boundaries were most likely fluid and changed position over time. In light of this, the tribal boundaries recorded by European people at, or after, the point of contact can only be considered as current to that period and were probably quite different prior to European observation (Flood 1980, pp.2). To make things more ambiguous, the few European accounts of Aboriginal groups in the Armidale region are limited in detail, often confused in regard to Aboriginal group names and give varying interpretations of territorial boundaries.

Norman Tindale (1974) in his landmark analysis of the 'Aboriginal Tribes of Australia' cites the PA area as the traditional home of the Kwiambal people (refer to *Figure 5.1*).



Figure 5.1 Map of Tribal Boundaries in Aboriginal Australia, PA indicated with blue circle (Tindale 1974)

5.2 Regional Archaeological Background

While many archaeological studies have been carried out in the region, much is focused on the coastal areas of northern NSW rather than the regional area containing the PA. A significant and early example of these studies was carried out by McBryde in 1974 (OzArk EHM 2011). This study dated a rockshelter site to 6,400±300BP, which is the oldest occupation date so far recorded in the coastal region.

Most hinterland studies in the region have been related to development related assessments, these include a series of transmission line assessments carried out by OzArk in the 2000s and at least once by Navin Offer in the 1990s (OzArk EHM 2011). These studies as well as others in the region suggest that the most common Aboriginal site type is an 'open site' (or stone artefact scatter), although art sites, modified trees, rockshelters, grinding grooves, and quarry sites have also been recorded. It has been noted that sites are most likely to appear on ridgelines, spurs, and slopes rather than along creek lines (Navin Officer 1990).

In 2013 Tenterfield Shire Council commissioned an Aboriginal Heritage Study (AMBS 2013) which aimed to identify places of significance within the LGA and to inform the listing of heritage items on the LEP. The study was carried out in consultation with Aboriginal stakeholders in the area. It identified 174 previously recorded Aboriginal cultural heritage sites within the LGA, noting that sites general occurred near watercourses in elevated areas. The study also noted that the distribution of sites that had been recorded reflected locations of previous studies, carried out as part of development activities or for academic purposes, rather than reflecting an accurate or complete distribution of sites likely to be found across the landscape (AMBS 2013).

5.3 Local Archaeological Context

There has been limited previous archaeological research within the PA; however at least one development project has been undertaken in its proximity which has required cultural heritage assessment. The results of this investigation helps to provide an indication of the range, nature and distribution of archaeological sites within the local area.

5.3.1 Dumaresq to Lismore Transmission Line Aboriginal and Historic Heritage Assessment

Between 2009 and 2011 OzArk EHM undertook Aboriginal and Historic heritage assessments as part of a project which proposed a transmission line between the Dumaresq Switching Station (approximately 75m from the PA's western boundary) and the Lismore Substation in far north NSW (OzArk 2011). In 2011, an assessment was carried out on the section of the proposed transmission line easement. Prior to completing the assessment information about previously recorded sites in proximity were gathered from AHIMS, it was noted that 49.2% of previously recorded sites in the area were open sites / artefacts scatters, and 15.2% were modified (scarred) trees. The assessment identified 50 previously unrecorded Aboriginal cultural heritage sites. The assessment found that most cultural heritage sites were located near a water source and that sites in the area are likely to be artefact scatters or modified trees.

5.3.2 OEH AHIMS Register

The AHIMS database provides information concerning previously recorded Aboriginal sites in NSW. AHIMS stores data regarding a sites location, site type, site features and a unique site identification number for all registered Aboriginal heritage sites in NSW. Mapping of an AHIMS database search results will identify any known sites which could be impacted by proposed works as well as help to determine the overall pattern of Aboriginal sites in an area. A summary of the various site types that could be located in the PA can be found in *Table 5.1* and will aid in the development of a site prediction model for the PA.

Page 17

Table 5.1 Cultural Heritage Site Types Definitions

Site types	Definition
Stone artefact scatters (or open camp sites)	Stone artefact scatter sites, also known as open camp sites, are usually indicated by surface scatters of stone artefacts and sometimes fire blackened stones and charcoal. Where such sites are buried by sediment they may not be noticeable unless exposed by erosion or disturbed by modern activities. The term camp site is used as a convenient label which, in the case of open sites, does not necessarily imply that Aboriginal people actually camped on the sites; rather it indicates only that some type of activity was carried out there.
Isolated finds	Sites consisting of only one identified stone artefact, isolated from any other artefacts or archaeological evidence. They are generally indicative of sporadic past Aboriginal use of an area.
Shell middens	Middens consist of accumulations of shell that represent the exploitation and consumption of shellfish by Aboriginal people. Shell species may be marine, estuarine or freshwater depending on the environmental context and middens may also include other faunal remains, stone artefacts, hearths and charcoal.
Shelter sites	Sandstone shelters and overhangs were used by Aboriginal people to provide camp sites sheltered from the rain and sun. The deposits in such sites are commonly very important because they often contain clearly stratified material in a good state of preservation.
Grinding grooves	Grooves resulting from the grinding of stone axes or other implements are found on flat areas of suitable sandstone. They are often located near waterholes or creek beds as water is necessary in the sharpening process. In areas where suitable outcrops of rock were not available, transportable pieces of sandstone were used.
Quarries	These are areas where stone was obtained for flaked artefacts or ground-edge artefacts, or where ochre was obtained for rock paintings, body decoration or decorating wooden artefacts.
Art sites	Aboriginal paintings, drawings and stencils are commonly to be found where suitable surfaces occur in sandstone shelters and overhangs. These sites are often referred to as rock shelters with painted art.
	Rock engravings, carvings or peckings are also to be found on sandstone surfaces both in the open and in shelters. These are referred to as rock engraving sites.
Scarred trees	Scarred trees bear the marks of bark and wood removal for utilisation as canoes, shields, boomerangs or containers. It is commonly very difficult to confidently distinguish between Aboriginal scars and natural scars or those made by Europeans.
Burial sites	Burials may be of isolated individuals, or they may form complex burial grounds.
Stone arrangements, carved trees and	These site types are often interrelated. Stone arrangements range from simple cairns or piles of rocks to more elaborate arrangements; patterns of stone laid out to form circles and other designs, or standing slabs of rock held upright by stones around the base.
ceremonial grounds	Carved trees are trees with intricate geometric or linear patterns or representations of animals carved into their trunks. Ceremonial grounds and graves were often marked by such trees. Bora grounds are a common type of ceremonial site and they are generally associated with initiation ceremonies. They comprise two circles, generally edged with low banks of earth but sometimes of stone, a short distance apart and connected by a path.

A targeted AHIMS search conducted for the PA on 5 March 2018 and again on 28 February 2019 revealed that one site had been previously recorded within the PA, and another nine sites had been previously recorded within 1.5 km of the PA.

The results of the AHIMS searches are shown in Tables 5.2 and 5.3, and Figure 5.2.

Table 5.2 AHIMS Extensive Search Details

Date	Client Service ID	Latitude	Longitude	Number of Sites
5 March 2018	224600	From: -29.2336	From: 151.2741	10
5 March 2016	331690	To: -29.1502	To: 151.4063	10

Table 5.3 AHIMS Registered Sites

Site ID	Site Name	Site Features	Site Types	Approximate Distance to Project Area	Status
11-3-0083	Tenterfield- Dumeresq OS14	Artefact	Open Site	Within PA	Valid
11-3-0041	S19	Artefact (Isolated Find)	Open Site	>1.5 km	Valid
11-3-0047	EL35	Artefact	Open Site	>1.5 km	Valid
11-3-0063	EL34	Artefact	Open Site	>1.5 km	Valid
11-3-0064	EL33	Art (Pigment or Engraved)	Open Site	>1.5 km	Valid
11-3-0065	EL32	Artefact	Open Site	>1.5 km	Valid
11-3-0066	EL36	Artefact	Open Site	>1.5 km	Valid
11-3-0080	TSR Dumaresq River OCS	Artefact (500)	Open Site	>1.5 km	Valid
11-3-0084	Tenterfield- Dumeresq OS15	Artefact	Open Site	>1.5 km	Valid
11-3-0093	Tenterfield Dumaresq OS 4 & PAD	Artefact	Open Site	>1.5 km	Valid

5.3.2.1 Discussion of AHIMS Site #11-3-0083

Only one previously registered site was identified within the PA. The site, identified as AHIMS #11-3-0083 (Tenterfield-Dumaresq OS14) is recorded as an open artefact scatter, containing at least one culturally modified tree. Details from the site recording card indicate that the site covers six (6) discrete scatters, identified within 50 m of each other. The size of the site complex is estimated to be 100 m by 80 m (see *Figure 5.2*). The following description of the site is provided in the site card:

TD-OS14 is located at the confluence of two unnamed water courses 750m to the east of the Dumaresq Switching Station. Artefacts were recorded on both the east bank, central promontory between the water courses and on the west bank of the water courses. Artefact densities appear to be greater on the east bank and to the south of the easement: although, within the proposed transmission line easement, artefacts are present on both banks of the water courses, albeit in lower densities. Across TD-OS14 there appeared to be very little soil depth and it is assessed that there is a low probability of subsurface archaeological deposits being present. The site is also affected by erosion and stock movement .Recorded artefacts were of a local dark chert (that was noted in the water courses adjacent to TD-OS14), quartz, quartzite and a fine-grained indurated mudstone. Modified tree at Northern end.

The description provided in the site recording card suggests that AHIMS Site #11-3-0083 consists of an open camp site, comprised of a number of discrete scatters that may represent activity areas. Considering the location of the site at the confluence of two smaller creeks, there is a possibility that the site extends beyond the recorded boundary.

AHIMS #11-3-0083 was assessed as having moderate to high cultural significance and moderate to low scientific significance at the time of recording.





Figure 5.2 Reference Map from Site Recording Card #11-3-0083 (OzArk 2011)



5.4 Implications for the Project Area – Predictive Model

It is important to note that the level of archaeological potential relates to the likelihood of discovering an Aboriginal object or site, within a location. Further description should then be made as to the potential condition and integrity of the soil matrix and potential site itself. Only once all these factors have been considered, can scientific value start to be assessed for an area with potential. Therefore, whilst scientific value and potential are linked, it must be noted that these values and potentials are not the same and can differ substantially for any single site or area with potential.

Areas with archaeological potential were assessed according to the definitions in *Table 5.4* and includes a consideration of landforms, distance to resources, historic disturbances and past land use, and the results of previous surveys. Proximity to a permanent water supply and landform is the primary factor appearing to determine the location of Aboriginal campsites within the region and scar trees can occur anywhere that trees of a suitable age remain.

Rank	Definition	Example
Very Low potential	Artefacts are very unlikely to occur in situ.	Eroded landforms, reconstructed landscapes, hazardous landscape, developed areas.
Low potential	Artefacts are not normally found in comparable contexts but could occur in low densities making detection unlikely.	Landforms with no specific focus for use, i.e. areas not associated with or in proximity to water sources.
Moderate potential	Artefacts are known to occur in comparable landforms in detectable densities (~1artefact/m²) and there is an unknown possibility for detection.	Landforms with an environmental focus which may have seen seasonal visitation.
High potential	Artefacts are consistently found in comparable landforms or similar environmental contexts and thus will certainly be found in any ground breaking works.	Landforms with known environmental focus encouraging repeat visitation to specific locale, i.e. margins of swamp or near high order creeks.

Table 5.4 Definitions of Archaeological Potential

Archaeological site formation is a complex combination of factors, such as bioturbation and environmental conditions like erosion or the burial of sites through soil movement. Once discarded on the grounds surface, artefacts are often readily incorporated into the topsoil horizons through the process of bioturbation. It is common for dense artefact deposits to exist hidden beneath the grounds surface (cf. Wandsnider and Camilli 1992). Archaeological assessments that do not employ appropriate methods for prediction cannot reliably define an area's archaeological content.

Frequently, only the eroded component of a larger subsurface deposit is detected and recorded as a site. Where soils are soft, sandy or in boggy conditions, artefacts can occur at greater depths below surface level. Therefore, it is crucial that the nature of an area's soils, sands and geomorphology are defined correctly in an archaeological assessment and the resulting archaeological implications identified. An understanding of these factors, linked further to the notions of site integrity and condition, results in an understanding of an area or site's archaeological potential.

The knowledge gained from examining landforms, geology, regional archaeological patterns, and prior archaeological reports have enabled a set of parameters to be established to predict the potential location of Aboriginal sites within the PA. The topographical nature, geology, flora and fauna of the PA would have provided shelter from the elements and a range of subsistence and lithic resources to Aboriginal people. The PA's proximity to a permanent water source, the Dumaresq River and its numerous associated tributaries that run through the PA would have made it a prime source of water and food resources for Aboriginal groups in the area. In line with this and the archaeological context as discussed above, the PA is likely to contain low density stone artefact scatters across all of its landform units. Although the upper layers of soil horizons across the PA have likely been disturbed by non-Aboriginal agricultural activities such as ploughing, *in situ* archaeological deposits may be present within deeper stratigraphic layers. It is also likely that stone artefacts have not been moved far from their original depositional position by ploughing.

5.4.1 Predictive Model

Based on the background studies and comparative analysis, the following predictions are made for Aboriginal cultural heritage within the PA:

- Aboriginal sites are most likely to be stone artefact sites;
- stone artefact sites are most likely to occur within 400 m of a permanent water course, although smaller creek margins may display scatters of stone artefacts, particularly if near to larger water courses;
- surface expressions of artefacts are most likely to be found on flat or gently sloping terrain;
- surface expressions of artefacts are most likely to be found within proximity of water, or within elevated landforms such as ridge crests or spur line crests, or within proximity of open drainage depressions;
- high density open camp sites will likely be representative of a permanent or seasonal occupational camp, and will likely be located within flat raised terrain that is defensible and sheltered from the elements and has access to resources (water, food, wood and potentially stone);
- lower density sites or isolated finds are likely to contain a 'background scatter' and be representative of a more transient movement throughout the landscape, or intermittent activities (hunting, food procurement, etc.);
- elevated areas that afford views of the surrounding landscape are likely to contain stone artefact sites;
- cultural modifications or scars may occur on mature trees with the PA;
- areas of subsurface stone artefact deposits (with or without a display of surface level stone artefacts) may be present in the PA. Such sites are likely to be located at depths of between 0 and 30 cm below ground surface level;
- human burials are rare, and are not likely to be present in the PA, but if present would most likely be in the alluvial soils that make up the PA's flood plain, creek and river terraces or found in crests and hill tops;
- ceremonial sites (bora grounds) may be present on hill tops in the PA, though are very unlikely
 due to their rarity and previous disturbances across the landscape; and
- the most common stone materials used for the manufacturing of artefacts within the local region of the PA are likely to be silcrete, chert and quartz (with some quartzite and volcanic material).

Much of the material used by Aboriginal people to produce survival equipment (such as wood, bone, shell and fibre material) is highly perishable and does not often survive in the archaeological record. Material culture that has survived, often found in locations where Aboriginal people camped, are generally stone artefacts and scarred trees. Stone artefacts, and to a lesser extent scarred trees, are the most likely artefact types to be located within the PA. Aboriginal archaeological sites within the PA are likely to have been impacted upon (disturbed) due to past European farming practices, the development of transport infrastructure (such as road constructions) and trenching for utility services.

6. HISTORICAL CONTEXT

This chapter considers the potential non-Aboriginal heritage values for the PA. It includes a review of primary and secondary resources including available heritage assessments, reports, publications, historical maps and aerial imagery for the local area. This material will be used to help the PA's history and development over time. The following databases were also searched to determine whether known non-Aboriginal ('historic') heritage sites are located within the PA:

- Commonwealth Heritage List (CHL);
- NSW State Heritage Register and Inventory;
- Inverell Local Environmental Plan (LEP) 2012; and
- The National Trust of Australia.

6.1 Historical Overview

The New England region and Inverell district's first European explorers were Allan Cunningham (1827) and John Oxley (1818), who entered the area on separate expeditions (OzArk 2011). Squatters then moved into the Inverell area around 1835, some establishing large holdings throughout the area, including Campbell in 1837, a significant landholder in Inverell (OzArk 2011; Inverell Shire Council 2018). Campbell, having arrived in Australia from Scotland in 1824 and named the area "Inv" a meeting place and "Ell" meaning swans, as a large amount of these birds were in the area at the time (Inverell Shire Council 2018).

In June 1838 the Myall Creek massacre occurred 35 km from Inverell, becoming a famous case in Australia as the first time that Europeans were hanged for killing Aborigines (Inverell Shire Council 2018). Unfortunately it was not the only massacre in the region at this time, however subsequent massacres and violent episodes went unreported or punished (Inverell Shire Council 2018).

The town of Inverell was incorporated as a municipality in 1872 and tin mining in the region ensured the town's economic growth and remained an important industry in the area for the next century (Inverell Shire Council 2018). The Closer Settlements Acts introduced in NSW between 1901 and 1909 helped to create new small towns in the area in an effort to break large landholding squatter's dominance in country areas (Inverell Shire Council 2018).

6.1.1 Township of Bonshaw

It is not clear when the small township of Bonshaw was formed, however it was built to support local agricultural holdings. The earliest cattle station is thought to have been established in the Bonshaw area in the early 1840s and a track from Bonshaw to Ashford developed in the 1850s (Main Roads 1968). A hotel was constructed at Bonshaw prior to 1867, in time for the bushranger, Captain Thunderbolt (Fred Ward) to rob it in May that year (The Armidale Express 1867). Other community buildings were also built in the town including a church, school, and memorial hall (refer to *Photographs 6.1* and *6.2*). In 1878 £750 was provided for the construction of a track from Tenterfield to Bonshaw (Main Roads 1968). Further roads from Bonshaw to other large towns in the area were established and upgraded over the following decades, including to Ashford, Texas, and Inverell, leading to the formation of State Highway No. 16 in 1938 which became NSW's most northern highway from the coast to the Queensland border at Goondiwindi, it was later renamed Bruxner Highway in 1959 (Main Roads 1968).



Photograph 6.1 School at Maidenhead Station, Bonshaw, c. 1880s (Trove Item #48117813)



Photograph 6.2 Memorial Hall, Bonshaw (Trove Item #252402940)

6.1.2 The Project Area

While little is known about the history of the PA, historical parish maps dating from at least 1909 show that the land was used for pastoral purposes. In 1909, a Charles Strathley Jr was the landholder (refer to *Figure 6.1*). No further detail is included on the map (or any later parish maps) to indicate that any buildings or dwellings were present on the site. Strathley is shown to hold a number of titles within this parish and it is possible that he resided elsewhere and used the land purely for grazing purposes. Strathley is listed as a grazier in several historical sources including the Sydney Stock and Station Journal (5 April 1918). According to later parish maps Strathley was still the landholder in 1939. By 1967 a B.F. Hartley held the land title and continued to do so until at least 1979 (NSWLRS 1965; 1979).




Figure 6.1 1909 Parish of Bowman map, PA is indicated in red (NSWLRS 1909)

6.2 Historical Heritage Database Searches

A search of heritage databases was undertaken in August 2018 and again on 27 February 2019 to determine whether any historical heritage items have previously been recorded within the Project Area. The results are outlined below.

6.2.1 Commonwealth Heritage List

There are no places listed on the CHL within or near the Project Area.

6.2.2 State Heritage Register

A search of the State Heritage Inventory (SHI) indicated that there are no places listed under the NSW Heritage Act (NSW State Heritage Register) within or near the Project Area.

6.2.3 Local Environmental Plan

There are hundreds of places of heritage significance items located within the Inverell Council area that are listed on the LEP Schedule 5: Part 1 Heritage Items. In Bonshaw itself, three places are listed, all of which are located in Bonshaw village approximately 16 km NNW of the PA, and these are:

- Bonshaw Cemetery, Spark Street (Item #I019);
- Church, 10986 Bruxner Highway (Item #I018); and
- Memorial Hall, Miller Street (Item #I021).

No items of local historic heritage are located within or near the general Project Area.

6.3 Historical Heritage Predictive Model

There are no registered or known significant historic heritage sites in or near the PA. There may be evidence of early agricultural activities, timber harvesting, domestic dwellings and domestic remains in the PA. However, the historical background suggests that the PA was used as grazing land. Given the long term pastoral grazing and ongoing site disturbance associated with the installation of and maintenance of the transmission line through the southern section of the PA, it is unlikely that there will be substantial historical remains identified within the AP.

7. SURVEY METHODOLOGY AND RESULTS

This chapter provides an overview of the archaeological surveys of the PA that were undertaken between the 11 and 13 September 2018.

7.1 Field Survey Methodology

The September 2018 field survey was conducted according to the survey methodology developed and sent to RAPs on 9 August 2018 (*Appendix B*). The archaeological survey aimed to assess the ground surface of the PA and targeted all soil exposures and zones with low vegetation such as areas of erosion and any tracks or paths.

Where Aboriginal cultural heritage sites were identified, these were mapped and recorded by the survey team for content, GPS location, landscape features and digitally photographed. Notes were made of soil conditions, evidence of ground disturbance and possible spatial extent of sites.

Visibility refers to the amount of ground upon which artefacts could be seen. The presence of vegetation, leaf litter and other variables can obscure visibility, which is expressed as a percentage. An exposure is defined as an area in which ground surface disturbance (usually in the form of erosion) results in the removal of ground cover and soils and permits the detection of archaeological material that was formerly contained within a surface or subsurface context. The level of exposure is determined as a percentage. As a descriptive tool, *Table 7.1* has been devised which indicates the level of ground surface visibility. It is a subjective method of assessment, but provides a useful tool when attempting to describe the level of ground surface visible during field surveys or inspection.

Table 7.1 Ground Surface Visibility Rating

	Description	GSV Rating %
Very Poor	Heavy vegetation, scrub foliage or debris cover, dense tree of scrub cover. Soil surface of the ground very difficult to see.	0-9%
Poor	Moderate level of vegetation, scrub, and / or tree cover. Some small patches of soil surface visible in the form of animal tracks, erosion, scalds, blow outs etc., in isolated patches. Soil surface visible in random patches.	10-29%
Fair	Moderate levels of vegetation, scrub and / or tree cover. Moderate sized patches of soil surface visible, possibly associated with animal, stock tracks, unsealed walking tracks, erosion, blow outs, etc. Soil surface visible as moderate to small patches across a larger section of the PA.	30-49%
Good	Moderate to low level of vegetation, tree or scrub cover. Greater amount of areas of soil surface visible in the form of erosion, scalds, blow outs, recent ploughing, grading or clearing.	50-59%
Very Good	Low levels of vegetation / scrub cover. Higher incidence of soil surface visible due to recent or past land-use practices such as ploughing, grading, mining, etc.	60-79%
Excellent	Very low to non-existent levels of vegetation/scrub cover. High incidence of soil surface visible due to past or recent land use practices, such as ploughing, grading, mining, etc.	80-100%

Each of the different landforms identified in the PA were surveyed, which included slopes, upper flats, crests and plains (refer to *Table 7.3*). Creek lines, mature trees, erosion scours, and vehicle and animal access tracks were all inspected. In order to ensure the highest likelihood of finding Aboriginal sites, the field survey focussed on areas of highest ground visibility although it is noted that the ground surface visibility was poor (10-29%) to fair (30-49%) across the majority of the survey area.

7.2 Field Survey Results

The PA was surveyed over three days in September 2018 by Katherine Deverson and representatives of the RAPs as outlined in *Table 7.2*. The field survey methodology was adopted to pursue the discovery of new archaeological sites, ensure the accurate recording of such sites and provide sufficient information to provide an assessment of the PA's cultural significance. Discussion also included Aboriginal intangible values and the importance of Aboriginal sites to the local community.

Table 7.2 Field Survey Attendees

Name	Organisation
Vicky Duncan	Aboriginal Cultural Sites Services
Samantha Duncan	Edgerton Kwiambal
Darren Daley	Moombahlene LALC

7.2.1 Description of the Project Area

The PA generally consisted of grazing paddocks with dense grass traversing gently undulating terrain, plains, and crest landforms (refer to *Table 7.3* and *Photographs 7.1* and *7.2*). There was generally a low level of ground surface visibility with some ground exposures under trees, in areas of soil erosion and along tracks. Disturbances observed include the development of fencing, farm dams and vegetation clearance. Several woodland areas, along with a number of small unnamed creeks were also located in the PA (*Photograph 7.3*). Exposures associated with tracks, dams, woodland areas and other disturbances were examined for artefacts.

Soils across the PA range from alluvial soils adjacent to watercourses with thin sandy-silty Aeolian soil in colours of deep orange and red, to a light yellow/brown and dark brown. Disturbance to the soil profile has occurred during past episodes of vegetation clearance.

Table 7.3 Landform Summary

Landform	% of landform effectively surveyed	Number of Sites
Slope	1.5%	6
Upper Flat/Crest	1.75%	1
Plain/Open Depression	2.88%	28



Photograph 7.1 Densely grassed paddock in PA, view to creek line (ERM 2018)



Photograph 7.2 Area of exposure along creek to the left, view to south of PA (ERM 2018)



Photograph 7.3 View to north, along unnamed creek (ERM 2018)

7.2.2 Survey Coverage

The PA was examined in 32 survey units, based on fenced paddocks/areas throughout the PA for ease of recording and analysis. These survey units are identified as Survey Unit (SU) 1 to SU32. The location of these survey units is shown in *Figure 7.1* and detailed in *Table 7.4*.




Table 7.4 Survey Coverage

Survey Unit	Landform	Visibility %	Exposure %	Effective Coverage %	Number of Sites
SU1	Plain	Poor 10%	10%	1%	2
SU2	Plain/Mid Slope	Poor 15%	10%	1.5%	2
SU3	Upper Slope/Crest	Poor 10%	10%	1%	0
SU4	Plain/Open Depression	Poor 20%	10%	2%	2
SU5	Plain/Mid Slope	Poor 10%	10%	1%	2
SU6	Plain/Mid Slope	Poor 10%	10%	1%	0
SU7	Plain	Poor 20%	10%	2%	1
SU8	Plain	Poor 15%	10%	1.5%	1
SU9	Upper Slope/Crest	Poor 25%	10%	2.5%	0
SU10	Mid Slope	Poor 10%	10%	1%	0
SU11	Mid Slope	Poor 10%	10%	1%	0
SU12	Plain	Poor 20%	10%	2%	0
SU13	Plain	Poor 10%	10%	1%	0
SU14	Plain	Poor 10%	10%	1%	2
SU15	Plain	Poor 10%	10%	1%	1
SU16	Plain/Mid Slope	Poor 15%	10%	1.5%	1
SU17	Mid Slope	Poor 10%	10%	1%	0
SU18	Plain	Poor 10%	10%	1%	0
SU19	Plain/Mid Slope	Poor 10%	10%	1%	0
SU20	Plain/Open Depression/Mid Slope	Poor 15%	10%	1.5%	0
SU21	Plain	Poor 15%	10%	1.5%	1
SU22	Plain	Poor 10%	10%	1%	2
SU23	Mid Slope	Poor 10%	10%	1%	1
SU24	Plain/Slope/Upper Flat	Good 50%	30%	15%	4
SU25	Plain	Fair 30%	10%	3%	2
SU26	Plain	Poor 10%	10%	1%	1
SU27	Plain	Fair 40%	30%	12%	1
SU28	Plain/Mid Slope	Poor 20%	15%	3%	1
SU29	Plain	Fair 40%	20%	8%	4
SU30	Mid/Upper Slope	Fair 30%	15%	4.5%	2
SU31	Plain	Poor 25%	15%	3.75%	2
SU32	Plain	Poor 25%	15%	3.75%	1

7.2.3 Survey Results

7.2.3.1 Aboriginal Heritage

During the field survey 35 previously unidentified Aboriginal heritage sites were recorded. The sites were located within 3 km of the Dumaresq River to the north, often along small creek lines (*Figure 7.2.* 29 of these sites were stone artefacts including isolated finds and stone artefact scatters. Seven scarred trees were also identified, one of the scarred trees was identified as part of an artefact scatter site. The proximity to water sources and prevalence of artefact scatters or open sites is in line with the predictive model developed in *Section 5.4* and are representative of previously recorded sites in the area. It should be noted that while early studies suggest that sites in the region are less likely to occur along creek lines, this is not the case here and many artefact scatters were located along creeks, rather than in elevated areas or on ridgelines. These sites are described below in *Table 7.5*. Previously unrecorded Aboriginal cultural heritage sites located during the survey were recorded and artefacts and features identified were left *in situ*.

AHIMS Site #11-3-0083

AHIMS Site #11-3-0083 was identified within the PA in 2011 during surveys for the Dumaresq to Lismore 330kV Transmission Line. The 2018 survey conducted by ERM attempted to relocate site #11-3-008; however, this attempt was not successful which comprises of discrete artefact scatters and a scar tree. During the 2018 survey, the AHIMS site was attempted to be relocated. However, the scar tree, which was seen as a key identifier of AHIMS #11-3-0083 (as shown on to *Figure 5.2*) was unable to be relocated. In regards to the artefact scatters that also comprise AHIMS #11-3-0083, it is noted in *Table 7.4* above the ground surface visibility across the PA was generally poor, averaging 10-15% visibility and 1-3% effective survey coverage.

Additionally, as the site complex was recorded as a large series of artefact scatters (measuring 100 m by 80 m, high likelihood that sites located in the general area (between 50-100 m from the edges of the recording may form part of the same greater site. Therefore it was originally thought, during the EIS preparation process, that BSF27, BSF28 and BSF32 may form part of AHIMS #11-3-0083. However, through the assessment process of the SSD project, the Biodiversity and Conservation Division undertook a technical assessment of all the information recorded regarding AHIMS #11-3-0083. The result of this assessment suggests that a technical error in the recording of coordinates has resulted in a misunderstanding about the location of AHIMS #11-3-0038. Having projected the original coordinates, the Biodiversity and Conservation Division has determined that the area was identified during the 2018 survey and given a new site identifier. *Appendix D* provides further detail of the coordinate error and supporting Figures of the originally mapped AHIMS location and of the reprojected location.




Table 7.5 Aboriginal Cultural Heritage Field Survey Results

Site	Survey Unit	Landform	Description	Associated PAD
BSF1	SU31	Plain	Large Artefact Scatter / Open Camp This site comprises a high number of stone artefacts. It is not likely that all surface artefacts were identified and recorded at this site, however more than 50 artefacts were noted, across an area of approximately 350 m (east to west) by 200 m (north to south). The site is located at northern boundary of property along small creek line and vehicle access track. The site is located on a plain set in a wider landscape context of rolling hills. BSF1 is considered to have a moderate potential for further archaeological deposits due to the number of artefacts found with a corresponding low ground surface visibility, as well as its landscape setting and association with the nearby water course. Artefacts located included chert, tuff, basalt and quartz material, a silcrete retouched axe head and quartz round grindstone. Given its location on the creek that (to the south) contains a site also considered to be an occupation site (BSF14) indicating extended or repeated use, and its subsurface potential, it is considered likely that BSF1 may be part of a complex of occupation sites along the creek beginning at BSF1 and continuing south through BSF14 to BSF22.	Yes
BSF2	SU32	Plain	Artefact Scatter This small artefact scatter of approximately six stone artefacts, including silcrete and basalt flakes and cores, and potentially a quartzite grindstone is located across an area of approximately 75 m (east to west) by 25 m (north to south). The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No

Site	Survey Unit	Landform	Description	Associated PAD
			The site may have been formed as a result of wash down from the nearby hill to the south.	
BSF3	SU30	Upper Slope	Artefact Scatter This small artefact scatter of four stone artefacts, including dilite/diorite flakes was identified by RAPS on a slope near the hillcrest on a track. It is located across an area of approximately 75 m (east to west) by 25 m (north to south). The site is located on an upper slope set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from the nearby hill top.	No
BSF4	SU31	Plain	Isolated Find Broken Granite manuport artefact. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No
BSF5	SU14	Plain	Scarred Tree During the survey RAPs identified a felled scarred tree located within proximity to another standing but dead scarred tree (BSF6). The oval shaped scar was approximately 1.4 m in length and 40 cm wide. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	No

Site	Survey Unit	Landform	Description	Associated PAD
BSF6	SU14	Plain	Scarred Tree During the survey RAPs identified a standing (but dead) scarred tree, with an oval shaped scar. Within proximity to another felled scarred tree (BSF5). The oval shaped scar was approximately 1.2 m in length and 25 cm wide. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	No
BSF7	SU30	Mid Slope	Isolated Find Silcrete core located on slope set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No
BSF8	SU28	Mid Slope	Artefact Scatter Small artefact scatter consisting of at least seven blue chert flakes scattered within a 3 m x 5 m area may have been formed as the result of a single knapping event. The site is located on a slope set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No

Site	Survey Unit	Landform	Description	Associated PAD
BSF9	SU29	Plain	Large Artefact Scatter / Open Camp This site comprises a widespread surface artefact scatter evident along a track and yard across an area of approximately 500 m (north to south) by 100 m (east to west), the site also contains an associated scar tree. There is no evidence for subsurface material and it is possible that the site could have been caused by wash down from the nearby hill to the west of from the nearby occupation sites at BSF1 and BSF14. It is located near creek line and on a vehicle access track. The site is located on a plain set in a wider landscape context of rolling hills. The site contains more than 40 artefacts which are mostly flakes including silcrete, chert, and quartz, flakes; basalt and tuff artefacts were also identified. The site associated with a scar tree identified by RAPs during survey which has been recorded as part of the site. Given its location near the creek that (to the north and south) contains a sites that are considered to be occupation sites (BSF1 and BSF14), it is considered likely that BSF9 may form part of an area of wider occupation sites at BSF1 and continuing south through BSF 9, BSF14, and BSF22.	No

BONSHAW SOLAR FARM

Cultural Heritage Assessment

Site	Survey Unit	Landform	Description	Associated PAD
BSF10	SU29	Plain	Artefact Scatter Small artefact scatter, comprising 2 x basalt hammerstones. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No

BONSHAW SOLAR FARM

Cultural Heritage Assessment

Site	Survey Unit	Landform	Description	Associated PAD
BSF11	SU26	Plain	Scarred Tree During the survey RAPs identified a standing, live, scarred tree. The roughly oval shaped scar was approximately 1 m in length, 40 cm wide, and very close to the ground. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	No
BSF12	SU29	Plain	Isolated Find Silcrete core located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No

Site	Survey Unit	Landform	Description	Associated PAD
BSF13	SU29	Plain	Isolated Find Silcrete core located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No
BSF14	SU25	Plain	Large Artefact Scatter / Open Camp This site comprises a high number of stone artefacts. It is not likely that all surface artefacts were identified and recorded at this site, however more than 50 artefacts were noted, across an area of approximately 110 m (east to west) by 330 m (north to south). Located along and extending between two small creek lines and vehicle access track. The site is located on a plain set in a wider landscape context of rolling hills. The site is considered to have a high potential for further archaeological deposits due to the number of artefacts found with a corresponding low ground surface visibility, as well as its landscape setting and association with the nearby water course. Sub-surface artefacts were visible in the soil profile along eroded creek line at southern end of site. Artefacts located included chert, granite, basalt and quartz material. Given its location on the creek that (to the south) contains a site also considered to be an occupation site (BSF14) indicating extended or repeated use, and its subsurface potential, it is considered likely that BSF1 may be part of a complex of occupation sites along the creek beginning at BSF1 and continuing south through BSF14 to BSF22. It is not likely that all surface artefacts were identified and recorded at this site, however more than 50 artefacts were noted, across an	Yes

Site	Survey Unit	Landform	Description	Associated PAD
			area of approximately 350 m (east to west) by 200 m (north to south).	
BSF15	SU15	Plain	Artefact Scatter This site comprises1 x silcrete flake and 1 x broken silcrete hammerstone. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No
BSF16	SU16	Plain	Artefact Scatter This site comprises1 x silcrete flake and 1 x silcrete hammerstone. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No

Site	Survey Unit	Landform	Description	Associated PAD
BSF17	SU23	Mid Slope	Artefact scatter This site comprises approximately eight stone artefact including blue chert flakes, silcrete flakes and one core, across an area of approximately 80 m (NW to SE) by 30 m (SW to NE). The site is located on a slope set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from BSF14 which is nearby to the north.	No
BSF18	SU22	Plain	Scarred Tree During the survey RAPs identified a standing, live, scarred tree. The oval shaped scar was approximately 1.7 m in length, 40 cm wide, and had significant overgrowth. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	No

BONSHAW SOLAR FARM

Cultural Heritage Assessment

Site	Survey Unit	Landform	Description	Associated PAD
BSF19	SU8	Plain	Scarred Tree During the survey RAPs identified a standing, live, scarred tree with a 1.5 m (length) narrow scar that was approximately 40 cm from the ground and showed significant overgrowth. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	No
BSF20	SU7	Plain	Artefact Scatter This site comprises a three chert flakes. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from larger sites, such as BSF14 which is along the same creek line to the north.	No

Site	Survey Unit	Landform	Description		Associated PAD
BSF21	SU24 & SU 27	Plain	Artefact Scatter This site comprises five stone artefacts, including a quartz flake, a large coarse grained flake with evidence for retouch, a large blue/grey coarse grained flake, a chert flake displaying at least 50% cortex, and a basalt axe. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from nearby sites along the creek to the north (such as BSF14) or from a small lightly wooded rise immediately to its north, although no artefacts were identified on the rise.		No
BSF22	SU22	Plain	Artefact Scatter This site comprises a small artefact scatter with approximately 20 artefacts across a ground surface area on approximately 50 m (NW to SE) by 30 m (SW to NE). The site contains mostly basalt flakes and cores, and at least one chert flake. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site is considered to have potential for further archaeological deposits due to noted evidence for sub-surface artefacts in the soil profile in the eroded creek line along the western edge of site. The presence of the site on the high bank of a creek that (further to the north) contains occupation sites indicating extended or repeated use, and its noted subsurface potential means that it may also be an occupation site and a southern extension of what may be a complex of occupation sites along the creek beginning at BSF1 and continuing south through BSF14 to BSF22.	IFRAO Don	Yes

BONSHAW SOLAR FARM Cultural Heritage Assessment

Site	Survey Unit	Landform	Description	Associated PAD
BSF23	SU5	Crest/Upper Flat	Artefact Scatter This site comprises two artefacts a smooth grained modified stone artefact and a granite hammerstone. The site is located on a rocky upper flat set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	No
BSF24	SU25	Plain	Artefact Scatter This site comprises a small artefact scatter of four fine grained (possibly chert) flakes across an area 75 m 40 m. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from nearby sites along the creek to the north (such as BSF14).	No
BSF25	SU24	Mid Slope	Artefact Scatter This site comprises two artefacts, 1 x basalt hammerstone and 1 x granite axe. The site is located on a slope set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from nearby sites and terraced landforms to the west.	No

BONSHAW SOLAR FARM Cultural Heritage Assessment

Site	Survey Unit	Landform	Description		Associated PAD
BSF26	SU24	Plain	Isolated Find This site comprises a silcrete flake. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.		No
BSF27	SU24	Plain	Artefact Scatter This site comprises a small artefact scatter of eight stone artefacts including five chert flakes, two basalt flakes and a hammerstone granite. The site is located in a ploughed field across an area of approximately 25 m (SW to NE) by 50 m (NW to SE), on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.		No
BSF28	SU21	Plain	Artefact Scatter This site comprises a small artefact scatter of three artefacts with a chert, and quartz flakes identified and a quartzite core, across an area of approximately 110 m (SW to NE) by 50 m (NW to SE). The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified. The site may have been formed as a result of wash down from nearby sites along the creek to the north (such as BSF14).	IFRAO 18 de	No

BONSHAW SOLAR FARM

Cultural Heritage Assessment

Site	Survey Unit	Landform	Description		Associated PAD
BSF29	SU5	Plain	Scarred Tree During the survey RAPs identified a standing, but dead, scarred tree. The long oval shaped scar was approximately 2 m in length and 40 cm wide. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.		No
BSF30	SU1	Plain	Isolated Find This site comprises a granite artefact. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	TRAO-2-2	No

BONSHAW SOLAR FARM

Cultural Heritage Assessment

Site	Survey Unit	Landform	Description		Associated PAD
BSF31	SU1	Plain	Isolated Find This site comprises a grindstone. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.		No
BSF32	SU4	Plain	Artefact scatter – This site comprises a small artefact scatter, 1 x basalt scraper and 1 x basalt flake. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	IFRO NE	No
BSF33	SU4	Plain	Artefact scatter – This site comprises a small artefact scatter, 1 x fine-grained core and 1 x basalt core. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	FRAO 19 SE	No

BONSHAW SOLAR FARM

Cultural Heritage Assessment

Site	Survey Unit	Landform	Description		Associated PAD
BSF34	SU2	Mid Slope	Isolated Find This site comprises a grindstone. The site is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.		No
BSF35	SU2	Plain	Artefact scatter This site comprises a small artefact scatter comprising 3 x blue chert flakes identified in ploughed field located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for additional artefacts, though none were identified.	IFRAO 1022	No

7.2.3.2 Historic Heritage

Two areas with historical items were observed during the field survey:

- A surface scatter of several miscellaneous historical items; and
- A smaller surface scatter of glass bottles to the north of the first surface scatter.

The surface scatter of historical items (refer to *Photographs 7.4* to 7.7) included three small ceramic fragments, car parts, bottle fragments, and other metal domestic items. The items were sparsely strewn across an area of approximately 25 m x 20 m on the high bank side of an unnamed small creek. Items appeared to approximately date from (after) the 1940s, which is evidenced by the presence of bottles labelled as property of the Australasian Pickle Company Ltd. There was no evidence of any previous structure at the site nor any indication of a PAD.



Photograph 7.4 Domestic dump site (ERM2018)



Photograph 7.5 Ceramic fragments at domestic dump site (ERM 2018)



Photograph 7.6 Car parts at domestic dump site (ERM 2018)



Photograph 7.7 Discarded bottle, inscriptions says "This bottle always remains the property of the Australasian Pickle Company Ltd" (ERM 2018)

The separate surface scatter of bottles, located approximately 230 m NNW of the domestic dump, consists of five complete bottles sitting on the ground surface (refer to *Photograph 7.8*). The bottles include:

- one brown glass long necked beer bottle;
- one clear cylindrical bottle labelled "Pick-me-up (regd trade mark)" "this bottle is the property of Pick-Me-Up Condiment Co. Ltd Sydney 1946"; and
- three oblong shaped clear bottles, one of which is labelled as "Clements Tonic" (refer to *Photograph 7.9*).

Given the date on the cylindrical bottle it is likely that this smaller bottle dump dates to a similar period as the domestic dump to the south (from/after the late 1940s). Frederick Moore Clements began manufacturing Clements Tonic in Newtown, Sydney in c.1886 and was commercially successful (Haines 1981). The product is still available today (though very likely uses a different formula than the original. The bottles were laying directly on top of the ground surface with no indication of further subsurface items.



Photograph 7.8 Small bottle dump (ERM 2018)



Photograph 7.9 Clements Tonic bottle (ERM 2018)

8. SIGNIFICANCE ASSESSMENT

The heritage values significance assessment for the PA has been assessed in accordance with the Guide to investigating, assessing and reporting on Aboriginal cultural heritage in NSW (OEH 2011), the Code of Practice for the Archaeological Investigation of Aboriginal Objects in NSW (DECCW 2010) and the Aboriginal cultural heritage consultation requirements for proponents (DECCW 2010). This assessment encompasses the four values outlined in the Burra Charter (social, historical, scientific and aesthetic) (Australia ICOMOS 2013), as shown in Figure 8.1.




Figure 8.1 Diagrammatic Presentation of the Significance Assessment Process (OEH 2011)

8.1 Assessment of Aboriginal Cultural Significance

Aboriginal heritage sites, objects and places hold value for communities in many different ways. The nature of those heritage values is an important consideration when deciding how to manage a heritage site, object or place and balance competing land use options. Assessing the cultural significance of a place means identifying the reasons why a place is culturally important.

The OEH Guideline for investigating, assessing and reporting on Aboriginal Cultural Heritage in NSW (2011) states that analysing cultural heritage significance involves two main steps:

- identifying the range of values present, including social, historic, scientific and aesthetic values;
 and
- assessing why they are important.

The Australia ICOMOS Burra Charter provides seven key criteria against which significance can be assessed, providing information about why sites are significant for the community. These criteria are identified as:

- Rarity: whether any or all aspects of a site (type, location, integrity, content and archaeological potential) can be considered common or rare within a local, regional or national context;
- Representativeness: the comparative rarity of the site when considered and contrasted against other similar sites conserved at the local and/or regional level;
- Archaeological landscapes: the study of the cultural sites relating to Aboriginal peoples within the
 context of their interactions in the wider social and natural environment they inhabited.
 Landscapes can be large or small depending upon specific contexts (i.e. local or regional
 conditions); they may also may be influenced by Aboriginal social and demographic factors
 (which may no longer be apparent);

- Connectedness: whether the site can be connected to other sites at the local or regional level through aspects such as type, chronology, content (i.e. materials present, manufacturing processes), spatial patterning or ethno-historical information;
- Integrity & condition: integrity refers to the level of modification a site has been subject to (the cultural and natural formation process) and whether the site could yield intact archaeological deposits, which could be spatially meaningful. Condition takes into account the state of the material, which is especially relevant for organic materials;
- Complexity: the demonstrated or potential ability of a site to yield a complex assemblage (stone, bone and/or shell) and/or features (hearths, fire pits, activity areas); and
- Archaeological potential: the potential to yield information (from sub-surface materials which retain integrity, stratigraphical or not) that will contribute to an understanding of contemporary archaeological interest, or which could be saved for future research potential.

The significance assessment presented in this Section and summarised in Table 8.1 utilises these criteria.

8.1.1 Social Significance

Cultural/social significance concerns the value/s of a place, feature or site to a particular community group/s, in this case the local Aboriginal communities. Aspects of cultural or social significance are relevant to sites, objects and landscapes that are important or have become important to local Aboriginal communities. This importance involves both traditional links with specific areas as well as an overall concern by Aboriginal people for sites and their continued protection. As such, Aboriginal cultural significance can only be determined by the Aboriginal community.

During the field survey component of this assessment and in accordance with the relevant Aboriginal consultation guidelines, Aboriginal representatives were queried about the cultural significance (to individuals and the community more broadly) of the PA generally, specific locations within it and at identified sites. It was indicated by all RAP representatives that the PA holds a high level of cultural significance to Aboriginal people as it is situated within areas that were used for hunting, gathering and camping by past Aboriginal groups and therefore represents Aboriginal occupation of the region, a past way of life and a direct link to their ancestors. The wider landscape, particularly the flora, fauna and water courses associated with the PA are significant as they formed part of an economic resource environment.

The large and somewhat dense artefact scatters BSF1, BSF9, and BSF14 close to and extending along creek lines are thought by RAPs to represent large camping sites or meeting places.

8.1.2 Historic Significance

The PA has not demonstrated historic values of significance to the local Aboriginal community, as identified through consultation. Although the PA has been identified as holding cultural/social significance for the local Aboriginal community, the PA cannot be linked to specific historical events which are significant to the community.

8.1.3 Scientific Significance

The majority of areas of archaeological interest occurring in the PA are common site types within the region. Stone artefact sites including open camp sites (or artefact scatters) and isolated finds are the most common regional sites types, and that is reflected in the results of the field survey undertaken for the Project. The description of the low density scatters do not place them as of exceptionally high standard in terms of condition or content. They are therefore assessed as having a low archaeological significance.

The three largest artefact scatter / open camp sites are fairly artefact dense sites that all extend along and near creek lines and could represent extended or repeated occupation of the area. The sites may also be closely related and possible represent one large occupation site. BSF1 and BSF14 are associated with Potential Archaeological Deposits (PADs) and are particularly dense and contain stone artefacts that are not only the by-product of stone tool making and tool use but also several tools themselves, including hammerstones and a retouched axe head found at BSF1. BSF1 and BSF14 are therefore assessed as having moderate significance.

Scarred trees, while less common, are known to be found in the area. Seven scarred trees were identified by RAPs (during the surveys) (refer to *Figure 7.2* and *Table 7.5*), however they do not present high scientific value due to the apparent age of the trees and relative commonness of the site type, and are assessed as having a low archaeological significance.

8.1.4 Aesthetic Significance

The stone artefact scatters identified within the PA do not reach the threshold of aesthetic significance and do not contribute to the aesthetic value of the broader landscape. The scarred trees identified have a moderate aesthetic value, as visual representations of cultural heritage practices of the local Aboriginal people.

8.2 Aboriginal Heritage Statement of Significance

The PA contains 36 Aboriginal sites and three associated PADs (35 recorded as part of this assessment). The majority of these sites are stone artefact sites including open camp sites and isolated finds. The sites recorded during this study have been assigned scientific significance in terms of rarity, representativeness, archaeological landscape, connectedness, integrity and condition, complexity, and archaeological sensitivity.

The significance rating of the identified stone artefact sites is higher or lower based on the presence of particular stone artefact types, formal tool types, diverse or unusual raw stone materials and the potential for stratified subsurface deposits. The sites identified within the PA are common site types at a local and regional level. Stone artefact sites are the main site type represented in the region and those located within the PA have not demonstrated a significantly greater diversity or complexity in comparison to other known sites within the region. Scarred tree sites have also been identified as part of this assessment.

It is for this reason that all but two sites (BSF1 and BSF14) located within the PA have been assessed as having low archaeological significance (separate to cultural significance). Additionally, AHIMS Site #11-3-0083 (within the development footprint) was previously assessed as having moderate to low archaeological significance.

BSF1, BSF14, BSF22 and AHIMS #11-3-0083 are all within close proximity to watercourses and located within flat terrain with relative shelter from the elements (areas of known Aboriginal occupation) were identified as having moderate potential for subsurface archaeological deposits based on observations of possible subsurface artefacts in soil profiles. These areas incorporate prominent landscape types within the PA (i.e. flat terrain and slightly sloping areas near a water source). Such areas are likely to contain as yet unrecorded Aboriginal sites and/or objects. Careful detailed design of the solar farm following initial heritage survey results has successfully avoided these two sites, however BSF22 and AHIMS #11-3-0083 may still be impacted by the Project. Given the BSF22's proximity to other significant sites (BSF1 and BSF14) thought to indicate extended or repeated occupation on the banks of the same creek as BSF22 to the north, it is possible that though the surface expression of artefacts at BSF22 is less dense than at sites to the north (BSF1, BSF9, and BSF14), it may also be an occupation site and a southern extension of what may be a complex of occupation sites along the creek.

 Table 8.1
 Summary Significance of Aboriginal Heritage Sites

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF1	It is considered that all Aboriginal sites are significant to the local Aboriginal community. No specific comments have been received regarding the social significance of this site. Significance: High	BSF1 does not meet the threshold for significance under this criterion.	BSF1 is a large artefact scatter containing formal tool types and has been assessed as having potential to contain subsurface archaeological material, and therefore may have some research value. It has been identified in proximity two other large scatters long a creek line indicating it may be part of a larger complex. Significance: Moderate	BSF1 does not meet the threshold for significance under this criterion.	Moderate
BSF2	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF2 does not meet the threshold for significance under this criterion.	BSF2 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF2 does not meet the threshold for significance under this criterion.	Low
BSF3	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF3 does not meet the threshold for significance under this criterion.	BSF3 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF3 does not meet the threshold for significance under this criterion.	Low
BSF4	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF4 does not meet the threshold for significance under this criterion.	BSF4 consists of a single manuport stone, transported in the landscape by human action. Significance: Low	BSF4 does not meet the threshold for significance under this criterion.	Low

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF5	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF5 does not meet the threshold for significance under this criterion.	BSF5 is a felled scarred tree identified by RAPs. The site type is relatively common in the region. Significance: Low	BSF5 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate	Moderate
BSF6	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF6 does not meet the threshold for significance under this criterion.	BSF6 is a scarred tree identified by RAPs. The site type is relatively common in the region. Significance: Low	BSF6 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate.	Moderate
BSF7	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF7 does not meet the threshold for significance under this criterion.	BSF7 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF7 does not meet the threshold for significance under this criterion.	Low
BSF8	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF8 does not meet the threshold for significance under this criterion.	BSF8 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF8 does not meet the threshold for significance under this criterion.	Low
BSF9	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF9 does not meet the threshold for significance under this criterion.	BSF9 is a large artefact that has been identified in proximity (between) two other large scatters near a creek line indicating it may be part of a larger complex. Significance: Low	The scarred tree at BSF9 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate	Moderate

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF10	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF10 does not meet the threshold for significance under this criterion.	BSF10 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF10 does not meet the threshold for significance under this criterion.	Low
BSF11	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF11 does not meet the threshold for significance under this criterion.	BSF11 is a scarred tree identified by RAPs. The site type is relatively common in the region. Significance: Low	BSF 11 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate	Moderate
BSF12	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF12 does not meet the threshold for significance under this criterion.	BSF12 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF12 does not meet the threshold for significance under this criterion.	Low
BSF13	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF13 does not meet the threshold for significance under this criterion.	BSF13 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF13 does not meet the threshold for significance under this criterion.	Low
BSF14	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF14 does not meet the threshold for significance under this criterion.	BSF14 is a large artefact scatter containing formal tool types and has been assessed as having potential to contain subsurface archaeological material, and therefore may have some research value. It has been identified in proximity two other large scatters long a creek line indicating it may be part of a larger complex. Significance: Moderate	BSF14 does not meet the threshold for significance under this criterion.	Moderate

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF15	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF15 does not meet the threshold for significance under this criterion.	BSF15 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF15 does not meet the threshold for significance under this criterion.	Low
BSF16	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF16 does not meet the threshold for significance under this criterion.	BSF16 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF16 does not meet the threshold for significance under this criterion.	Low
BSF17	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF17 does not meet the threshold for significance under this criterion.	BSF17 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF17 does not meet the threshold for significance under this criterion.	Low
BSF18	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF18 does not meet the threshold for significance under this criterion.	BSF18 is a scarred tree identified by RAPs. The site type is relatively common in the region. Significance: Low	BSF18 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate	Moderate
BSF19	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF19 does not meet the threshold for significance under this criterion.	BSF19 is a scarred tree identified by RAPs. The site type is relatively common in the region. Significance: Low	BSF19 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate	Moderate

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF20	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF20 does not meet the threshold for significance under this criterion.	BSF20 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF20 does not meet the threshold for significance under this criterion.	Low
BSF21	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF21 does not meet the threshold for significance under this criterion.	BSF21 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF21 does not meet the threshold for significance under this criterion.	Low
BSF22	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF22 does not meet the threshold for significance under this criterion.	BSF22 presents a common low density scatter which has low potential to contribute to archaeological research in the region, however it has been assessed as having potential to contain subsurface archaeological material, and therefore may have some research value. Significance: Moderate - Low	BSF22 does not meet the threshold for significance under this criterion.	Moderate to Low
BSF23	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF23 does not meet the threshold for significance under this criterion.	BSF23 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF23 does not meet the threshold for significance under this criterion.	Low
BSF24	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF24 does not meet the threshold for significance under this criterion.	BSF24 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF24 does not meet the threshold for significance under this criterion.	Low

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF25	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF25 does not meet the threshold for significance under this criterion.	BSF25 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF25 does not meet the threshold for significance under this criterion.	Low
BSF26	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF26 does not meet the threshold for significance under this criterion.	BSF26 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF26 does not meet the threshold for significance under this criterion.	Low
BSF27	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF27 does not meet the threshold for significance under this criterion.	BSF27 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF27 does not meet the threshold for significance under this criterion.	Low
BSF28	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF28 does not meet the threshold for significance under this criterion.	BSF28 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF28 does not meet the threshold for significance under this criterion.	Low
BSF29	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF29 does not meet the threshold for significance under this criterion.	BSF29 is a scarred tree identified by RAPs. The site type is relatively common in the region. Significance: Low	BSF29 is a visual representation of cultural heritage practices of the local Aboriginal people. Significance: Moderate.	Moderate

Cultural	Heritage	Assessment

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF30	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF30 does not meet the threshold for significance under this criterion.	BSF30 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF30 does not meet the threshold for significance under this criterion.	Low
BSF31	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF31 does not meet the threshold for significance under this criterion.	BSF31 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF31 does not meet the threshold for significance under this criterion.	Low
BSF32	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF32 does not meet the threshold for significance under this criterion.	BSF32 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF32 does not meet the threshold for significance under this criterion.	Low
BSF33	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF33 does not meet the threshold for significance under this criterion.	BSF33 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF33 does not meet the threshold for significance under this criterion.	Low
BSF34	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF34 does not meet the threshold for significance under this criterion.	BSF34 consists of an isolated stone artefact, which holds very little research potential and is unlikely to assist in demonstrating depositional processes or land usage. Significance: Low	BSF34 does not meet the threshold for significance under this criterion.	Low

Site ID	Social Significance	Historical Significance	Scientific Significance	Aesthetic Significance	Overall Significance Level
BSF35	It is considered that all Aboriginal sites are significant to the local Aboriginal community. Significance: High	BSF35 does not meet the threshold for significance under this criterion.	BSF35 presents a common low density scatter which has low potential to contribute to archaeological research in the region. Significance: Low	BSF35 does not meet the threshold for significance under this criterion.	Low
AHIMS Site #11- 3-0083	It is considered that all Aboriginal sites are significant to the local Aboriginal community. The 2011 report (OzArk EHM) assessed this site as being of moderate to high cultural significance. Significance: Moderate - High	11-3-0083 does not meet the threshold for significance under this criterion.	The 2011 report (OzArk EHM) assessed this site as being of low to moderate scientific cultural significance. Significance: Moderate - Low	11-3-0083 does not meet the threshold for significance under this criterion.	Moderate

8.3 Assessment of Historic Heritage Significance

Two surface scatters of historical items were identified during the 2018 surveys, however no research into these sites has been undertaken. As these sites are fairly sparse scatter with limited historic artefacts and no evidence for structures or subsurface expressions were noted during the survey, it is considered that these are dumpsites that were utilised once (or twice) by a local household sometime in the last 70 years. No items identified are considered rare or representative, or of containing any value for further research. It is considered likely that other properties in the area contain similar sites. Although the identification of five complete historic glass bottles is interesting, these sites are not considered to reach the threshold for local historic heritage significance.

9. IMPACT ASSESSMENT

The proposed works involve the following actions that have the potential to impact on Aboriginal heritage sites and values:

- site establishment including the provision of access and construction compounds and laydown areas/material storage facilities;
- topsoil stripping and stockpiling in windrows along the edge of the trench;
- trench excavation and stockpiling of the spoil on the opposite side to the topsoil;
- installation of a sand bed layer;
- laying of conductor and earth cables, incorporating direct bury of cable joints (located approximately every 500 m to 1000 m) and marking of those with electronic marker devises; and
- trench reinstatement consisting of placement of sand coverage of cables, installation of a hard cover, then backfilling/compaction of spoil and respread of topsoil;
- the grading of roads and upgrading of existing access roads;
- vehicle movement across eroded tracks:
- the development of new access roads;
- clearance of vegetation;
- the construction of hardstands and laydown areas; and
- on-site equipment storage areas for the construction period.

36 Aboriginal heritage sites (and three associated PADs) have been recorded within the PA. 30 of these sites were stone artefacts including isolated finds and stone artefact scatters. Seven scarred trees were also identified, one of the scarred trees was identified as part of an artefact scatter site. Careful detailed design of the Project footprint has successfully avoided several of these sites, including BSF1 and BSF14 which are considered to have moderate archaeological significance (refer to *Table 9.1* and *Figure 9.1*).

As the Project is still at concept design phase, and no detailed positioning of solar arrays has been developed, there remains some uncertainty around impact to a number of identified Aboriginal heritage sites. It is evident that these sites are within the Project footprint; however, careful avoidance measures applied and integrated at detailed design phase could avoid smaller artefact scatters or isolated finds. In order to facilitate this process, a detailed methodology for avoidance and procedures for surface collection, test excavation and archaeological salvage will be written into the Cultural Heritage Management Plan. Where impact to sites is possible, but has not been confirmed, these sites have been identified as 'Potentially Impacted'.

All scarred trees within the Project footprint will be avoided during detailed design phase and no harm will result from proposed works.


Table 9.1 Summary Impact Assessment of Aboriginal Heritage Sites

Site ID	AHIMS	Overall Significance Level	Assessment of Potential Impact
BSF1	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this site and PAD
BSF2	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF3	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF4	TBD	Low	No Impact Careful detailed design of the development footprint has successfully avoided this site
BSF5	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this scarred tree
BSF6	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this scarred tree
BSF7	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF8	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF9	TBD	Moderate	Partial Impact Careful detailed design of the development footprint has successfully avoided most of this site, however a small section of the site may be impacted
BSF10	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF11	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this scarred tree
BSF12	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF13	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF14	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this site and PAD
BSF15	TBD	Low	No Impact Careful detailed design of the development footprint has successfully avoided this site
BSF16	TBD	Low	No Impact Careful detailed design of the development footprint has successfully avoided this site

Site ID	AHIMS	Overall Significance Level	Assessment of Potential Impact
			Partial Impact
BSF17	TBD	Low	Careful detailed design of the development footprint has successfully avoided most of this site, however a small section of the site may be impacted
BSF18	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this scarred tree
BSF19	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this scarred tree
BSF20	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF21	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF22	TBD	Low	No Impact Careful detailed design of the development footprint has successfully avoided this site
			Potential Impact
BSF23	TBD	Low	Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF24	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF25	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF26	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF27	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF28	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF29	TBD	Moderate	No Impact Careful detailed design of the development footprint has successfully avoided this scarred tree
BSF30	TBD	Low	No Impact Careful detailed design of the development footprint has successfully avoided this site
BSF31	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF32	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF33	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.

Page 68

Site ID	AHIMS	Overall Significance Level	Assessment of Potential Impact
BSF34	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
BSF35	TBD	Low	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.
Tenterfield- Dumaresq OS14	11-3-0083	Moderate	Potential Impact Artefact scatter may be impacted by project works. Avoidance measures have been recommended.



Page 70

10. CONCLUSIONS AND RECOMMENDATIONS

10.1 Conclusions

36 Aboriginal heritage sites and three associated PADs have been recorded within the PA. 30 of these sites were stone artefacts including isolated finds and stone artefact scatters. Seven scarred trees were also identified, one of the scarred trees was identified as part of an artefact scatter site. Careful detailed design of the Project footprint has successfully avoided nine of these sites, including BSF1 and BSF14 which are considered to have moderate archaeological significance.

10.2 Recommendations

The following recommendations are made to assist in ongoing management of identified heritage sites. The management recommendation statements are made in light of:

- the results of background desktop investigation as outlined in this report;
- predictive modelling;
- results of the field survey;
- a heritage significance assessment;
- legislative requirements as outlined in this report; and
- consultation with the relevant Aboriginal organisations.

10.2.1 Aboriginal Heritage

Based on the information presented in this report, specific recommendations for each identified Aboriginal site is detailed in *Table 10.1* and the following general recommendations have been developed:

- personnel involved with ground breaking activities in the PA should undertake cultural awareness training in line with the recommendations below;
- during works, the location of all recorded Aboriginal heritage sites should be clearly marked on all construction plans for the PA, physically and visibly demarcated on the ground, and site foremen informed of their presence and the need to avoid disturbance;
- if suspected Aboriginal heritage objects are found during works, the Unexpected Find Procedure outlined below should be followed and applies to the entire PA;
- continuing Aboriginal involvement in the Bonshaw Solar Farm project. Ongoing consultation with the Aboriginal community and registered Aboriginal stakeholders for the Project should occur during the construction of the Project. The triggers for consultation with the community during construction may include:
- Additional heritage assessments for any changes in Project scope;
- The implementation of the Unexpected Finds Procedure;
- Preparation of a Cultural Heritage Management Plan (CHMP) for implementation during construction;
- Endorsement by RAPs of the heritage information to be contained in the Project induction material;
- a copy of this report should be provided to each of the Aboriginal organisations who expressed an interest in the Project; and
- a copy of this report should be provided to the relevant OEH regional branch.

Page 71

10.2.1.1 Cultural Awareness Training

In order to comply with best practice principles, it is recommended that any personnel involved with ground breaking activities in the PA undertake basic cultural awareness training. This training should be carried out in the form of a pre-work workshop by a qualified heritage specialist in consultation with a Aboriginal Stakeholders, or by a relevant Aboriginal Stakeholder. The heritage induction material, including an unexpected finds procedure should be developed by an appropriately qualified heritage specialist. This training should:

- include information on the Aboriginal archaeological and cultural heritage values of the PA;
- legal responsibilities and statutory obligations for heritage under the NPW Act and the Heritage Act:
- outline the location and type of archaeological sites within the PA including significant landforms and give instructions not to disturb these sites;
- outline the procedures for the discovery of previously unrecorded Aboriginal objects, through the chance finds procedure; and
- only information endorsed for sharing by the registered Aboriginal stakeholders should be included within the induction package for all workers.

10.2.1.2 Unexpected Finds Procedure

If any heritage objects and/or relics, as protected under NSW legislation, are uncovered during the Project, then the following steps should be followed:

- all activity in the immediate area should cease;
- an area of 2 m surrounding the find should be cordoned off with hazard tape to avoid further disturbance;
- an appropriately qualified archaeologist should be consulted;
- OEH should be immediately contacted;
- If the find consists of Aboriginal cultural heritage, the local Aboriginal stakeholder groups should be notified; and
- an appropriately qualified archaeologist should record the location and attributes of the site and determine the significance of the find.

Additional unexpected finds protocols relating to Historical Heritage finds are detailed in Section 10.2.2.

In the event of the discovery of human skeletal material (or suspected human skeletal material) during project activities in the PA the following steps should be followed:

- all activities and/or works in the immediate area must cease;
- the NSW Police must be contacted along with the OEH; and
- any sand/soils removed from the near vicinity of the find must be identified and set aside for assessment by the investigating authorities.

Project No.: 0464261 www.erm.com Version: 3.0 Client: GAIA Australia 19 June 2020

10.2.1.3 Aboriginal Community Endorsement and Recommendations

A draft copy of the CHA report was provided to all Aboriginal parties who registered an interest in the project on 9 May 2019, for the purposes of receiving written or oral general comments and more specific comments on the cultural significance of the PA and the identified sites and the recommended management and mitigation measures in the ACHA. Feedback was received from two of the RAPs (also refer to *Section 3.4* and *Appendix A*).

10.2.1.4 Cultural Heritage Management Plan

In addition to the cultural heritage induction for contractors, ERM recommends that a Cultural Heritage Management Plan (CHMP) be prepared prior to construction commencing. The CHMP would respond to statutory requirements and Conditions of Consent to ensure appropriate management of cultural heritage during the course of construction and operation of the Project. The CHMP would include protocol for surface collection and salvage, test or salvage excavations, and construction monitoring (as required). The CHMP would also provide a methodology for physical demarcation of sites prior to construction commencing, allowing for RAP participation in this process to ensure all significant cultural heritage sites are appropriately managed.

The CHMP should be prepared in consultation with the RAPs and should be endorsed by all RAP groups prior to finalisation.

10.2.1.5 Dissemination of Information

It is recommended that a copy of the final report be provided to each of the Aboriginal organisations who participated in the project. A copy of this report is to be provided to OEH upon finalisation.

10.2.1.6 Identified Aboriginal Heritage Sites

Direct harm to some sites listed in *Table 10.1* is considered unavoidable. A large number of individual artefacts were identified in the sites listed in *Table 10.1*. It is considered that the cumulative impacts resulting from harm to the sites has the potential to significantly and negatively impact on the Aboriginal cultural heritage values of the area. Therefore, it is considered essential to develop management strategies to minimise this impact as far as possible.

The management measures proposed in *Table 10.1* were formulated during discussions on site with Aboriginal representatives. The recommended mitigation of these direct impacts is to facilitate salvage of artefacts prior to harm occurring, followed by reburial (return to country) outside of impact areas, as close to original location as possible. This management strategy has been formulated with consideration of principles of Ecologically Sustainable Development (ESD), in particular intergenerational equity that holds that various aspects of the environment (including cultural heritage) are available for the benefit of future generations.

If impact to BSF22 is unavoidable it is recommended that subsurface testing be carried out to identify the nature and extent of sub-surface Aboriginal objects at the site in accordance with the *Code of Practice*. The presence of the site on the high bank of a creek that (further to the north) contains occupation sites indicating extended or repeated use, and its noted subsurface potential means that it may also be an occupation site and a southern extension of what may be a complex of occupation sites along the creek beginning at BSF1 and continuing south through BSF14 to BSF22. Test excavations at this site may contribute to the understanding of site characteristics and local and regional prehistory. It may also be used to better inform mitigation measures within the PA.

Where there is a time lapse between collection of artefacts and associated reburial, it is proposed that the Project archaeologist undertaking the artefact collection retain any collected artefacts until such time as they can be reburied.

Note that management measures would be carried out in accordance with the SEARS, and any requirements of the RAPs and the OEH, and therefore the specifics may differ from that presented below.

Table 10.1 Management Measures for Identified Aboriginal Heritage Sites				
Site ID	Overall Significance Level	Potential Impact?	Management Measure	
BSF1	Moderate	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.	
BSF2	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.	
BSF3	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.	
BSF4	Low	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.	
BSF5	Moderate	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.	
BSF6	Moderate	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.	
BSF7	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.	
BSF8	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.	
BSF9	Moderate	Partial	Avoidance. Prior to construction commencing, exclusion fencing and exclusion signage should be erected, around the section of the site that will not be impacted by project works and the scarred tree, by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed. It is recommended that if proposed project works cannot avoid the entire site that surface collection/salvage of the section of the site that will be impacted be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.	

Site ID	Overall Significance Level	Potential Impact?	Management Measure
BSF10	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF11	Moderate	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF12	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF13	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF14	Moderate	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF15	Low	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF16	Low	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF17	Low	Partial	Avoidance. Prior to construction commencing, exclusion fencing and exclusion signage should be erected, around the section of the site that will not be impacted by project works and by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed. It is recommended that if proposed project works cannot avoid the entire site that collection/salvage of the section of the site that will be impacted be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.

Site ID	Overall Significance Level	Potential Impact?	Management Measure
BSF18	Moderate	No	Avoidance.
			Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the scarred tree by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF19	Moderate	No	Avoidance.
			Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the scarred tree by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF20	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF21	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF22	Moderate to	No	Avoidance.
	Low		Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed. It is recommended if proposed project works cannot avoid the entire site that subsurface testing be carried out to identify the nature and extent of subsurface Aboriginal objects at the site. Further to this, surface collection/salvage of the section of the site that will be impacted should be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF23	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF24	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF25	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF26	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF27	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF28	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.

Site ID	Overall Significance Level	Potential Impact?	Management Measure
BSF29	Moderate	No	Avoidance. Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the scarred tree by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF30	Low	No	Avoidance. Prior to construction commencing, if works, access tracks or laydowns areas are planned in the same paddock, exclusion fencing and exclusion signage should be erected around the site by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.
BSF31	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF32	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF33	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF34	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
BSF35	Low	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.
11-3-0083	Moderate	Yes	It is recommended that surface collection/salvage be undertaken by Aboriginal stakeholder groups and an appropriately qualified archaeologist prior the commencement of works.

10.2.2 Historic Heritage

While the historic items identified during the this assessment have been assessed as not meeting the threshold for local historic heritage significance, the items at these sites should be carefully collected and offered to a local heritage museum or organisation prior to commencement of project works.

10.2.2.1 Unexpected Historic Heritage Finds Procedure

If any heritage objects and/or relics, as protected under NSW legislation, are uncovered during the Project, then the following steps should be followed:

- all activity in the immediate area should cease;
- and an appropriately qualified heritage professional should be consulted;
- OEH should be immediately contacted; and
- an appropriately qualified heritage professional should record the location and attributes of the site and determine the significance of the find.

In the event of the discovery of human skeletal material (or suspected human skeletal material) during project activities in the PA the following steps should be followed:

- all activities and/or works in the immediate area must cease;
- the NSW Police must be contacted along with the OEH; and
- any sand/soils removed from the near vicinity of the find must be identified and set aside for assessment by the investigating authorities.

REFERENCES

Barwick, D. E. (1984) 'Mapping the Past: An Atlas of Victorian Clans 1835 – 1904', Aboriginal History 8 pp. 100-132.

Clark, I. D. (1990) **Aboriginal Languages and Clans: An Historical Atlas of Western and Central Victoria, 1800-1900.** *Monash Publications in Geography, No.3, Department of Geography and Environmental Science.* Melbourne: Monash University.

DECCW. (2010a) Code of Practice for Archaeological Investigation of Aboriginal Objects in New South Wales.

DECCW. (2010b) Aboriginal Cultural Heritage Consultation Requirements for Proponents.

Flood, J. (1980) **The Moth Hunters: Aboriginal Prehistory of the Australian Alps**. Australian Institute of Aboriginal Studies, Canberra.

Flood, J. (1995) Archaeology of the Dreamtime. Angus and Robertson, Sydney.

Gregory Haines (1981) **'Clements, Frederick Moore (1859–1920)**', Australian Dictionary of Biography, National Centre of Biography, Australian National University, http://adb.anu.edu.au/biography/clements-frederick-moore-5682/text9601, published first in hardcopy 1981, accessed online 27 February 2019

Horton, D. (1996) **Aboriginal Australia** [cartographic material]; Australian Surveying and Land Information Group.

Hudson, S (2006) **Aboriginal Land Use in the Armidale Area**, accessed on 27 February 2019, from https://www.nesst.org.au/wp-content/uploads/2018/06/ABORIGINAL-LAND-USE-IN-THE-ARMIDALE-AREA.pdf

Lampert, R.J. (1971) **Burrill Lake and Currarong: coastal sites in southern New South Wales**. Australian National University, Research School of Pacific Studies, Canberra.

Ossa, P. (1995) **New Guinea II Cave: a Pleistocene site on the Snowy River, Victoria**, Papers by Paul co-authored with Brendan Marshal & Cathie Webb, *Archaeology in Oceania* (1995) Volume: 30, Issue: 1, Pages: 22-35

Main Roads, (1968). **Bruxner Highway,** Extract from March, 1968 Issue of 'Main Roads, Journal of Department of Main Roads New South Wales. Accessed 25 February 2019 from http://www.shop.nsw.gov.au/shopnsw/documents/agency22/RTA003.pdf

McBryde, I (1974). Aboriginal Prehistory in New England – An Archaeological Survey of Northeastern New South Wales. Sydney University Press

NSW NPWS. (2003) **The Bioregions of New South Wales: their biodiversity, conservation and history**. NSW National Parks and Wildlife Service, Hurstville.

Speight, JG. (2009) Landform. In Australian Soil and land survey field handbook (3rd edition). The National Committee on Soil and Terrain (CSIRO Publishing Melbourne).

Sutton, P. 1989. Postvocalic V in an Australian English dialect. Aust. Ling., vol.9, pp.161-163.

The Armidale Express (1867), **Thunderbolt Again**, Saturday May 18th 1867. The Armidale Express. Accessed 26 February 2019 at https://trove.nla.gov.au/newspaper/article/187931693?#

Tindale, N.B. 1974. Aboriginal Tribes of Australia. University of California Press: Berkeley.

Unknown, (n.d).(Photograph) **School on "Maidenhead" Station - Bonshaw, NSW** url=https://trove.nla.gov.au/work/37032664 | accessdate=27 February 2019

Unknown, (n.d).(Photograph)**Bonshaw Memorial Hall** url=https://trove.nla.gov.au/work/229575195 | accessdate=27 February 2019

Walker, G. T. (1977) **Preliminary Report on the Geomorphology of Natural Lagoons in the New England District: Nature, Origin, Modification and Significance**. Unpublished (University of New England: Armidale).

19 June 2020

Page 78

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia

BONSHAW SOLAR FARM	
APPENDIX A	ABORIGINAL STAKEHOLDER CONSULTATION LOG

AGENCIES CONTACTED

Body/Group	Contact Details	Date Sent	Response	Date
OEH Branch : Coffs Harbour	NSW Office of Environment and Heritage Federation House, 24 Moonee Street, Coffs Harbour NSW 2450 info@environment.nsw.gov.au	3/7/18 Via email and post	Refers ERM to contact Mr Craig Archibald and Mr Aaron Broad	11 th July Via post
The Registrar, Aboriginal Lands Right Act	Office of the Registrar, Aboriginal Land Rights Act PO Box 112 Glebe NSW 2037 adminofficer@oralra.nsw.gov.au	3/7/18 Via email and post	No Registered Aboriginal Owners within project Area on the Register of Aboriginal Owners. Refers ERM to contact Toomelah Local Aboriginal Land Council	11 th July Via email
National Native Title Tribunal (NNTT)	enquiries@nntt.gov.au	3/7/18 Via email and post	Overlap report	5/7/18 Via email
Native Title Services Corporation (NTS Corp)	Native Title Services Corporation PO Box 2105 Strawberry Hills NSW 2012 information@ntscorp.com.au	3/7/18 Via email and post		
Inverell Shire Council	Inverell Shire Council Administration Centre 144 Otho Street PO Box 138 Inverell NSW 2360	3/7/18 Via email and post	Referred to LALCs in the Inverell LGA	17 th July 2018 Via post

Body/Group	Contact Details	Date Sent	Response	Date
Tenterfield Shire Council		3/7/18 Via email and post		4/7/18 Via email
Local Land Services – Northern Tablelands	Local Land Services Northern Tablelands 15 Vivian Street PO Box 411 Inverell NSW harry.white@lls.nsw.gov.au	3/7/18 Via email and post		4 th July Via post
Moombahlene Local Aboriginal Land Council	299 Rouse St, Tenterfield NSW 2372 moombahlenelalc1@bigpond.com	3/7/18 Via email and post	` '	4/7/18 Email response

NEWSPAPER ADVERTISMENT

Newspaper	Contact	Date Sent	Date Published
	The Inverell Times		
	t 0267 200 100		
	classifieds.invtimes@fairfaxmedia.co		
	m.au		
Inverell Times	166 Byron Street, Inverell, NSW 2360	03/07/2018	06/07/2018
	The Tenterfield Star		
	t 02 6776 0501		
	classifieds.tentstar@fairfaxmedia.com		
Tenterfield Star	.au	03/07/2018	11/07/2018

Newspaper	Contact	Date Sent	Date Published
	115 Faulkner Street, Armidale, NSW		
	2350		
	www.tenterfieldstar.com.au		

LIST OF STAKEHOLDERS GROUPS IDENTIFIED & PROJECT NOTIFICATION SENT

Organisation/Person	Contact Provided	Date Sent	Comment
Mr Craig Archibald	27 Margaret Street	24 July 2018	No Response - Letter returned 30/08/2018
	Teralba NSW 2284		marked as "RTS no longer as this address"
Aaron Broad	1 Waratah Ave	24 July 2018	No Response
	Albion Park Rail NSW 2527		
	minnamunnung@gmail.com		
Toomelah Local Aboriginal	Toomelah LALC	Phone call and follow	Correspondence from the LLS has
Land Council	0746762348	up email 7 August 2018	suggested that the site is located within the
	toomelahlandcouncil@gmail.com		Toomelah LALC boundaries. Our review of
			the boundaries places the site within
			Moombahlene LALC. Request was sent to
			confirm that it is not in Toomelah region.

REGISTERED ABORIGINAL PARTIES

Organisation/Person	Contact	Date Registered	How the registration was received & any comments
	Vicky Duncan		
Aboriginal cultural	Diane MARLOW		Email response registering an interest in being
sites services	acsworknsw@gmail.com	8 July 2018	consulted

	Samantha Duncan and Liza Talbot. PH: 0422098648 6 Bala st, Ashford N.S.W 2361		Empil management in hairs
Edgerton Kwiambal	duncans528@gmail.com	9 July 2018	Email response registering an interest in being consulted
<u> </u>	Moombahlene LALC) july 2010	Constitution
	PO Box 70		
	Tenterfield ,2372		
	Ph: 0267363219		
	Fax: 0267361486		
	Helen Duroux	4 July 2018 and 17	Email response registering an interest in being
Helen Duroux (CEO)	moombahlenelalc1@bigpond.com	July 2018	consulted

Presentation of information about the proposed project

Presentation of proposed project information

Aboriginal Organisation/Person	Date Sent	Date Reply	Comments, outcomes and/or issues	Notes
Aboriginal cultural sites services	09/08/2018	10/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology and invitation to survey included in letter.	VD & DM responded via email accepting invitation to the survey, no comments on methodology or known cultural heritage values in email.
Edgerton Kwiambal	09/08/2018	13/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology and invitation to survey included in letter.	SD responded via email accepting invitation to the survey, no

Aboriginal Organisation/Person	Date Sent	Date Reply	Comments, outcomes and/or issues	Notes
				comments on methodology or known cultural heritage values in email.
Moombahlene LALC	09/08/2018	03/09/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology and invitation to survey included in letter.	ERM called on 30/08/18 to enquire if LALC was available for fieldwork on these dates or had any feedback on methodology and project information. No answer, left message. Followed up with email stating the same. HD responded via email 03/09/2018.

The record of agreed outcomes and/or contentious issues should be supplied to all registered Aboriginal parties.

Provision of the proposed assessment methodology to the RAP

Aboriginal Organisation/Person	Date Sent	Date Reply	Comment and details of how input has been considered	Notes
Aboriginal cultural sites services	09/08/2018	10/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology, cultural heritage assessment workshop and invitation to survey included in letter.	VD & DM responded via email accepting invitation to the survey, no

Aboriginal Organisation/Person	Date Sent	Date Reply	Comment and details of how input has been considered	Notes
				comments on methodology or known cultural heritage values in email.
Edgerton Kwiambal	09/08/2018	13/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology, cultural heritage assessment workshop and invitation to survey included in letter.	SD responded via email accepting invitation to the survey, no comments on methodology or known cultural heritage values in email.
Moombahlene LALC	09/08/2018	03/09/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology, cultural heritage assessment workshop details included in letter.	ERM called on 30/08/18 to enquire if LALC was available for fieldwork on these dates or had any feedback on methodology and project information. No answer, left message. Followed up with email stating the same. HD responded via email 03/09/2018, stating that they were interested in the methodology

Aboriginal Organisation/Person	Date Sent	Date Reply	Comment and details of how input has been considered	Notes
				and looking forward to fieldwork.

Field Survey or opportunity for RAP to visit the proposed project site

RAP	Representative	Date	Comments
Aboriginal cultural sites services	Vicky Duncan	11/09/2018, 12/09/2018 & 13/09/2018	
Edgerton Kwiambal	Samantha Duncan	11/09/2018, 12/09/2018 & 13/09/2018	
Moombahlene LALC	Darren	11/09/2018, 12/09/2018 & 13/09/2018	

Seek information from RAP on (a) the presence of Aboriginal objects of cultural value and (b) places of cultural value

RAP	Date	Comments	Cultural values identified
Aboriginal cultural sites services	09/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology, cultural heritage assessment workshop and a request to be provided with any information on the (a) the presence of	No values identified to heritage consultation

		Aboriginal objects of cultural value and (b) places of cultural value, if known or appropriate to divulge.	
Edgerton Kwiambal	09/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology, cultural heritage assessment workshop and a request to be provided with any information on the (a) the presence of Aboriginal objects of cultural value and (b) places of cultural value, if known or appropriate to divulge.	
Moombahlene LALC	09/08/2018	Presentation of project information sent with information on known heritage background, proposed survey methodology, cultural heritage assessment workshop and a request to be provided with any information on the (a) the presence of Aboriginal objects of cultural value and (b) places of cultural value, if known or appropriate to divulge.	

ONGOING CONSULTATION

RAP	Contact Made by	Date	Comments	Notes / Follow up
Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	03/09/2018	KD called VD to advise that following further desktop it was believed that further fieldwork days were needed and invited RAP to attend fieldwork one day earlier (Tuesday 11 September) and advised likely that fieldwork would now extend to Friday 14 September. VD	KD sent follow up email to confirm extension of fieldwork.

			confirmed availability to attend.	
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	03/09/2018	KD called SD to advise that following further desktop it was believed that further fieldwork days were needed and invited RAP to attend fieldwork one day earlier (Tuesday 11 September) and advised likely that fieldwork would now extend to Friday 14 September. No answer message left	KD sent follow up email to invite Edgerton Kwiambal to two further fieldwork days.
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	03/09/2018	KD called VD to advise that following further desktop it was believed that further fieldwork days were needed and invited RAP to attend fieldwork one day earlier (Tuesday 11 September) and advised likely that fieldwork would now extend to Friday 14 September. HD advised she would contact LALC heritage officer and advise ERM of their availability.	KD sent follow up email to confirm extension of fieldwork.
Edgerton Kwiambal – Samantha Duncun	Edgerton Kwiambal – Samantha Duncun	04/09/2018	SD emailed to advise she would attend all fieldwork days from 11/09/2018, and email Insurance information	KD emailed to thank SD for her response.
Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	18/09/2018	KD emailed to thank everyone for their	

			participation in fieldwork and advise of invoicing details	
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	18/09/2018	KD emailed to thank everyone for their participation in fieldwork and advise of invoicing details	
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	18/09/2018	KD emailed to thank everyone for their participation in fieldwork and advise of invoicing details	
Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	09/05/2019	KD emailed draft ACHA to RAPs requesting any comments by 07/06/2019.	
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	09/05/2019	KD emailed draft ACHA to RAPs requesting any comments by 07/06/2019.	
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	09/05/2019	KD emailed draft ACHA to RAPs requesting any comments by 07/06/2019.	
Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	12/06/2019	KD emailed checking if VD had any comments on the draft ACHA.	
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	12/06/2019	KD emailed checking if SD had any comments on the draft ACHA.	SD emailed endorsement of the ACHA 12/06
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	12/06/2019	KD emailed checking if HD had any comments on the draft ACHA.	
Edgerton Kwiambal – Samantha Duncun	Edgerton Kwiambal – Samantha Duncun	12/06/2019	SD replied to KD's email advising 100% endorsement of the ACHA draft	KD emailed reply thanking SD for her response.

Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	21/06/2019	KD emailed checking if VD had any comments on the draft ACHA.	VD replied with comment on project recommendations.
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	21/06/2019	KD emailed checking if HD had any comments on the draft ACHA.	
Aboriginal cultural sites services – Vicky Duncan	Aboriginal cultural sites services – Vicky Duncan	21/06/2019	VD replied with comment on project recommendations, that sites should be avoided otherwise salvaged, and that the area appeared to be a high occupancy area used by her ancestors.	KD emailed replied thanking VD for her response, advising that comments would be included in the ACHA.
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	10/07/2019	KD emailed on behalf of the proponent to advise that three sites (scarred trees) BSF18, BSF19 and BSF29 were currently within the projects impact area and to ask for comment on the possibility of the removal of these trees.	
Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	10/07/2019	KD emailed on behalf of the proponent to advise that three sites (scarred trees) BSF18, BSF19 and BSF29 were currently within the projects impact area and to ask for comment on the possibility of the removal of these trees.	VD replied on 13/07/2019, see entry below
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	10/07/2019	KD emailed on behalf of the proponent to advise that	HD replied 11/07/2019, see below entry

			three sites (scarred trees) BSF18, BSF19 and BSF29 were currently within the projects impact area and to ask for comment on the possibility of the removal of these trees.	
Moombahlene LALC – Helen Duroux	Moombahlene LALC – Helen Duroux	11/07/2019	HD emailed to advise that LALC did not support the removal of any scarred trees.	KD replied thanking HD for her response.
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	12/07/2019	KD emailed to advise that significance of scarred trees sites BSF5, BSF6, BSF9, BSF11, BSF18, BSF9, and BSF29 had been revised from low overall significance to moderate significance as a result of corrections in the draft ACHA to reflect the aesthetic and social/cultural values of the sites. Updated version of ACHA also sent.	
Aboriginal cultural sites services – Vicky Duncan	ERM – Katherine Deverson	12/07/2019	KD emailed to advise that significance of scarred trees sites BSF5, BSF6, BSF9, BSF11, BSF18, BSF9, and BSF29 had been revised from low overall significance to moderate significance as a result of corrections in the draft ACHA to reflect the	

			aesthetic and social/cultural values of the sites. Updated version of ACHA also sent.	
Moombahlene LALC – Helen Duroux	ERM – Katherine Deverson	12/07/2019	KD emailed to advise that significance of scarred trees sites BSF5, BSF6, BSF9, BSF11, BSF18, BSF9, and BSF29 had been revised from low overall significance to moderate significance as a result of corrections in the draft ACHA to reflect the aesthetic and social/cultural values of the sites. Updated version of ACHA also sent.	
Aboriginal cultural sites services – Vicky Duncan	Aboriginal cultural sites services – Vicky Duncan	13/07/2019	VD emailed to advise that ACSS did not support the removal of any live scarred trees and that in the event the removal of dead scarred trees notify Kwiembal Elders Group ,Edgerton Kwiembal environment cultural heritage aboriginal corporation also local land Council must be notified for salvage.	KD replied thanking VD for her response.
Edgerton Kwiambal – Samantha Duncun	ERM – Katherine Deverson	25/07/2019	KD emailed checking if SD had any comments on the email regarding possible removal of scarred trees.	

Edgerton Kwiambal – Samantha Duncun	Edgerton Kwiambal – Samantha Duncun	25/07/2019	SD emailed to advise the EK agreed with comments provided by other RAPs that removal of scarred trees should be avoided, however if removal was to occur that it should be done in consultation with RAPs and with them present for salvage.	KD replied thanking SD for her response.
--	--	------------	--	--

Elise Caldwell

From: Katherine Deverson

Sent: Friday, 21 June 2019 12:38 PM

To: Helen Duroux Cc: Amanda Antcliff

Subject: FW: 0464261 - Bonshaw ACHA Draft report for you review

Attachments: 0464261 - Bonshaw SF ACHA Draft.pdf

Hi Helen,

I am just checking in again to see if you had any comments on the Bonshaw Solar Farm draft ACHA that I sent through last month?

It you have had a chance to review it, and are able to email back an endorsement of the report and/or any comments you may have, that would be great!

Kind regards,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



ERM The business of sustainability

From: Katherine Deverson

Sent: Wednesday, June 12, 2019 1:13 PM

To: Helen Duroux <moombahlenelalc1@bigpond.com> **Cc:** Amanda Antcliff <Amanda.Antcliff@erm.com>

Subject: FW: 0464261 - Bonshaw ACHA Draft report for you review

Hi Helen,

I am just checking in to see if you had any comments on the Bonshaw Solar Farm draft ACHA that I sent through last month?

It you have been able to review it, it would be great if you could email back with your endorsement of the report and/or any comments you may have.

Kind regards,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia T 02 8584 8813 or 02 6126 5311 | M 0466 224 250 E katherine.deverson@erm.com | W www.erm.com



From: Katherine Deverson

Sent: Thursday, May 9, 2019 3:15 PM

To: Helen Duroux <moombahlenelalc1@bigpond.com>

Subject: 0464261 - Bonshaw ACHA Draft report for you review

Hi Helen,

Please find attached the Draft Bonshaw Solar Farm ACHA report which was completed following the surveys we carried out in September 2018.

Please review this report and provide any comments by Friday 7 June 2019.

Feel free to contact me anytime if you would like to discuss anything.

Regards,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



The business of sustainability

Elise Caldwell

From: Katherine Deverson

Sent: Friday, 21 June 2019 2:23 PM acsworknsw@gmail.com

Cc: Amanda Antcliff

Subject: RE: 0464261 - Bonshaw ACHA Draft report for you review

Hi Vicky,

Thanks so much for your comment, I will include it in the draft ACHA. As a State Significant Development the Bonshaw Wind Farm does not go through the approvals process that requires an AHIP, however this ACHA will inform the governing body during their development application consideration process and it is expected that the recommendations from the ACHA will be incorporated into the approval, so it is similar to an AHIP in the mitigation requirements for the proponents.

For this project we have recommended, as you have suggested, that the project area be moved where possibly to avoid impacting cultural heritage sites, and so far this has successfully occurred in those high occupancy sites along the creek line at sites like BSF1 (which you will probably remember is the one up near the road where we found lots of artefacts including the retouched axe head).

The draft ACHA recommends avoidance of all sites where possible however direct impact of the project to some sites is considered unavoidable. The draft ACHA recommends that;

- prior to construction all sites that are to be avoided by project works should be fenced so that they are not harmed by construction works or any associated traffic or lay down areas etc.;
- that surface collection/salvage be undertaken at those sites that cannot be avoided by the project by Aboriginal stakeholder groups and an appropriately qualified archaeologist;
- that any artefacts removed from site as part of the salvage works are safely stored until they can be reburied/returned to country in a place agreed upon by RAPs; and
- that ongoing consultation occurs with RAPs throughout the project.

Thanks again for your comment, I really appreciate your input. Feel free to call or email me anytime if you have any further comments or would like to discuss anything in the draft ACHA or to do with the project.

Have a great weekend!

Regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



From: acsworknsw@gmail.com <acsworknsw@gmail.com>

Sent: Friday, June 21, 2019 1:49 PM

To: Katherine Deverson < Katherine. Deverson@erm.com>

Subject: Re: 0464261 - Bonshaw ACHA Draft report for you review

Hi Katherine.

Yes comment on site I recommend that an AHIP BE ON the sites that thay will construct the solor panels. Remove to a safe spot, or find another area for farmas the area is a high occupancy area that was used my my Ancestors.

Kind regards Vicky Duncan

---- Reply message -----

From: "Katherine Deverson" < Katherine.Deverson@erm.com>
To: "acsworknsw@gmail.com" < acsworknsw@gmail.com>

Cc: "Amanda Antcliff" < Amanda.Antcliff@erm.com>

Subject: 0464261 - Bonshaw ACHA Draft report for you review

Date: Fri., Jun. 21, 2019 12:39 PM

Hi Vicky,

I am just checking in again to see if you had any comments on the Bonshaw Solar Farm draft ACHA that I sent through last month?

It you have had a chance to review it, and are able to email back an endorsement of the report and/or any comments you may have, that would be great!

Kind regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



From: Katherine Deverson

Sent: Wednesday, June 12, 2019 1:15 PM

To: acsworknsw@gmail.com

Cc: Amanda Antcliff < Amanda. Antcliff@erm.com>

Subject: FW: 0464261 - Bonshaw ACHA Draft report for you review

Hi Vicky,

I am just checking in to see if you had any comments on the Bonshaw Solar Farm draft ACHA that I sent through last month?

It you have been able to review it, it would be great if you could email back with your endorsement of the report and/or any comments you may have.

Kind regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com

-	Standing and Adapta Standing Standing Standing Adapta Standing
×	

From: Katherine Deverson

Sent: Thursday, May 9, 2019 3:17 PM

To: acsworknsw@gmail.com

Cc: Amanda Antcliff < <u>Amanda.Antcliff@erm.com</u>>

Subject: 0464261 - Bonshaw ACHA Draft report for you review

Hi Vicky,

Please find attached the Draft Bonshaw Solar Farm ACHA report which was completed following the surveys we carried out in September 2018.

Please review this report and provide any comments by Friday 7 June 2019.

Feel free to contact me anytime if you would like to discuss anything.

Regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy Policy

Elise Caldwell

From: Katherine Deverson

Sent: Wednesday, 12 June 2019 4:41 PM

To: Samantha Duncan

Subject: RE: FW: 0464261 - Bonshaw ACHA Draft report for you review

Hi Samantha

Thanks so much for getting back to me. I really appreciate your input on this project.

Kind regards,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



ERM The business of sustainability

From: Samantha Duncan <duncans528@gmail.com>

Sent: Wednesday, June 12, 2019 3:00 PM

To: Katherine Deverson < Katherine. Deverson@erm.com>

Subject: Re: FW: 0464261 - Bonshaw ACHA Draft report for you review

Hi Katherine. I apologise for the delay.

I have read the report and 100% endorse the draft.

Kind regards,

Samantha Duncan.

On 12 Jun 2019 1:17 PM, "Katherine Deverson" < <u>Katherine Deverson@erm.com</u>> wrote:

Hi Sam,

I am just checking in to see if you had any comments on the Bonshaw Solar Farm draft ACHA that I sent through last month?

It you have been able to review it, it would be great if you could email back with your endorsement of the report and/or any comments you may have.

Kind regards,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



ERM The business of sustainability

From: Katherine Deverson

Sent: Thursday, May 9, 2019 3:19 PM

To: Samantha Duncan < <u>duncans528@gmail.com</u>> **Cc:** Amanda Antcliff < <u>Amanda.Antcliff@erm.com</u>>

Subject: 0464261 - Bonshaw ACHA Draft report for you review

Hi Sam,

Please find attached the Draft Bonshaw Solar Farm ACHA report which was completed following the surveys we carried out in September 2018.

Please review this report and provide any comments by Friday 7 June 2019.

Regards, Kat Katherine Deverson Heritage Consultant
Katherine Deverson
Heritage Consultant
Environmental Resources Management Australia Pty Ltd
15 Tench Street Kingston ACT 2604 Australia
T 02 8584 8813 or 02 6126 5311 M 0466 224 250
E <u>katherine.deverson@erm.com</u> W <u>www.erm.com</u>
<u></u>
ERM The business of sustainability
This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED B

This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy Policy

Elise Caldwell

From: Katherine Deverson

Sent: Thursday, 11 July 2019 3:56 PM

To: Helen Duroux Cc: Amanda Antcliff

Subject: RE: Bonshaw Solar Farm

Hi Helen,

Thank you for your feedback.

I very much appreciate it.

Kind regards, Katherine

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | M 0466 224 250

E katherine.deverson@erm.com | W www.erm.com



ERM The business of sustainability

From: Helen Duroux <moombahlenelalc1@bigpond.com>

Sent: Thursday, July 11, 2019 3:37 PM

To: Katherine Deverson < Katherine. Deverson@erm.com>

Subject: RE: Bonshaw Solar Farm

Hi Katherine,

We don't agree to having the trees removed and would advise against removing them at all costs,

Regards

Helen D

From: Katherine Deverson [mailto:Katherine.Deverson@erm.com]

Sent: Wednesday, 10 July 2019 10:39 AM

To: Helen Duroux **Cc:** Amanda Antcliff

Subject: Bonshaw Solar Farm

Hi Helen,

Thank you for your review of the draft ACHA for the Bonshaw Solar Farm.

As you know from our fieldwork and the draft ACHA, numerous sites (including scarred trees and artefact scatters) were located and recorded during our fieldwork.

In considering ecological and heritage constraints within the Project Area the proponent has adjusted the Project's footprint, effectively moving it away from several larger sites located along the creek line to avoid impacting the archaeological record in these areas. However, the Project's proposed footprint currently impacts three scarred trees.

The draft ACHA recommends the following for each of these sites:

Avoidance.

Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the scarred tree by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.

The Proponent would like your comment on the possibility of removing these three scarred trees.

The scarred trees as recorded in the draft ACHA are:

BSF18 Scarred Tree

During the survey RAPs identified a standing, live, scarred tree. The oval shaped scar was approximately 1.7 m in length, 40 cm wide, and had significant overgrowth.

The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.



BSF19 Scarred Tree

During the survey RAPs identified a standing, live, scarred tree with a 1.5 m (length) narrow scar that was approximately 40 cm from the ground and showed significant overgrowth.

The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.



BSF29 | Scarred Tree

During the survey RAPs identified a standing, but dead, scarred tree. The long oval shaped scar was approximately 2 m in length and 40 cm wide.

The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.



If you could kindly review the above information and provide your comments, it would be greatly appreciated.

Kind regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia **T** 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250 **E** katherine.deverson@erm.com | **W** www.erm.com



This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy Policy



This email has been checked for viruses by Avast antivirus software. www.avast.com
 From:
 Katherine Deverson

 To:
 acsworknsw@gmail.com

 Cc:
 Amanda Antcliff

Subject: RE: Bonshaw Solar Farm

Date: Monday, 15 July 2019 5:48:00 AM

Attachments: image001.png

HI Vickv.

Thank you for your response and comment. It is greatly appreciated.

Kind regards,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia **T** 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250 **E** katherine.deverson@erm.com | **W** www.erm.com



From: acsworknsw@gmail.com <acsworknsw@gmail.com>

Sent: Saturday, July 13, 2019 3:34 PM

To: Katherine Deverson < Katherine. Deverson@erm.com>

Cc: Amanda Antcliff < Amanda. Antcliff@erm.com>

Subject: Re: Bonshaw Solar Farm

Hi Kat

Aboriginal cultural site services recommend that live scarred trees stay and notify Kwiembal Elders Group ,Edgerton Kwiembal environment cultural heritage aboriginal corporation also local land Council on dead scare tree to salvage.

Kind regards Vicky Duncan

---- Reply message -----

From: "Katherine Deverson" < Katherine.Deverson@erm.com>

To: "acsworknsw@gmail.com" <acsworknsw@gmail.com>

Cc: "Amanda Antcliff" < Amanda. Antcliff@erm.com >

Subject: Bonshaw Solar Farm Date: Fri., Jul. 12, 2019 4:52 PM

Hi Vicky,

We have recently incorporated your comments into the ACHA and prepared a final version, in doing so we have corrected the overall significance of the scarred tree sites from low to moderate significance which takes into consideration their aesthetic and social/cultural significance.

I have attached the latest version of the ACHA with these changes included. No recommendations have changed since the draft ACHA.

Please let me know if you have any further comments or questions.

Have a great weekend, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia **T** 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250 **E** katherine.deverson@erm.com | **W** www.erm.com



From: Katherine Deverson

Sent: Wednesday, July 10, 2019 1:12 PM

To: acsworknsw@gmail.com

Cc: Amanda Antcliff < <u>Amanda.Antcliff@erm.com</u>>

Subject: RE: Bonshaw Solar Farm

Hi Vicky,

The Draft ACHA is now attached.

Cheers, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia **T** 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250

E <u>katherine.deverson@erm.com</u> | **W** <u>www.erm.com</u>



From: acsworknsw@gmail.com>

Sent: Wednesday, July 10, 2019 1:07 PM

To: Katherine Deverson < <u>Katherine.Deverson@erm.com</u>>

Cc: Amanda Antcliff < <u>Amanda.Antcliff@erm.com</u>>

Subject: Re: Bonshaw Solar Farm

Hi Kath

Can u send report to me please.

---- Reply message -----

From: "Katherine Deverson" < Katherine.Deverson@erm.com>

To: "acsworknsw@gmail.com" <acsworknsw@gmail.com>

Cc: "Amanda Antcliff" < <u>Amanda.Antcliff@erm.com</u>>

Subject: Bonshaw Solar Farm

Date: Wed., Jul. 10, 2019 12:12 PM

Hi Vicky.

No, I didn't get them. When did you send them through?

Cheers,

Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250 **E** katherine.deverson@erm.com | **W** www.erm.com



From: acsworknsw@gmail.com>

Sent: Wednesday, July 10, 2019 12:11 PM

To: Katherine Deverson < <u>Katherine.Deverson@erm.com</u>>

Cc: Amanda Antcliff < <u>Amanda.Antcliff@erm.com</u>>

Subject: Re: Bonshaw Solar Farm

Hi Kath.

Did you get all picks that I took of project area.

---- Reply message -----

From: "Katherine Deverson" < Katherine.Deverson@erm.com>

To: "acsworknsw@gmail.com" <acsworknsw@gmail.com>

Cc: "Amanda Antcliff" < Amanda. Antcliff@erm.com >

Subject: Bonshaw Solar Farm

Date: Wed., Jul. 10, 2019 10:37 AM

Hi Vicky,

Thank you for your review and comment on the draft ACHA for the Bonshaw Solar Farm.

As you know from our fieldwork and the draft ACHA, numerous sites (including scarred trees and artefact scatters) were located and recorded during our fieldwork.

In considering ecological and heritage constraints within the Project Area the proponent has adjusted the Project's footprint, effectively moving it away from several larger sites located along the creek line to avoid impacting the archaeological record in these areas. However, the Project's proposed footprint currently impacts three scarred trees.


The draft ACHA recommends the following for each of these sites:

Avoidance.

Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the scarred tree by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.

The Proponent would like your comment on the possibility of removing these three scarred trees.

The scarred trees as recorded in the draft ACHA are:



	were identified.	
BSF19	Scarred Tree During the survey RAPs identified a standing, live, scarred tree with a 1.5 m (length) narrow scar that was approximately 40 cm from the ground and showed significant overgrowth. The tree is located on a plain set in a wider	Photo I
BSF29	landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	Photo1
	During the survey RAPs identified a standing, but dead, scarred tree. The long oval shaped scar was approximately 2 m in length and 40 cm wide.	?
	The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	

If you could kindly review the above information and provide your comments, it would be greatly appreciated.

Kind regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia **T** 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250

E <u>katherine.deverson@erm.com</u> | **W** <u>www.erm.com</u>



This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy Policy

This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy Policy

This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy Policy

From: Katherine Deverson
To: Samantha Duncan
Subject: RE: Bonshaw Solar Farm

Date: Thursday, 25 July 2019 7:40:00 PM

Attachments: image001.png

image005.jpg image006.jpg image007.jpg

Hi Sam.

Thanks so much for sending through your comments, it is greatly appreciated.

Have a great weekend.

Regards,

Kat

Katherine Deverson

Senior Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street | Kingston | ACT 2604 | Australia

T 02 8584 8813 or 02 6126 5311 | **M** 0466 224 250

E <u>katherine.deverson@erm.com</u> | **W** <u>www.erm.com</u>



From: Samantha Duncan <duncans528@gmail.com>

Sent: Thursday, July 25, 2019 7:19 PM

To: Katherine Deverson < Katherine. Deverson@erm.com>

Subject: RE: Bonshaw Solar Farm

Hi Kat, sorry for the late reply.

I have read through the attachment and agree with the other parties in relation to the scarred trees. If removal can be avoided that would be great due to cultural and spiritual reasons. However, if it is absolutely necessary that the trees be removed than its vital that T.O's (Traditional Owners) of the boundary areas be present and involved in relocation and the removal process.

Hopefully removal can be avoided as it would be extremely disheartening as an indigenous person with cultural connections to the area and objects within it. After all, our history and our future is all depending on the little historical ARTE facts that remain in every tribal area.

Once again I sincerely apologise for the delay.

Kind regards,

Samantha Duncan..

On 25 Jul 2019 7:08 AM, "Katherine Deverson" < Katherine. Deverson@erm.com > wrote:

Hi Sam,

Just checking in to see if you wanted to make a comment about the scarred tree query in the email below. I have received a response from Vicky but wasn't sure if you also wanted to submit any comments?

I have attached the most recent version of the report to this email, although it is the same copy that I sent through to you on 12 July.

I hope you are well.

Kind regards,

Kat

Katherine Deverson

Senior Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street|Kingston|ACT 2604|Australia T 02 8584 8813 or 02 6126 5311 | M 0466 224 250 E katherine.deverson@erm.com | W www.erm.com



From: Katherine Deverson

Sent: Wednesday, July 10, 2019 10:39 AM

To: Samantha Duncan <<u>duncans528@gmail.com</u>> **Cc:** Amanda Antcliff <<u>Amanda.Antcliff@erm.com</u>>

Subject: Bonshaw Solar Farm

Hi Sam,

Thank you for your review and comment on the draft ACHA for the Bonshaw Solar Farm.

As you know from our fieldwork and the draft ACHA, numerous sites (including scarred trees and artefact scatters) were located and recorded during our fieldwork.

In considering ecological and heritage constraints within the Project Area the proponent has adjusted the Project's footprint, effectively moving it away from several larger sites located along the creek line to avoid impacting the archaeological record in these areas. However, the Project's proposed footprint currently impacts three scarred trees.

The draft ACHA recommends the following for each of these sites:

Avoidance.

Prior to construction commencing, exclusion fencing and exclusion signage should be erected around the scarred tree by Aboriginal stakeholder groups and an appropriately qualified archaeologist. At the completion of construction exclusion zone fencing will be removed.

The Proponent would like your comment on the possibility of removing these three scarred trees.

The scarred trees as recorded in the draft ACHA are:

The sca	rred trees as record	ed in the draft ACHA are:
BSF18	Scarred Tree	Photo3
	During the survey RAPs identified a standing, live, scarred tree. The oval shaped scar was approximately 1.7 m in length, 40 cm wide, and had significant overgrowth. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and surrounding area was checked	?
	for stone artefacts, though none were identified.	
BSF19	Scarred Tree	Photo1
	During the survey RAPs identified a standing, live, scarred tree with a 1.5 m (length) narrow scar that was approximately 40 cm from the ground and showed significant overgrowth. The tree is located on a plain set in a wider landscape context of	?
	rolling hills. The exposure and surrounding area was checked for stone artefacts, though none were identified.	
BSF29	Scarred Tree During the survey RAPs identified a standing, but dead, scarred tree. The long oval shaped scar was approximately	Photo 1
	2 m in length and 40 cm wide. The tree is located on a plain set in a wider landscape context of rolling hills. The exposure and	

If you could kindly review the above information and provide your comments, it would be greatly appreciated.

Kind regards, Kat

Katherine Deverson

Heritage Consultant

Environmental Resources Management Australia Pty Ltd

15 Tench Street|Kingston|ACT 2604|Australia T 02 8584 8813 or 02 6126 5311 | M 0466 224 250 E katherine.deverson@erm.com | W www.erm.com



This electronic mail message may contain information which is (a) LEGALLY PRIVILEGED, PROPRIETARY IN NATURE, OR OTHERWISE COVERED BY LAW FROM DISCLOSURE, and (b) intended only for the use of the Addressee (s) names herein. If you are not the Addressee (s), or the person responsible for delivering this to the Addressee (s), you are hereby notified that reading, copying, or distributing this message is prohibited. If you have received this electronic mail message in error, please contact us immediately and take the steps necessary to delete the message completely from your computer system. Environmental Resources Management Australia Pty Ltd (ERM) has systems in place to encourage a virus free software environment, however we cannot be liable for any loss or damage, corruption or distortion of electronically transmitted information, or for any changes made to this information during transferral or after receipt by the client.

Please visit ERM's web site: http://www.erm.com. To find out how ERM manages personal data, please review our Privacy_Policy





November 2018

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia 19 June 2020

9 August, 2018

Helen Duroux Moombahlene LALC PO Box 70 Tenterfield NSW 2372

moombahlenelalc1@bigpond.com

Our Reference: 0464261_SURVEY METHODOLOGY_LALC.DOCX

Dear Helen Duroux,

RE: SOLAR FARM, BONSHAW NSW

Thank you for registering an interest in being consulted for this project.

In accordance with the NSW Office of Environment and Heritage's (OEH) Aboriginal Cultural Heritage Consultation Requirements for Proponents 2010 (the Guidelines), and the Code of Practice for the Archaeological Investigation of Aboriginal Objects in NSW 2010 (Code of Practice), Environmental Resources Management Australia (ERM) is undertaking an Aboriginal cultural heritage assessment (ACHA) associated with the construction and operation of 500 MW capacity solar farm connecting to the existing Dumaresq Substation (Figure 1).

This letter provides the relevant information about the proposed project, the current known heritage values of the study area and the proposed archaeological survey methodology.

In accordance with *Section 4.1.6* of the Guidelines, we would also like to provide Moombahlene LALC with a list of the registered groups for this Project:

- Aboriginal Cultural Sites Services;
- Edgerton Kwiambal; and
- Moombahlene LALC.

Annex A also includes a copy of the Newspaper advertisement that was placed in the Inverell Times on 6 July 2018 and the Tenterfield Star on 11 July 2018.

1. PROPOSED DEVELOPMENT

The proponent for the proposed works is GAIA Australia Pty Ltd (GAIA).

The proposed Project Area (PA) is approximately 1097 hectares (ha) with a development footprint of approximately 650 ha. The land is privately owned rural land, currently used for grazing.

Environmental Resources Management Australia Pty Ltd


15 Tench Street Kingston ACT 2604

PO Box 4160 Kingston ACT 2604

Telephone +61 2 6253 6888 Facsimile +61 2 9299 7502

www.erm.com





With specific reference to Aboriginal cultural heritage and the assessment of potential impacts to the heritage values of the PA, the project will involve the construction, operation and maintenance of a solar PV generation facility and associated infrastructure with a capacity of up to 500 MW, supplying electricity to the national electricity grid.

The Project would include:

- a network of PV modules in a fixed or tracking arrangement;
- a site office (two proposed options);
- three potential access tracts from Bruxner Highway and one from Glenrock Road;
- underground or overhead cabling for connection between arrays and inverters and transformers;
- parking and internal access tracks;
- perimeter security fencing;
- battery storage; and
- two grid connection options to the 330 kV TransGrid Dumaresq Substation, on the boundary of the PA.

2. HERITAGE BACKGROUND

A preliminary due diligence assessment was undertaken by ERM in June 2018. The assessment included a desktop review of the heritage values which have the potential to impact the PA.


The review was limited to matters identified in the following online resources accessed on Tuesday 6 March 2018:

- Australian Heritage Database register search, including: World Heritage List, National Heritage List, Commonwealth Heritage List, National Trust Register, Register of the National Estate (archive);
- NSW State Heritage Inventory includes items listed on the State Heritage Register (SHR) and provides a list of places and items of State Significance made under the Heritage Act. Heritage Council approval is required for works proposed to an item on the SHR;
- Inverell LEP 2012 Schedule 5 and Conservation Areas (Locally listed items);
- Historic Heritage Information Management System (HHIMS) search;

- Aboriginal Heritage Information Management System (AHIMS);
- Schedule 14 of the National Parks and Wildlife Act; and
- Native Title Search.

One active Native Title claim (Comeroi People #NC2011/006) is currently registered within the LGA, and is located on the western side of Rocky Creek Road adjacent to the PA. The PA is not included in this Native Title claim.

An extensive AHIMS register search was conducted for each allotment on 5 March 2018, which identified three previously recorded Aboriginal sites located within the PA. All three of these sites (AHIMS #11-3-0083, #11-3-0084 and #11-3-0093) are reported as artefact scatters, with one potential scar tree also noted at the northern extent of #11-3-0083 (Refer to *Figure* 2). They were recorded as part of the Far North NSW (Dumaresq to Lismore 330kV Transmission Line) Project in 2011 (Ozark, 2011).



3. FIELDWORK METHODOLOGY

Based on the results of the due diligence assessment, the archaeological survey will be undertaken over two days and will aim to identify all Aboriginal sites present within the proposed impact area including the identification of any PADs.

The proposed methodology for the archaeological survey includes:

- the survey will be undertaken on foot where possible with Registered Aboriginal Parties in attendance;
- the survey will consist of all participants traversing the transmission line corridor using walking transects approximately 5 m apart to ensure the entire survey area is covered (subject to surface visibility and accessibility);
- the survey will target each landform in the PA;
- areas of potential such as raised landforms in close proximity to a semipermanent water source will also be targeted;
- areas of exposure and ground visibility will be targeted;
- any areas of interest to the Registered Aboriginal Parties will be targeted;
- the previously recorded sites will be targeted, re-identified and recorded; and
- any cultural heritage information for the PA held by Aboriginal parties will be recorded during the field survey. Any cultural knowledge provided by Aboriginal Stakeholders will be treated in confidence and the information will be distributed according to their wishes.

ERM propose the dates of **12 and 13 September 2018** to undertake this archaeological survey. One representative from your organisation is invited to attend. Details of the meeting time and location for each day will be provided once we have confirmed attendance.

Each representative is responsible for providing their own Personal Protective Equipment (PPE), including sturdy walking boots, long pants and long sleeve shirts (and wet weather gear, if required). Please ensure that you have sun protection and enough water for the day.

Should you wish to participate, you will need to supply ERM with:

- the name of the individual who will attend as your representative;
- a Certificate of Currency for Public Liability insurance, valid for the period of work; and

a Certificate of Currency for Workers Compensation insurance, valid for the period of work.

4. FEEDBACK

ERM requests that you respond to this information package and fieldwork methodology prior to Friday 7 September 2018 with any comments you may have and confirmation of availability for fieldwork.

Please provide feedback to **Katherine Deverson** on the following contact details:

Post: PO Box 4160, Kingston, ACT 2604 Phone: 02 8584 8813 or 02 6126 5311 Email: katherine.deverson@erm.com

If you hold any knowledge of sites within or near the study area or have any specific information concerning the cultural values of the study area, we would be grateful if you could let us know. Our contact details are listed above. Any cultural knowledge provided by Aboriginal Stakeholders will be treated in confidence and the information will be distributed according to their wishes.

Ral Bughen

Yours faithfully,

for Environmental Resources Management Australia Pty Ltd

Katherine Deverson

Paul Douglass **ERM Partner** ERM Archaeologist



APPENDIX C AHIMS EXTENSIVE SEARCH RESULTS

November 2018

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia 19 June 2020



AHIMS Web Services (AWS) Extensive search - Site list report

Your Ref/PO Number: 0450869

Client Service ID: 331690

SiteID 11-3-0041	SiteName S19	Datum AGD	Zone 56	Easting 335382	Northing 6770551	Context Open site	<u>Site Status</u> Valid	SiteFeatures Artefact: -	SiteTypes Isolated Find	<u>Reports</u>
11 0 0011	Contact	Recorders		ert Paton	0,,0001	o pen site	vana	Permits	1001000 1 1110	
11-3-0047	EL35	AGD	56	342100	6767700	Open site	Valid	Artefact : -		
	Contact	Recorders	Rob	ert Paton				<u>Permits</u>		
11-3-0063	EL34	AGD	56	342150	6767550	Open site	Valid	Artefact : -		
	Contact	Recorders	Rob	ert Paton				<u>Permits</u>		
11-3-0064	EL33	AGD	56	342100	6767500	Open site	Valid	Art (Pigment or Engraved) : -		
	Contact	Recorders		ert Paton				<u>Permits</u>		
11-3-0065	EL32	AGD	56	342000	6767000	Open site	Valid	Artefact : -		
	Contact	Recorders	Rob	ert Paton				<u>Permits</u>		
11-3-0066	EL36	AGD	56	342100	6767500	Open site	Valid	Artefact : -		
	Contact	Recorders	Rob	ert Paton				<u>Permits</u>		
11-3-0080	TSR Dumaresq River OCS	GDA	56	335700	6772930	Open site	Valid	Artefact: 500		
	<u>Contact</u> Searle	Recorders	Miss	s.Karen Glove	er			<u>Permits</u>		
11-3-0083	Tenterfield-Dumaresq OS14	GDA	56	338395	6768085	Open site	Valid	Artefact : -		
	Contact	Recorders	ozA	rk Environm	ental and Herit	age Managemen	t,Mr.Ben Churcher	<u>Permits</u>		
11-3-0084	Tenterfield-Dumaresq OS15	GDA	56	342072	6768358	Open site	Valid	Artefact : -		
	Contact	Recorders	OzA	rk Environm	ental and Herit	age Managemen	t,Mr.Ben Churcher	<u>Permits</u>		
11-3-0093	Tenterfield Dumaresq OS 4 & PAD	GDA	56	342136	6768402	Open site	Valid	Artefact : -		
	Contact	Recorders	OzA	rk Environm	ental and Herit	age Managemen	t,Mr.Ben Churcher	<u>Permits</u>		



APPENDIX D BIODIVERSITY AND CONSERVATION DIVISION LETTER JUNE 2020

www.erm.com Version: 3.0 Project No.: 0464261 Client: GAIA Australia 19 June 2020



Our Ref: DOC20/426271 Your Ref: SSD 9438

Director Energy Assessments
Department of Planning, Industry and Environment
Locked Bag 5022
Parramatta NSW 2124

Attention: Mr Iwan Davies

Dear Ms Brewer

RE: Bonshaw Solar Farm - SSD9438 - Outstanding Aboriginal Cultural Heritage Issues

I am writing to provide you advice about the outstanding issues regarding the Aboriginal Cultural Heritage (ACH) assessment process carried out to inform the proposed Bonshaw Solar Farm project (SSD 9438).

The Biodiversity and Conservation Division (BCD) of the Environment, Energy and Science Group in the Department of Planning, Industry and Environment has been involved, in an advisory capacity, in assisting with the determination process for this project.

On 2 December 2019 we provided a letter outlining several recommendations we considered should be addressed prior to the determination of the development application for this project, including that a concerted effort be made to rediscover the previously recorded Aboriginal Heritage Information Management System (AHIMS) site 11-3-0083, which we considered highly important to the contextual understanding of the ACH values of the project area.

The ACH assessment for the project that we had reviewed stated that the consultant archaeologist had been unable to relocate the site in question and that the site had originally been identified as being of high significance. It was recorded as containing an Aboriginal scar tree, surface artefacts and likely subsurface deposits and representative of a much broader occupation complex likely to exist along the entire riparian corridor at this location.

On 21 February 2020 we received an updated Aboriginal Cultural Heritage Assessment Report in response to our submissions. Our review of that report noted that the issues we had raised about AHIMS 11-3-0038 had been overlooked.

On 21 May 2020 we participated in a teleconference with yourself and the Department's Mr Javier Canon, where it was agreed that our recommendation had not been addressed and that further discussions with the project archaeologist may be helpful in understanding the intensity and locations of the survey effort undertaken to relocate the site.

The BCD North East Branch Archaeologist has since made his own technical assessment of all the information recorded to date regarding AHIMS 11-3-0038. The result of his assessment suggests that technical errors on behalf of both the original recording archaeologist in 2011 and the current consultant archaeologists have resulted in a misunderstanding about the location of AHIMS 11-3-0038.

The original recording of the site, carried out in 2011, included coordinates, visual and textual information identifying the site location. After review of this information the BCD has determined the original coordinates of the site were recorded in one projection (AGD) but were mislabelled on the Aboriginal site recording form as another projection (GDA). This error would place the site approximately 210m southeast of its actual location on the ground. The error on the original recording form was translated into the AHIMS register by the Office of Environment and Heritage as per the form submitted.

When the consultant archaeologist for the project carried out their desktop analysis and field assessment, they failed to correlate the coordinates with the other information regarding the site location and therefore failed to identify the correct location of the site.

The BCD has reprojected the original coordinates into the GDA system and determined that the area identified in the 2011 survey and registered as AHIMS 11-3-0038 was identified during the most recent ACH survey and given a new site identifier as it was not recognised as the existing site. Figures 1 and 2 at Attachment 1 to this letter show the recorded location of AHIMS 11-3-0038 as previously understood and the actual location as determined by the BCD. Based on this information, we are confident that the original site, as recorded by OzArk in 2011, has been extensively surveyed as part of the most recent ACH Assessment for the Bonshaw Solar Farm.

However, this also means it now appears that a recorded Aboriginal scar tree may have been removed or destroyed in the last 10 years, which, if true, may constitute an offence under the *National Parks & Wildlife Act 1974*. Accordingly, the matter has been referred to the BCD North East Branch Compliance Team, which has advised that an investigation into the matter has commenced.

The BCD recommends that:

- 1. the consultant archaeologist must update the ACHAR for the project to reflect the information set out in this letter and include a map showing the re-projected location of AHIMS 11-3-0038.
- 2. conditions of approval should be included to:
 - a. require the preparation of an Aboriginal Cultural Heritage Management Plan in consultation with the project registered Aboriginal parties and the Senior Team Leader Planning, North East Branch, Biodiversity and Conservation Division, to the satisfaction of the Department, before ground-disturbing activities commence on the project.
 - b. set out a chance-finds procedure for Aboriginal Cultural Heritage.
 - c. require the identification of an appropriate area surrounding the re-projected location of AHIMS 11-3-0038 in consultation with the Senior Team Leader Planning, North East Branch, Biodiversity and Conservation Division, where there will be no impacts or management actions, to enable any compliance investigation considered necessary to proceed. Works can only commence in the identified area after written advice is received from the Biodiversity and Conservation Division to this effect.

The BCD has no further issues to raise about the level of ACH assessment for the project and would support a conditional approval of the development application as set out in our recommendations above. If you have any questions about this advice, please do not hesitate to contact Mr Roger Mehr, Archaeologist, at roger.mehr@environment.nsw.gov.au or 6773 7005.

Yours sincerely

11 June 2020


DIMITRI YOUNG


Senior Team Leader Planning, North East Branch

Biodiversity and Conservation

Enclosure: Attachment 1: Figures Demonstrating recorded and actual of AHIMS 11-3-0038

Attachment 1: Figures demonstrating recorded and actual of AHIMS 11-3-0038





NB: New AHIMS sites recorded by ACH survey for the project are shown in orange hatching – actual tree location is within northern-most survey site

ERM has over 160 offices across the following countries and territories worldwide

The Netherlands Argentina Australia New Zealand Belgium Norway Brazil Panama Canada Peru Chile Poland China Portugal Colombia Puerto Rico France Romania Germany Russia Hong Kong Singapore India South Africa Indonesia South Korea Ireland Spain Italy Sweden Japan Switzerland Kazakhstan Taiwan Kenya Thailand Malaysia UAE Mexico UK Mozambique US Myanmar Vietnam

ERM Sydney

Level 15, 309 Kent Street SYDNEY NSW 2000 Locked Bag 3012

AUSTRALIA SQUARE NSW 1215

T: +61 2 8584 8888 F: +61 2 8584 8800

www.erm.com

