

Dr Ian Baird  
3 Waimea St  
Katoomba NSW 2780  
02 47826755  
[ianbaird2@hotmail.net.au](mailto:ianbaird2@hotmail.net.au)

---

23 May 2014

Mining and Industry Projects  
NSW Department of Planning & Infrastructure  
GPO Box 39  
Sydney NSW 2001

## **Re. State Significant Development 5602**

### **Objection to proposed Angus Place Colliery Longwall Mine Extension Project, Newnes Plateau, NSW, by Centennial Angus Place Pty Ltd.**

I am writing to object to the proposed extension of the Angus Place Colliery longwall coal-mine on the Newnes Plateau by Centennial Angus Place Pty Ltd. This submission focuses specifically upon the likely impacts on endangered, groundwater dependent swamp ecosystems and their dependent species.

This project has the potential to lead to the irrevocable degradation and potential destruction of several Newnes Plateau Shrub Swamps (NPSS), an Endangered Ecological Community (EEC) (NSW Scientific Committee 2005b), which forms part of the Commonwealth listed Temperate Highland Peatswamps on Sandstone (THPSS) EEC (Threatened Species Scientific Committee 2005), with resultant negative impacts upon their associated groundwater dependent species. These species may include threatened flora and fauna, such as the endangered Giant Dragonfly (*Petalura gigantea*), endangered Blue Mountains Water Skink (*Eulamprus leuraensis*), and vulnerable Dean's Boronia (*Boronia deanei deanei*), all of which are obligate mire (peat swamp) dwelling species (Benson & Baird 2012). These species have been recorded from most NPSS (including new localities of *P. gigantea* and *E. leuraensis* recorded in January 2014 by Ian Baird) (see Atlas of NSW Wildlife; Baird 2012: Appendix 1; Benson & Baird 2012) and there are likely to be additional occurrences within the project area. These groundwater dependent swamp ecosystems (NSW Government 2002; Serov et al. 2012) are important habitat for these and other groundwater dependent species. Based upon extensive observation of the habitat of these species, and surveys of the three NPSS which occur above the proposed longwall panels, I can state that these swamps contain potential habitat for *P. gigantea* and *E. leuraensis*. Some patches of Newnes Plateau Hanging Swamps (NPHS) (DEC 2006), which also form part of the THPSS EEC, and which occur across the project area, may also provide habitat for these groundwater dependent species. My objection is grounded in over 10 years of research studying mire ecosystems, particularly in the Blue Mountains region, with a focus on *P. gigantea* (e.g., Baird 2012; Baird 2013; Baird in press; Baird & Burgin 2013; Benson & Baird 2012).

This submission includes a supporting literature review which focuses upon the specific threats to the groundwater dependent peat swamps and their associated groundwater dependent species (Part A), comments in relation to specific related issues identified in the EIS documentation (Part B) and recommendations to address identified issues (PART C).

## Part A: Literature Review

### Threats to groundwater dependent species

#### *Petalura gigantea*

Like most of the 11 species of petalurid dragonflies worldwide, the endangered (TSC Act 1995) *Petalura gigantea* (Petaluridae) is characterised by a burrowing larval lifestyle (Baird 2012: Chapter 1; in press; Tillyard 1911) that is unique in the Odonata (Corbet 1999). Studies to date suggest a larval stage of at least six years, and possibly more than 10 in *P. gigantea* (Baird 2012: Chapter 1). The long larval stage and the unpredictability of emergence events (Baird 2012: Chapter 7) means that the species is most unsuitable for the purpose of monitoring ecological changes associated with mining subsidence. Thus, any deleterious effects upon populations of the species, as a result of lowered water tables, will not be detectable until well after the event and thus too late.

Successful *P. gigantea* breeding sites are characterised by a groundwater regime that provides sufficient surface moisture to minimise risk of desiccation of eggs and early larval instars, supports development of organic-rich mire soils suitable for larval burrowing, and maintains a water table height that larvae can access within established burrows or adaptively through burrow deepening. Extensive burrow investigations and observation of hundreds of oviposition and burrow locations across a range of mire types in the Blue Mountains confirm *P. gigantea* as an obligate, groundwater dependent, mire-dwelling species (Baird 2012: Chapter 6). All investigated burrows included groundwater in the burrows, usually throughout the majority of the burrow depth (Baird 2012: Chapter 1; in press; Benson & Baird 2012). Although there is some evidence of putative adaptive burrow deepening by late stadia larvae in some circumstances, it is availability of suitable microhabitat for ovipositing and early larval establishment which is critical to persistence of populations of the species. All observed oviposition occurred into waterlogged substrate, fissures in the substrate, amongst or under moist litter overlying the substrate, amongst roots at the base of plants in moist substrate, or into *Sphagnum*. Breeding habitat for the species is restricted to the wetter parts of these swamp systems where the water table is either emergent or causes saturation of the surface of the substrate, at least throughout the period of egg-laying, early larval development and burrow establishment. This is the critical life history stage and is dependent upon the presence of saturated organic-rich substrates associated with high, and frequently, emergent water tables.

Any long-term lowering of the water table will result in a proportional contraction or reduction in the spatial extent of potential breeding habitat within individual swamps, and potentially, to extirpation of the species from those swamps, with potential implications for metapopulation dynamics of this patchily distributed species (Baird 2012: Chapter 8).

#### *Eulamprus leuraensis*

The endangered (TSC Act 1995, EPBC Act 1999) lizard, *Eulamprus leuraensis* (Scincidae), is endemic to mid to upper elevation Blue Mountains Sedge Swamps (of Keith & Benson 1988) and NPSS (Dubey & Shine 2010a, b; Dubey & Shine 2011; Dubey et al. 2013; LeBreton 1996). The species is closely associated with, and dependent upon, the wetter parts of these swamp systems, where there is typically moist to saturated substrates and at least some surface water associated with an emergent water table. It is known to use burrows of the crayfish *Euastacus australasiensis* (Decapoda: Parastacidae), and possibly also of *P. gigantea* during and post fire, as both fire and predation refugia (pers. obs.). Any long-term reduction in groundwater levels in individual swamps will negatively impact upon the species through a reduction in the area of core habitat and in the abundance of the species within affected swamps. Although individuals may be recorded in

adjoining non-swamp habitats, these areas do not represent core habitat of this obligate mire-dwelling species.

A recent genetic study has identified low rates of genetic exchange and high genetic divergence between discrete swamp populations, and recommended that most populations be treated separately as discrete conservation units (Dubey & Shine 2010b). Any loss of discrete swamp populations may contribute to eroding its genetic potential and capacity to adapt to projected climate change. Predicted contractions in the extent of the mire habitats of this species under projected climate change scenarios (see Ramp & Chapple 2010), and associated degradation in habitat value for the species, will further threaten the viability of populations of this species (Dubey & Shine 2011). These effects mirror, and would compound, those that will occur as a result of any lowering of water tables within these swamps as a result of subsidence associated with longwall mining. The distinctiveness of individual swamp populations is mirrored in Hose's (2009) work on genetics of stygofauna on the Woronora Plateau.

### ***Euastacus australasiensis***

The burrowing crayfish *Euastacus australasiensis* occurs in streams and upland swamps of the Sydney region (Growths & Marsden 1998; Merrick 1998; Morgan 1997) and can be abundant in the wetter parts of the swamps of the Newnes Plateau (pers. obs.); on current knowledge it is the only indigenous upland swamp dwelling crayfish within the Blue Mountains (see Merrick 1998; Morgan 1997).

Burrow investigations within Blue Mountains swamp systems have not been undertaken, but investigations of Tasmanian mire dwelling crayfish (*Parastacoides* spp.) have revealed complex burrow structures, groundwater dependence and unique functional roles (Growths & Richardson 1988; Richardson & Swain 1980, 1991). It is likely that *E. australasiensis* performs similar functional roles in NPSS. For example, the groundwater in its burrows may provide habitat for other specialised groundwater dependent fauna, in addition to providing refugia for *E. leuraensis* during and following fire events. The burrows of these crayfish can be referred to the Type 2 burrows (burrows connected to the water table) of the ecological classification system for burrows of Australian freshwater crayfish (Horwitz & Richardson 1986).

Studies of freshwater crayfish worldwide have revealed that their groundwater filled burrows may provide habitat for a diversity of organisms, including pholoteros (Lake 1977; Lake & Newcombe 1975), stygofauna (Gilbert et al. 1994a; Humphreys 2008), hibernating frogs (Irwin et al. 1999), aestivating commensal crayfish (Johnston & Robson 2009), and dragonfly larvae during drought and winter months (Harrington 2010; Pintor 2000). These diverse ecological relationships, and the shared reliance of different taxa upon groundwater, highlight the complexity of functional roles of these burrowing groundwater dependent organisms and their importance in these groundwater dependent ecosystems (GDEs). A lowering of water tables in these swamps can be expected to reduce and potentially eliminate the habitat available to *Euastacus* in affected swamps and negatively affect any associated species.

### **Additional groundwater dependent fauna**

Our knowledge of the groundwater dependent invertebrate fauna in Australian mire ecosystems is negligible. Additional groundwater dependent species within these Blue Mountains swamp ecosystems may include stygofauna (e.g., Gilbert et al. 1994b; Hose 2009; Humphreys 2008), pholoteros (e.g., Creaser 1931; Lake 1977; Lake & Coleman 1977; Lake & Newcombe 1975; Suter & Richardson 1977), and other specialised mire (e.g., Larson & House 1990; Rosenberg & Danks 1987; Spitzer & Jaroš 1993) or moorland (e.g., Butterfield & Coulson 1983; Gardner 1991; Greenslade &

Smith 1999) invertebrates. A lowering of water tables in any swamp is likely to have significant impacts upon any such groundwater dependent fauna, with possible extirpation from swamps where the water table is permanently lowered below a threshold level.

### ***Boronia deanei deanei***

The rare and vulnerable (TSC Act 1995) shrub, *Boronia deanei deanei* (Rutaceae), is restricted to NPSS and some bogs on the Boyd Plateau. It is associated with particular groundwater conditions and any lowering of water tables can be expected to reduce the distribution and abundance of the species in affected swamps, with implications for the persistence of the species more broadly (Benson & Baird 2012).

### ***Dillwynia stipulifera***

*Dillwynia stipulifera* (Fabaceae) is another small shrub restricted to swamps on the Newnes Plateau, with disjunct occurrences on the South Coast. Any contraction in its abundance or distribution on the Newnes Plateau will also have implications for its persistence in the area (Benson & Baird 2012). Benson and Baird (2012) have suggested that the species may qualify for listing as vulnerable under the TSC Act 1995.

## **Threats to peat swamps associated with longwall coal mining**

Subsidence associated with Longwall coal mining is listed as a Key Threatening Process (KTP) under the TSC Act 1995. This determination recognises that subsidence may cause deleterious changes to the quality and/or quantity of groundwater available to GDEs and surface streams. Mine dewatering may have similarly negative effects on surface ecosystems due to the alteration of surface water flow regimes and water quality. The impact of long wall coal mining, in particular, upon groundwater dependent swamp and stream ecosystems is of concern (NSW Scientific Committee 2005a; Young 1982; Young & Wray 2000) and controversy, consistent with the increasingly well-documented negative impacts of longwall coal mining worldwide (e.g., Bell et al. 2000; Booth 2006; Booth & Bertsch 1999; Booth et al. 1998; Karaman et al. 2001).

Extensive longwall coal mining under the Woronora Plateau and nearby areas in the Southern Coalfields has resulted in well documented cases of subsidence, with fracturing of stream beds and loss of groundwater from streams and endangered Coastal Upland Swamps. Threats from longwall mining have been specifically identified in respect of the EEC determinations for NPSS (NSW Scientific Committee 2005b) and Montane Peatlands and Swamps in NSW (NSW Scientific Committee 2004), which includes the Cops River Swamps (of Benson & Keith 1990). A large proportion of NPSS and Cops River Swamps occur above underground coal mining leases. This occurs particularly in Newnes and Ben Bullen state forests where the area has either been subjected to mining or such mining is proposed. Hydrological changes generally have been identified as threats in the EEC determinations for NPSS (NSW Scientific Committee 2005b) and THPSS (Threatened Species Scientific Committee 2005). Extensive mine de-watering into headwater streams and directly into swamp systems has occurred across this area, with significant and irreparable damage resulting. The well documented and irreparable damage to East Wolgan Swamp on the Newnes Plateau, resulting from loss of groundwater, associated with mine waste water discharge and subsidence following longwall coal mining by Centennial Coal, highlights the threat to all GDEs, including peat swamps, as a result of subsidence from long wall mining in this geological setting. 'Alteration to the natural flow regimes of rivers, streams, floodplains and wetlands' is also listed as a KTP on Schedule 3 of the TSC Act 1995 (NSW Scientific Committee 2002) and is applicable in the context of the potential loss of groundwater from these swamps and of base-flow to headwater streams.

Globally, threats to groundwaters and their dependent ecosystems are increasingly being recognised (Boulton 2005; Boulton et al. 2003; Danielopol et al. 2003; MacKay 2006). Hatton and Evans (1998) identified a range of ecosystem dependencies on groundwater. Their class of *Ecosystems with proportional dependence on groundwater* appears to include all mire ecosystems in Australia, including NPSS. For this class of ecosystem, they suggested that '*it is likely that a unit change in the amount of groundwater will result in a proportional change in the health or extent of that ecosystem*'. Commenting on the level of groundwater dependency of wetland ecosystems, Clifton and Evans (2001) highlighted the importance of maintaining adequate groundwater levels in unconfined aquifers and adequate groundwater discharge flux for most wetland ecosystems to maintain the necessary level of wetness or waterlogging for key ecological stages: '*Changes in water table level may have important implications for these communities. Prolonged lowering or raising of the water table is likely to result in changes in species composition, favouring species adapted to drier or wetter conditions, respectively*'.

Changes in groundwater regime will potentially impact the biota of GDEs at species and community level. This will be caused by changes to critical habitat attributes, either variably, or (potentially) across threshold levels (Clifton & Evans 2001). Where groundwater levels are lowered, even seasonally, effects may include changes in the wetland vegetation community composition (Boulton & Brock 1999; Breeuwer et al. 2009; Wheeler 1999). Although many of the plants which are obligate swamp dwellers in NPSS and other upland swamps may persist across a spectrum of the hydrological gradient, the effects of a lowering of the water table upon individual species may not be measurable for some time. This may be most pronounced for long-lived species such as Buttongrass, *Gymnoschoenus sphaerocephalus* (Cyperaceae) (pers. obs.). The effect upon groundwater fauna, however, will be more rapid. With longer term lowering of the water table, effects may include succession from swamps to drier heath, sedgeland, grassland or forest communities (as has occurred to a former swamp on Lamb's Creek located above longwall mining carried out by Angus Place Colliery); extirpation of groundwater dependent species; changes in groundwater quality and soil chemistry (Wheeler 1999); and degradation of peat (Shearer 1997) and peatlands (Moore 2002; Whittington & Price 2006). There may be increased competitive interactions for terrestrial swamp fauna (e.g., the swamp rat *Rattus lutreolus*, *Eulamprus leuraensis*) with otherwise generally allopatric congeners of adjoining non-mire habitats (Baird 2012). Such increased competitive interactions, in conjunction with reduced habitat suitability for the mire dwelling species, will further contribute to extirpation of populations of these species. Bioclimatic modelling indicates a progressive contraction in the extent of Blue Mountains swamps (Ramp & Chapple 2010), and swamps of the Woronora Plateau and the Sydney region more broadly (Keith et al. 2014; Keith et al. 2010), under projected climate change scenarios; these contractions mirror and would compound the effects that can be expected from any lowering of water tables in association with subsidence from longwall coal mining.

Additional potential impacts associated with a lowering of the water table include weed invasion and increased fire risk (Keith et al. 2006; Kodela et al. 2001; NSW Scientific Committee 2005a). Weed invasion of NPSS has been documented (Henson 2010) and anthropogenically disturbed and drying swamps, in particular, are most vulnerable to weed invasion. Where drying of organic-rich, peaty swamp substrates occur, fire effects may include burning of the organic soil component, resulting in the destruction of soil seed banks, rhizomes and lignotubers, thus also affecting re-sprouter species (Keith 1996). In the Blue Mountains region, including NPSS, substantial mortality of mature lignotubers of re-sprouter shrubs may occur as a result of fire during periods of reduced water levels in swamp soils (Benson & Baird 2012). Drying related oxidation and/or combustion of the organic component of these peatlands leads to their shrinkage and has the potential to leave sterilised and (often) hydrophobic sandy or peaty soils, with an increased risk of erosion due to surface water

flows. This, in turn, may lead to channelisation and further lowering of shallow water tables (e.g., Young & Wray 2000). Degradation of the hydrological function of these peatland ecosystems (capacity to store and slowly release water) will also occur (Keith et al. 2006; NSW Scientific Committee 2005a).

Fire is considered one of the main threats to Australian peatlands (Pemberton 2005). A more intense fire regime may cause unsustainable loss of peatland soils, as documented, for example, in moorland fires in the North York Moors (Maltby et al. 1990), for rare peatland in south-western Australia (Horwitz et al. 1999; Horwitz & Smith 2005; Semeniuk & Semeniuk 2005), in alpine peatland soils in Tasmania (Kirkpatrick & Dickinson 1984), in Buttongrass moorland (blanket bogs) in Tasmania (Bridle et al. 2003) and other organic terrains (peatswamps) in Tasmania and Victoria (Wein 1981). In addition to combustion of organic-rich peatland soils, fire induced changes include the potential for increased erosion, changes in water quality and loss of biodiversity (e.g., Horwitz & Sommer 2005). Fire effects are exacerbated where groundwater levels are lowered due to drought or anthropogenic influences (e.g., long wall mining) and by more intense fire regimes, for example, as reported also in New Zealand peatlands and bogs (Clarkson 1997; Johnson 2001; Timmins 1992). Similar effects have been reported for the upland swamps of the Woronora Plateau (Keith et al. 2006; Young 1982). Destruction of peat substrates as a result of fire in Blue Mountains Sedge Swamps has also been reported (Keith 1996; Stricker & Wall 1995). It may also lead to surface collapse over peatland piping, resulting in gully head initiation and channelisation (pers. obs.). An increase in periods where peatland soils are bare following fire events (or in response to drought or otherwise lowered water tables) will also contribute to increased rates of peatland photodegradation and loss of organic matter (Rutledge et al. 2010). A lowering of the groundwater table, for any reason, will compound these effects. The ecological damage resulting from any loss of groundwater from these swamp ecosystems as a result of subsidence from longwall mining will be compounded by subsequent long term effects of fire, and the potential compounding effects of predicted climate change (Baird 2012: Chapter 8; Benson & Baird 2012).

Fire is a natural and recurring event within the swamp communities in the Sydney region (Keith 1995; Keith et al. 2006). The temporal and spatial scales, and the intensity at which these fire disturbance events occur in particular swamp types are, however, critical to whether fire regimes result in long-term loss of organic terrains at a rate that exceeds net accumulation of organic matter, or exceeds fire regime thresholds for swamp plant (e.g., *Sphagnum* spp.) and animal species (e.g., *Rattus lutreolus*) (Keith 1996; Keith et al. 2002; Morrison 2002; Morrison et al. 1995; Watson 2006a, b). Fire impacts are directly correlated to swamp water levels and surface wetness (e.g., Horwitz & Smith 2005; Horwitz & Sommer 2005). Under a more intense fire regime, combined with reduced groundwater availability, long-term degradation and contraction of these ecosystems will occur. It will also threaten the persistence of groundwater dependent mire species such as *P. gigantea* and *E. leuraensis* (Baird 2012: Chapter 8; Benson & Baird 2012).

## Conclusion

Reduced groundwater availability as a result of subsidence from longwall mining will result in long-term degradation and contraction of these NPSS, compounding predicted effects of climate change. The result will be reduced spatio-temporal distribution of suitable breeding habitat for *Petalura gigantea*, and will threaten the persistence of other groundwater dependent species, including *Boronia deanei*, *Eulamprus leuraensis*, *Euastacus australasiensis*, and other organisms such as stygofauna. These unique and geographically restricted montane mire ecosystems must be protected. Maintenance of the necessary hydrological regime and groundwater levels will be fundamental to the persistence of these mire ecosystems and their groundwater dependent species.

In view of the extensive evidence of damage from subsidence associated with longwall coal mining in the western Blue Mountains, across the Sydney region more broadly, and internationally, the threat to these ecosystems and their dependent species from an expansion of longwall mining under the Newnes Plateau is too great. Assessing the likely impacts of this proposal must be informed by the Precautionary Principle and the substantial body of available evidence of its damaging effects.

## **Part B: Specific comments on EIS**

### **MAIN DOCUMENT Volume 1**

p.39 map (proposed longwall layout): Twin Gully Swamp, Tri Star Swamp, and Trail 6 swamp (NPSS) all contain some suitable habitat for *Petalura gigantea* (Pg) and may contain some core habitat for *Eulamprus leuraensis* (El).

p. 54, Figure 2.8: In the event that this proposal proceeds, LW1015 to LW1017 should be shortened in length to avoid Trail 6 Swamp completely as a precautionary measure, or further reduced in panel width, with increased pillar widths, to further minimise risk of subsidence. The longwall panels were similarly shortened to not go under Twin Gully Swamp and such shortening should also be applied to LW1004 to LW1006 in the case of Tri Star Swamp. If shortening is not feasible, then LW1004 to LW1006 should be further reduced in width, with increased pillar widths, to further minimise risk of subsidence. The proponent's justification for the currently proposed panel widths under these swamps is not substantiated.

p.97, s. 2.8.1: overstates the impact of selective logging across the project area.

p.98, s.2.8.2: fails to identify likely presence of El and Pg in the project area. Pg recorded by IRC Baird in Sunnyside Swamp. Trail 6 Swamp, Twin Gully Swamp and Tri Star Swamp contain some suitable habitat for Pg and El. Both species have been recorded in most NPSS in the Carne Creek catchment (see Atlas of NSW Wildlife; Baird 2012; Benson & Baird 2012)[Note that Volume 2, p.336, Table 10.9 does include record for El and identify Pg as potentially occurring; and Volume 2, p.350, s. 10.3.3.2 notes records for Pg in the project area].

p.101: Photograph 2.1.7 is not of a typical Newnes Plateau Shrub Swamp.

### **MAIN DOCUMENT Volume 2:**

p.352: states "Potential impacts to groundwater dependant ecosystems from longwall mining induced subsidence are the cracking of swamp bases causing consequent loss of water and the tilting and buckling of bedding planes and the ground surface. Cracks may divert some water temporarily from swamps, but will initially fill with water before eventually filling with silt/peat from within the swamp, so that there would be no long-term permanent impact on the groundwater level or flows in the swamp." There is no evidence on the Newnes Plateau or elsewhere in the Sydney Region that such natural filling of cracks with sediment will prevent long term damage to these swamps as a result of fracturing following subsidence or permit groundwater levels to return to pre-mining levels, in the medium to long term. Medium to long term lowering of water levels will result in either a reduction in the spatial extent of suitable habitat within affected swamps or extirpation of populations of various species, such as Pg and El, from affected swamps.

p.362, Table 10.14: This assessment is incorrect and flawed. Both Pg and El may be subject to local extirpation of individual swamp populations in the event of lowering of watertable in such swamps. The same applies to *Boronia deanei* ssp. *deanei* and El. Note also the work of Dubey and Shine (2010) which showed genetic diversity between El swamp populations, recommending they be treated as discrete conservation units, which has implications in terms of application of the 7 part test and consideration of likely impacts under the EPBC Act.

#### **APPENDIX G:**

p.viii: States "Remediation measures, such as grouting, in areas where fracturing of stream or swamp beds leads to protracted diversion of stream flow and/or drainage of pools". There is no evidence from the Sydney Basin that such grouting of fractured stream beds will achieve remediation objectives and return stream flows to a pre-mining condition; in fact, the opposite is true. The same applies in the case of fracturing of bedrock under swamps, where there is no evidence that this has ever been successfully implemented.

p.viii-ix: Discharge of what is effectively untreated, highly contaminated mine water to Kangaroo Creek via LDP001, and subsequently to the Coxs River, is totally inappropriate and the measures proposed to mitigate the ongoing and increasing damage to these aquatic ecosystems are inadequate. It is unfathomable that the company is currently permitted to degrade this stream system, let alone increase those negative effects; a complete redesign of the waste water management system, which pre-treats all mine waste water to a much higher level of water quality before discharge, is essential.

Emergency discharge points in the Wolgan River and Carne Creek must be eliminated and those discharge licences voided. The proponent must be held accountable and be responsible to ensure that their water management system is designed to cope with all scenarios and ensure that no waste water is ever transferred to these watercourses.

p.ix: States that any effects on potential populations of the endangered Adams Emerald Dragonfly will be insignificant. It is impossible to make this statement, considering that there have never been any ecological studies of this species and limited aquatic sampling was undertaken for this project. The assessment of no significant impact assumes that there will never be a discharge of contaminated water to a watercourse which may contain a population of this species or that there will never be a subsidence related, sustained loss of base flow to such a stream.

p.60-61: The consultant's recommendation to undertake more comprehensive, and better designed pre-mining surveying, and finer resolution taxonomic identification of stygofauna, must be implemented if this project proceeds to ensure that the diversity of stygofauna is properly assessed and potential risks of the project determined. The consultant later states that "Stygofauna associated with the perched surficial groundwater system could have been impacted by past mining-related activities, including discharge of mine water and construction of roads and access paths."

p. 76, s.7.4.2: I dispute the conclusion of the consultants that there will be no significant impact in relation to the identified Key Threatening Process (KTP) of subsidence associated with long wall coal mining. Their conclusion is based, in large part, upon the contentious conclusions of other consultants employed by Centennial Coal, in relation to the potential risk of subsidence induced lowering of water tables.



## APPENDIX H

p.iii: Pg and El have been recorded from NPSS immediately outside the project area in Sunnyside Swamp (Pg) and in most swamps cross the Carne Creek catchment. This should have been noted here, considering the project area includes potential swamp habitat for both species. It is possible that some swamps identified as NPHS in the project area may also provide some suitable habitat for these species.

p.iv: Fauna surveying was obviously inadequate, if they did not at least find swamp rats *Rattus lutreolus* (evidence observed by I.R.C. Baird in most NPSS) and *Antechinus* spp. in NPSS etc.

p.vi: As for Appendix G, the consultant here has based their conclusions on the contentious conclusions of other consultants, i.e., that there will be no significant impacts from subsidence upon GDEs and their associated species. These conclusions do not discount the possibility that such effects might occur, just that they are highly unlikely to occur.

p.vi: the consultant claims that there has been no significant damage to shrub swamps from subsidence, although the Main Report clearly identifies swamps significantly affected by mine subsidence, in addition to mine water discharge, e.g. East Wolgan Swamp.

p.vii: I strongly dispute the conclusion by the consultant that there will be no significant impact upon threatened species and EECs. They state that there is a “high level of confidence” that subsidence will not significantly affect threatened species such as the groundwater dependent Pg and El, and the NPSS and NPHS EECs. This is not good enough. Any risk is too much. A small, medium to long term lowering of the water table in a NPSS may result in a significant reduction in the potential habitat suitable for Pg and El, potentially leading to extirpation of one or both of these species from such a swamp. The consultant has demonstrated in their report a complete lack of understanding of the habitat requirements of both species which invalidates their conclusions. See notes above in relation to El and recommendation to treat each population as a discrete conservation unit (see Dubey & Shine 2010).

p.48: Photograph does not show a typical NPSS.

p.59: The Southern Emu-wren also occurs in NPSS and has been observed during fieldwork by I.R.C. Baird, including in Tri Star Swamp.

p.86: I reject the proposition that a lowering of water tables in swamps as a result of any subsidence will not negatively affect groundwater dependent species. Pg, for example, is reliant upon moist to waterlogged substrate for ovipositing and larval establishment (Baird 2012: Ch. 6). It only requires a drying of the surface peat to a shallow depth to reduce the spatial extent of potential breeding habitat in a particular swamp. The assumption of no significant impact by the consultant assumes that no significant subsidence and lowering of water tables will occur, which is contrary to the recognition elsewhere in the EIS that there will be some lowering of water tables in swamps.

p.94: Incorrectly states that since 2005 there has been no loss of EECs or populations of threatened species. This ignores East Wolgan Swamp, where subsidence is identified as a contributing factor to the damage to this NPSS, in addition to mine water discharge. To state that “the minor alterations to

the hydrological regime predicted are unlikely to modify the vegetation communities present in the short or long term” is totally unacceptable and inaccurate. “Unlikely” is not good enough. The previously undermined Junction Swamp, where EI was recorded in 2001, has been substantially degraded following a lowering of the water table in that swamp. EI has not subsequently been recorded there and the habitat may now be unsuitable for the species.

7 part tests of significance:

Assessment for *Boronia deanei* ssp. *deanei* is flawed. Any NPSS population of this species, which is subjected to medium to long term lowering of the water table, will be at risk of a reduction and potentially the loss of that population in the long term.

Assessment for *P. gigantea* is flawed and the consultant has misrepresented the information in the reference (Benson & Baird 2012) he cites. The requirement for moist to saturated substrate for successful oviposition and larval burrow establishment is the critical factor in the persistence of this species in swamps. Because of the long larval stage, probably in excess of 6 years, some late stadia larvae in established burrows may be able to persist until emergence, even after some lowering of water tables; however, successful reproduction, and thus persistence of populations, will be limited by the availability of suitable wet substrate for ovipositing and larval establishment (Baird 2012, in press; Benson & Baird 2013).

Assessment of *Eulamprus leuraensis* is flawed. The report states that EI also occurs in non-swamp habitat, however, any records for this species in non-swamp habitats are likely to be of wandering or foraging individuals temporarily outside their core swamp habitat. As above, the assessment of no significant impact upon this species is based on the flawed presumption that there will be no significant lowering of the water table in a particular swamp. A relatively minor loss of groundwater could have a deleterious effect on a local swamp population of this species.

Known and expected bird list: Southern Emu-wren has also been recorded in NPSS and was recently observed in Tri Star Swamp by I.R.C. Baird. This species favours swamp heaths and sedgelands.

## **PART C: Recommendations**

The mining footprint must be significantly lessened and mining methods reduced in intensity to protect the Wolgan River, Carne Creek, pagodas, cliffs and the nationally endangered swamps associated with these proposals. Centennial Coal must be required to consider alternative bord and pillar mining methods for its proposed Angus Place extension. Centennial’s Airly mine in the Capertee Valley operates to a depth of 405 metres underground in the same geology, with bad mine roof conditions, including many structural defects. If Centennial can operate Airly Colliery as a bord and pillar mine, then it can also operate the Angus Place mine in this manner.

The proposed Angus Place mine extension should not be granted development consent unless:

- Development consent is staged, with a review every five years;
- Staged approval with triggers requiring review of consent conditions should impacts be observed in the environmental matters of national and international significance;
- Subsequent approvals contingent on performance to consent conditions;

- Consent should be subject to performance standards triggers that ensure the health and integrity of receiving waters and heritage values;
- If the trigger levels are exceeded then consent should be immediately reviewed to address the failures at any time;
- Opportunity for third parties to provide effective input to avoid regulatory failure as State agencies are too busy to undertake the regulatory task and follow these issues in detail;
- A comprehensive, systematic pre-mining stygofauna survey must be implemented across the project area, with finer resolution taxonomic identification of stygofauna, to ensure that the diversity of stygofauna is properly assessed and potential risks of the project determined;
- No surface cracking of stream beds, under swamps or of pagodas, rock outcrops or cliffs;
- The intensity of longwall mining is reduced so that all nationally endangered swamps are protected – this includes shortening longwalls 1017 and 1016 to protect Trail 6 swamp and shortening longwalls 1004, 1005 and 1006 to protect the Tri Star Swamp complex;
- Narrowing and/or splitting the longwall panels 1007, 1008, 1009 and 1010 to prevent damage from fracturing in the Birds Rock Flora Reserve;
- Splitting longwalls 1013 and 1014 to prevent damage to pagodas and cliffs, as proposed;
- All proposed discharge of up to 43.8ML/day of mine effluent to the Cocks River via the Springvale-Delta Water Transfer Scheme (SDWTS) is treated by reverse osmosis technology to remove salt and metals to a standard that protects, the Cocks River, the downstream drinking water supply and near-pristine ecosystems in the World Heritage Area;
- Any malfunction of SDWTS, such as following a bushfire, must not result in emergency discharges to the World Heritage Area via Wolgan River or Carne Creek but be reinserted underground into the mine;
- Reinserted mine effluent must not then re-emerge in an unauthorised or unregulated manner but be properly treated;
- No emergence of near surface groundwater with elevated levels of salt or metal precipitate in Carne Creek;
- Representative sites for the piezometers to monitor groundwater in swamps and streams must be identified by a third party agency;
- Monitoring guidelines must clearly specify how the condition of groundwater dependent indicator plant species and the general condition of groundwater dependent ecosystems will be performed;
- All past tracks and trails created by Centennial Coal and its consultants, including those established by trail bikes, need to be recorded and plans set in place to rehabilitate these

trails on an on-going basis and as soon as practicable as part of the on-going rehabilitation program for this mine.

Subsidence monitoring should be by a third party agency, such as the Office of Environment and Heritage, and monitoring should be paid for by Centennial Coal.

Monitoring of surface flow and near-surface groundwater monitoring must create a comprehensive picture of the sub-catchments affected by mining.

Monitoring of changes in ecosystem condition must include well exposed, wide angle impacts of affected areas with GPS co-ordinates.

## References

- Baird I. R. C. (2012) The wetland habitats, biogeography and population dynamics of *Petalura gigantea* (Odonata: Petaluridae) in the Blue Mountains of New South Wales. PhD thesis, University of Western Sydney, Australia. Available from <http://handle.uws.edu.au:8081/1959.7/509925>.
- Baird I. R. C. (2013) Emergence behaviour in *Petalura gigantea* (Odonata: Petaluridae): confirmation of upright emergence. *International Journal of Odonatology* 16, 213-8.
- Baird I. R. C. (in press) Larval burrow morphology and groundwater dependence in a mire-dwelling dragonfly, *Petalura gigantea* (Odonata: Petaluridae). *International Journal of Odonatology*.
- Baird I. R. C. & Burgin S. (2013) An emergence study of *Petalura gigantea* (Odonata: Petaluridae). *International Journal of Odonatology* 16, 193-211.
- Bell F. G., Stacey T. R. & Genske D. D. (2000) Mining subsidence and its effect on the environment: some differing examples. *Environmental Geology* 40, 135-52.
- Benson D. & Baird I. R. C. (2012) Vegetation, fauna and groundwater interrelations in low nutrient temperate montane peat swamps in the upper Blue Mountains, New South Wales *Cunninghamia* 12, 267-307.
- Benson D. H. & Keith D. A. (1990) Natural vegetation of the Wallerawang 1:100,000 map sheet. *Cunninghamia* 2, 305-35.
- Booth C. J. (2006) Groundwater as an environmental constraint of longwall coal mining. *Environmental Geology* 49, 796-803.
- Booth C. J. & Bertsch L. P. (1999) Groundwater geochemistry in shallow aquifers above longwall mines in Illinois, USA. *Hydrogeology Journal* 7, 561-75.
- Booth C. J., Spande E. D., Pattee C. T., Miller J. D. & Bertsch L. P. (1998) Positive and negative impacts of longwall mine subsidence on a sandstone aquifer. *Environmental Geology* 34, 223-33.
- Boulton A. J. (2005) Chances and challenges in the conservation of groundwaters and their dependent ecosystems. *Aquatic Conservation: Marine and Freshwater Ecosystems* 15, 319-23.
- Boulton A. J. & Brock M. A. (1999) *Australian Freshwater Ecology: Processes and Management*. Gleneagles Publishing, Adelaide.

- Boulton A. J., Humphreys W. F. & Eberhard S. M. (2003) Imperilled subsurface waters in Australia: biodiversity, threatening processes and conservation. *Aquatic Ecosystem Health & Management* 6, 41-54.
- Breeuwer A., Robroek B. J. M., Limpens J., Heijmans M. M. P. D., Schouten M. G. C. & Berendse F. (2009) Decreased summer water table depth affects peatland vegetation. *Basic and Applied Ecology* 10, 330-9.
- Bridle K. L., Cullen P. & Russell M. (2003) *Peatland Hydrology, Fire Management and Holocene Fire Regimes in Southwest Tasmanian Blanket Bogs. Nature Conservation Report No. 03/07*. Nature Conservation Branch, Department of Primary Industries, Water and Environment, Hobart.
- Butterfield J. & Coulson J. C. (1983) The carabid communities on peat and upland grasslands in northern England. *Holarctic Ecology* 6, 163-74.
- Clarkson B. R. (1997) Vegetation recovery following fire in two Waikato peatlands at Whangamarino and Moanatuatua, New Zealand. *New Zealand Journal of Botany* 35, 167-79.
- Clifton C. & Evans R. (2001) *Environmental Water Requirements of Groundwater Dependent Ecosystems, Environmental Flows Initiative Technical Report Number 2*. Commonwealth of Australia, Canberra.
- Corbet P. S. (1999) *Dragonflies. Behaviour and Ecology of Odonata*. Cornell University Press, Ithaca, NY.
- Creaser E. P. (1931) Some cohabitants of burrowing crayfish. *Ecology* 12, 243-4.
- Danielopol D. L., Griebler C., Gunatilaka A. & Notenboom J. (2003) Present state and future prospects for groundwater ecosystems. *Environmental Conservation* 30, 104-30.
- DEC. (2006) The Vegetation of the Western Blue Mountains. Unpublished report funded by the Hawkesbury-Nepean Catchment Management Authority. Department of Environment and Conservation NSW, Hurstville.
- Dubey S. & Shine R. (2010a) Plio-Pleistocene diversification and genetic structure of an endangered lizard (the Blue Mountains water skink, *Eulamprus leuraensis*) in south-eastern Australia. *Journal of Biogeography* 37, 902-14.
- Dubey S. & Shine R. (2010b) Restricted dispersal and genetic diversity in populations of an endangered montane lizard (*Eulamprus leuraensis*, Scincidae). *Molecular Ecology* 19, 886-97.
- Dubey S. & Shine R. (2011) Predicting the effects of climate change on reproductive fitness of an endangered montane lizard, *Eulamprus leuraensis* (Scincidae). *Climatic Change* 107, 531-47.
- Dubey S., Sinsch U., Dehling M., Chevalley M. & Shine R. (2013) Population demography of an endangered lizard, the Blue Mountains Water Skink. *BMC Ecology* 13, 4.
- Gardner S. M. (1991) Ground beetle (Coleoptera: Carabidae) communities on upland heath and their association with heathland flora. *Journal of Biogeography* 18, 281-9.
- Gilbert J., Danielopol D. L. & Stanford J. A. (1994a) *Groundwater Ecology*. Academic Press, San Diego.

- Gilbert J., Stanford J. A., Dole-Olivier M.-J. & Ward J. V. (1994b) Basic attributes of groundwater ecosystems and prospects for research. In: *Groundwater Ecology* (eds J. Gilbert, D. L. Danielopol and J. A. Stanford), pp. 10-3. Academic Press, San Diego.
- Greenslade P. & Smith D. (1999) The epigaeic arthropod fauna of buttongrass moorland in Tasmanian Wilderness World Heritage Area. In: *The Other 99%: the Conservation and Biodiversity of Invertebrates* (eds W. F. Ponder and D. Lunney), pp. 90-4. Transactions of the Royal Zoological Society of New South Wales, Mosman, NSW.
- Growns I. O. & Marsden T. (1998) Altitude separation and pollution tolerance in the freshwater crayfish *Euastacus spinifer* and *E. australasiensis* (Decapoda: Parastacidae) in coastal flowing streams of the Blue Mountains, New South Wales. *Proceedings of the Linnean Society of NSW* 120, 139-45.
- Growns I. O. & Richardson A. M. M. (1988) Diet and burrowing habits of the freshwater crayfish, *Parastacoides tasmanicus tasmanicus* Clark (Decapoda : Parastacidae). *Marine and Freshwater Research* 39, 525-34.
- Harrington C. (2010) Return of the Dragon. A lost species reveals its secrets on a Wisconsin preserve. *Nature Conservancy* Summer 2010, 16.
- Hatton T. & Evans R. (1998) *Dependence of Ecosystems on Groundwater and its Significance to Australia*. LWRRDC Occasional Paper No. 12/98. Land and Water Resources Research & Development Corporation, Canberra.
- Henson M. (2010) Newnes Plateau Shrub Swamp Aerial Condition Assessment Report. Unpublished report. Save our Swamps project funded by Caring For Country and NSW Environmental Trust.
- Horwitz P., Pemberton M. & Ryder D. (1999) Catastrophic loss of organic carbon from a management fire in a peatland in southwestern Australia. In: *Wetlands for the Future* (eds A. J. McComb and J. A. Davis), pp. 487-501. Gleneagles Publishing, Adelaide, South Australia.
- Horwitz P. & Smith R. (2005) Fire and wetland soils and sediments on the Swan Coastal Plain: an introduction. *Journal of the Royal Society of Western Australia* 88, 77-9.
- Horwitz P. & Sommer B. (2005) Water quality responses to fire, with particular reference to organic-rich wetlands and the Swan Coastal Plain: a review. *Journal of the Royal Society of Western Australia* 88, 121-8.
- Horwitz P. H. J. & Richardson A. M. M. (1986) An ecological classification of the burrows of Australian freshwater crayfish. *Marine and Freshwater Research* 37, 237-42.
- Hose G. (2009) Stygofauna baseline assessment for Kangaloon borefield investigations – Southern Highlands, NSW. Supplementary report – stygofauna molecular studies. Report to Sydney Catchment Authority Access Macquarie Ltd, North Ryde.
- Humphreys W. F. (2008) Rising from down under: developments in subterranean biodiversity in Australia from a groundwater fauna perspective. *Invertebrate Systematics* 22, 85-101.
- Irwin J. T., Costanzo J. P. & Lee Jr R. E. (1999) Terrestrial hibernation in the northern cricket frog, *Acris crepitans*. *Canadian Journal of Zoology* 77, 1240-6.

- Johnson P. N. (2001) Vegetation recovery after fire on a southern New Zealand peatland. *New Zealand Journal of Botany* 39, 251-67.
- Johnston K. & Robson B. J. (2009) Commensalism used by freshwater crayfish species to survive drying in seasonal habitats. *Invertebrate Biology* 128, 269-75.
- Karaman A., Carpenter P. & Booth C. (2001) Type-curve analysis of water-level changes induced by a longwall mine. *Environmental Geology* 40, 897-901.
- Keith D. A. (1995) Mosaics in Sydney heathland vegetation: the roles of fire, competition and soils. *CALM Science Supplement* 4, 199-206.
- Keith D. A. (1996) Fire-driven extinction of plant populations: a synthesis of theory and review of evidence from Australian vegetation. *Proceedings of the Linnean Society of NSW* 116, 37-78.
- Keith D. A. & Benson D. H. (1988) Natural vegetation of the Katoomba 1:100,000 map sheet. *Cunninghamia* 2, 107-43.
- Keith D. A., Elith J. & Simpson C. C. (2014) Predicting distribution changes of a mire ecosystem under future climates. *Diversity and Distributions* 20, 440-54.
- Keith D. A., McGaw L. & Whelan R. J. (2002) Fire regimes in Australian heathlands and their effects on plants and animals. In: *Flammable Australia: the Fire Regimes and Biodiversity of a Continent* (eds R. A. Bradstock, J. E. Williams and A. M. Gill). Cambridge University Press, Cambridge.
- Keith D. A., Rodoreda S. & Bedward M. (2010) Decadal change in wetland - woodland boundaries during the late 20th century reflects climatic trends. *Global Change Biology* 16, 2300-6.
- Keith D. A., Rodoreda S., Holman L. & Lemmon J. (2006) Monitoring change in upland swamps in Sydney's water catchments: the roles of fire and rain. Sydney Catchment Authority Special Area Strategic Management Research and Data Program. Project number RD07: Long term responses of upland swamps to fire. Final Report. Department of Environment and Conservation, Hurstville, NSW.
- Kirkpatrick J. B. & Dickinson K. J. M. (1984) The impact of fire on Tasmanian alpine vegetation and soils. *Australian Journal of Botany* 32, 613-29.
- Kodala P. G., Sainty G. R., Bravo F. J. & James T. A. (2001) Wingecarribee Swamp flora survey and related management issues. Unpublished report to Sydney Catchment Authority, New South Wales.
- Lake P. S. (1977) Pholoteros - the faunal assemblage found in crayfish burrows. *Australian Society of Limnology Newsletter* 15, 57-60.
- Lake P. S. & Coleman D. J. (1977) On the subterranean syncarids of Tasmania. *Helictite* 15, 12-7.
- Lake P. S. & Newcombe K. J. (1975) Observations on the ecology of the crayfish *Parastacoides tasmanicus* (Decapoda: Parastacidae) from south-western Tasmania. *Australian Journal of Zoology* 18, 197-214.
- Larson D. J. & House N. L. (1990) Insect communities of Newfoundland bog pools with emphasis on the Odonata. *Canadian Entomologist* 122, 469-501.

- LeBreton M. (1996) Habitat and distribution of the Blue Mountains swamp skink (*Eulamprus leuraensis*). B. Zool. (Honours) thesis, University of New South Wales.
- MacKay H. (2006) Protection and management of groundwater-dependent ecosystems: emerging challenges and potential approaches for policy and management. *Australian Journal of Botany* 54, 231-7.
- Maltby E., Legg C. J. & Proctor M. C. F. (1990) The ecology of severe moorland fire on the North York Moors: effects of the 1976 fires, and subsequent surface and vegetation development. *The Journal of Ecology* 78, 490-518.
- Merrick J. R. (1998) *Endemic Crayfishes of the Sydney Region: Distribution, Biology and Management Options*. Macquarie University, North Ryde, NSW.
- Moore P. D. (2002) The future of cool temperate bogs. *Environmental Conservation* 29, 3-20.
- Morgan G. J. (1997) Freshwater crayfish of the genus *Euastacus* Clark (Decapoda: Parastacidae) from New South Wales, with a key to all species of the genus. *Records of the Australian Museum Supplement* 23 (1997), 110 pp.
- Morrison D. A. (2002) Effects of fire intensity on plant species composition of sandstone communities in the Sydney region. *Austral Ecology* 27, 433-41.
- Morrison D. A., Cary G. J., Pengelly S. M., Ross D. G., Mullins B. J., Thomas C. R. & Anderson T. S. (1995) Effects of fire frequency on plant species composition of sandstone communities in the Sydney region: inter-fire interval and time-since-fire. *Australian Journal of Ecology* 20, 239-47.
- NSW Government. (2002) *The NSW State Groundwater Dependent Ecosystems Policy*. Department of Land and Water Conservation and the State Groundwater Policy Working Group, Sydney.
- NSW Scientific Committee. (2002) Alteration to the natural flow regimes of rivers, streams, floodplains and wetlands. NSW Scientific Committee Key Threatening Process final determination. [Accessed December 10th 2009]. Available from <http://www.environment.nsw.gov.au/threatenedspecies/AlterationNaturalFlowKTPListing.htm>
- NSW Scientific Committee. (2004) Montane Peatlands and Swamps of the New England Tableland, NSW North Coast, Sydney Basin, South East Corner, South Eastern Highlands and Australian Alps bioregions - endangered ecological community listing. NSW Scientific Committee final determination. [Accessed August 13th 2008]. Available from <http://www.environment.nsw.gov.au/determinations/MontanePeatlandsEndSpListing.htm>
- NSW Scientific Committee. (2005a) Alteration of habitat following subsidence due to longwall mining. NSW Scientific Committee Key Threatening Process final determination. [Accessed May 29th 2005]. Available from <http://www.environment.nsw.gov.au/determinations/LongwallMiningKtp.htm>
- NSW Scientific Committee. (2005b) Newnes Plateau Shrub Swamp in the Sydney Basin Bioregion - endangered ecological community listing. NSW Scientific Committee final determination. [Accessed August 13th 2008]. Available from <http://www.environment.nsw.gov.au/determinations/NewnesPlateauShrubSwampEndSpListing.htm>
- Pemberton M. (2005) Australian peatlands: a brief consideration of their origin, distribution, natural values and threats. *Journal of the Royal Society of Western Australia* 88, 81-9.



Pintor L. (2000) The effect of drought on species interactions, behavior and habitat use by the Hine's emerald dragonfly, *Somatochlora hineana*. MSc Thesis. University of Illinois, Urbana-Champaign.

Ramp D. & Chapple R. (2010) Managing for Ecosystem Change in the GBMWhA: Final Report. ARC Project LP774833. University of NSW and Blue Mountains World Heritage Institute.

Richardson A. M. M. & Swain R. (1980) Habitat requirements and distribution of *Engaeus cisternarius* and three subspecies of *Parastacoides tasmanicus* (Decapoda: Parastacidae), burrowing crayfish from an area of south-western Tasmania. *Australian Journal of Marine and Freshwater Research* 31, 475-84.

Richardson A. M. M. & Swain R. (1991) Pattern and persistence in the burrows of two species of the freshwater crayfish, *Parastacoides* (Decapoda: Parastacidae), in southwest Tasmania. *Memoirs of the Queensland Museum* 31, 283.

Rosenberg D. M. & Danks H. V. (1987) Aquatic insects of peatlands and marshes in Canada. *Memoirs of the Entomological Society of Canada* 140, 174pp.

Rutledge S., Campbell D. I., Baldocchi D. & Schipper L. A. (2010) Photodegradation leads to increased carbon dioxide losses from terrestrial organic matter. *Global Change Biology* 16, 3065-74.

Semeniuk V. & Semeniuk C. A. (2005) Wetland sediments and soils on the Swan Coastal Plain, southwestern Australia: types, distribution, susceptibility to combustion, and implications for fire management. *Journal of the Royal Society of Western Australia* 88, 91-120.

Serov P., Kuginis L. & Williams J. P. (2012) *Risk assessment guidelines for groundwater dependent ecosystems, Volume 1 - the conceptual framework*. Department of Primary Industries, NSW Office of Water, Sydney.

Shearer J. C. (1997) Natural and anthropogenic influences on peat development in Waikato/Hauraki Plains restiad bogs. *Journal of the Royal Society of New Zealand* 27, 295-313.

Spitzer K. & Jaroš J. (1993) Lepidoptera associated with the Cervene Blato Bog (Central Europe): conservation implications. *European Journal of Entomology* 90, 323-36.

Stricker J. S. & Wall C. A. (1995) Wetlands of the Nepean – Hawkesbury Catchment. Sydney Water Corporation, Sydney.

Suter P. J. & Richardson A. M. M. (1977) The biology of two species of *Engaeus* (Decapoda : Parastacidae) in Tasmania. III. Habitat, food, associated fauna and distribution. *Marine and Freshwater Research* 28, 95-103.

Threatened Species Scientific Committee. (2005) Commonwealth listing advice on Temperate Highlands Peat Swamps on Sandstone. [Accessed August 12th 2008]. Available from <http://www.environment.gov.au/biodiversity/threatened/communities/temperate-highland-peat-swamps.html>

Tillyard R. J. (1911) Studies in the life-histories of Australian Odonata. 4. Further notes on the life-history of *Petalura gigantea* Leach. *Proceedings of the Linnean Society of NSW* 36, 86-96.

Timmins S. M. (1992) Wetland vegetation recovery after fire: Eweburn Bog, Te Anau, New Zealand. *New Zealand Journal of Botany* 30, 383-99.

Watson P. (2006a) *Hotspots Fire Project: Fire and the Vegetation of the Southern Rivers Region*. Nature Conservation Council of NSW and the NSW Government Environmental Trust, Sydney.

Watson P. (2006b) *Hotspots Fire Project: Fire Frequency Guidelines and the Vegetation of the Northern Rivers Region*. Nature Conservation Council of NSW and the NSW Government Environmental Trust, Sydney.

Wein R. W. (1981) Characteristics and suppression of fires in organic terrain in Australia. *Australian Forestry* 44, 162-9.

Wheeler B. D. (1999) Water and plants in freshwater wetlands. In: *Eco-hydrology: Plants and Water in Terrestrial and Aquatic Environments* (eds A. J. Baird and R. L. Wilby), pp. 127-80. Routledge, London.

Whittington P. N. & Price J. S. (2006) The effects of water table draw-down (as a surrogate for climate change) on the hydrology of a fen peatland, Canada. *Hydrological Processes* 20, 3589-600.

Young A. R. M. (1982) Upland swamps (dells) on the Woronora Plateau. PhD thesis. University of Wollongong.

Young R. W. & Wray R. A. L. (2000) The geomorphology of sandstones in the Sydney region. In: *Sandstone City: Sydney's Dimension Sandstone and other Sandstone Geomaterials: Proceedings of a Symposium held on 7 July 2000* (eds G. H. McNally and B. J. Franklin). Geological Society of Australia, Springwood.