A decorative background on the left side of the page featuring a topographic map with contour lines in a light grey color.

Normanhurst Loreto – Revised SSD application - Concept and envelopes Biodiversity Development Assessment Report

**Loreto Normanhurst c/o Carmichael Tompkins Property
Group**

DOCUMENT TRACKING

| | |
|------------------------|---|
| Project Name | Loreto Normanhurst Biodiversity Development Assessment Report |
| Project Number | 20SYD_17325 |
| Project Manager | Belinda Failes |
| Prepared by | Belinda Failes |
| Reviewed by | Nicole McVicar and Jennie Powell |
| Approved by | Nicole McVicar |
| Status | Final |
| Version Number | 6 |
| Last saved on | 16 December 2020 |

This report should be cited as 'Eco Logical Australia. 2020 Loreto Normanhurst Biodiversity Development Assessment Report. State Significant Development Resubmission Masterplan and Stage 1. Prepared for Loreto Normanhurst.'

ACKNOWLEDGEMENTS

This document has been prepared by Eco Logical Australia Pty Ltd with support from Dua Green from Allen Jack + Cottier

Disclaimer

This document may only be used for the purpose for which it was commissioned and in accordance with the contract between Eco Logical Australia Pty Ltd and Loreto Normanhurst. The scope of services was defined in consultation with Loreto Normanhurst, by time and budgetary constraints imposed by the client, and the availability of reports and other data on the subject area. Changes to available information, legislation and schedules are made on an ongoing basis and readers should obtain up to date information. Eco Logical Australia Pty Ltd accepts no liability or responsibility whatsoever for or in respect of any use of or reliance upon this report and its supporting material by any third party. Information provided is not intended to be a substitute for site specific assessment or legal advice in relation to any matter. Unauthorised use of this report in any form is prohibited.

Template 2.8.1

Executive Summary

Eco Logical Australia Pty Ltd (ELA) was engaged by Loreto Normanhurst on behalf of Carmichael Tompkins Property Group to update the existing Biodiversity Development Assessment Report (BDAR) in response to changes for the updated Masterplan and Stage 1 development at 91 - 93 Pennant Hills Road, Normanhurst, NSW.

Stage 1 will involve the construction of a new boarding house facility, a link road and carparking facilities. The proposed development is classified as State Significant Development and requires approval under State Significant Development Application SSD 17_8996 and is required to be assessed by the *Department of Planning and Environment* (DP&E). The preparation of a BDAR is a requirement of the Secretary's Environmental Assessment Requirements (SEARs).

The Loreto Normanhurst site (referred to as the 'subject site') covers an area of 13.14 ha. The Stage 1 and Concept Plan works (i.e. the 'development footprint') covers an area of 4.34 ha and the proposed development will result in the removal of 0.31 ha of non-native planted vegetation, 0.35 ha of native planted vegetation, and 0.05 ha of remnant non-planted vegetation. The remaining 3.62 ha represents buildings and cleared areas. The subject site is zoned R2 Low Density Residential under the *Hornsby Shire Council Local Environmental Plan 2013* (HLEP) which allows for residential development and educational establishments with council development consent.

The proposed development will impact upon biodiversity values within the subject site and as such a BDAR is required to assess the vegetation clearing under the NSW *Biodiversity Conservation Act 2016* (BC Act) in accordance with the SEARs. This report has been prepared to meet the requirements of the Biodiversity Assessment Method 2016 (BAM) established under Section 6.7 of the NSW BC Act 2016. Requirements of the HLEP and *Hornsby Development Control Plan 2013* have also been addressed in this report.

The vegetation within the subject site is highly disturbed with a mixed canopy of mature exotic and native planted species and some remnant regrowth canopy species which have been incorporated into horticultural landscape gardens. Although the vegetation in the subject site is highly modified and has been planted, under the BAM 2017 all vegetation native to NSW must be assigned a Plant Community Type (PCT) in accordance with the NSW Government's BioNet Vegetation Classification System. Where native vegetation has been planted and does not clearly confirm to any PCT, a 'best-fit' PCT must be assigned. Based on Office of Environment and Heritage (OEH) 2016 vegetation mapping, soil mapping and field validation of regrowth remnant vegetation retained within the subject site, it has been determined that the planted native vegetation conforms to PCT 1237 *Blue Gum High Forest*. PCT 1237 was also represented as an isolated remnant tree of which 0.001 ha will require minor trimming of trees as part of the development works. Although components of PCT 1237 corresponds to Blue Gum High Forest which is listed as a threatened ecological community (TEC), only the small amount of remnant vegetation (i.e. not planted) present within the subject site was considered part of the TEC under the NSW BC Act and none of the patches satisfied the criteria for listing under the *Commonwealth Environment Protection and Biodiversity Conservation Act 1999* (EPBC Act).

A second PCT, PCT 1281 *Sydney Turpentine-Ironbark Forest* was mapped in the southern section of the subject site and contains remnant regrowth vegetation in good condition and disturbed/weedy

condition. This patch of bushland has also been mapped on the NSW Government Biodiversity Values Map and Hornsby Council Terrestrial Biodiversity layer in the HLEP. PCT 1281 correspond to Sydney Turpentine-Ironbark Forest which is listed as critically endangered under the EPBC Act and endangered under the BC Act. Blue Gum High Forest (BGHF) and Sydney Turpentine-Ironbark Forest (STIF) are both listed Serious and Irreversible Impacts (SAIL) candidate entities and are impacted by this proposal. Consideration of SAIL candidates have been assessed as part of this BDAR.

Despite the presence of native vegetation, no threatened flora or fauna species were recorded within the subject site. Three widely cultivated species which also represents threatened entity listed under the BC Act and/or EPBC Act were located in landscaped gardens. This includes *Grevillea juniperina*, *Eucalyptus scoparia* (Wallangarra White Gum) and *Syzygium paniculatum* (Magenta Lilly Pilly). In accordance with the BAM 2020 Appendix D it was determined that these species were not planted for rehabilitation or recovery purposes and therefore, do not require additional assessment under the BC Act. An assessment under the EPBC Act was conducted for *Eucalyptus scoparia* (Wallangarra White Gum) and *Syzygium paniculatum* (Magenta Lilly Pilly)

Targeted surveys were conducted to determine if threatened microchiropteran (microbat) species utilise the roof cavity of a historic building to be demolished for Stage 1 works. Some microbat species are considered dual ecosystem credit and species credit species; however, no species credit species habitat was identified as potential habitat within the subject site. However, the BAM also requires targeted surveys as part of assessing Prescribed Impacts, this includes man-made structures. There was no evidence of microbat species recorded during targeted surveys.

This BDAR outlines the measures taken to avoid, minimise and mitigate impacts to the vegetation and species habitat present within the subject site and methodologies to minimise impacts during construction and operation of the development. Following consideration of all the above aspects, the residual unavoidable impacts of the project were calculated in accordance with the BAM by utilising the Biodiversity Assessment Method Credit calculator (BAMC). Under the BAM, seven (7) ecosystem credits are required to offset the removal of 0.35 ha of PCT 1237 (integrity score of 35.1) and one (1) ecosystem credits for PCT 1281 for the removal of 0.05 ha (integrity score of 39.7).

One Matter of National Environmental Significance (MNES) was identified as having potential to be adversely affected by the proposed works. *Pteropus poliocephalus* (Grey-headed Flying-fox) is listed as Vulnerable under the EPBC Act and it is considered that this species is likely to use some of the subject site for foraging. Two planted threatened species listed under the EPBC Act, *Eucalyptus scoparia* (Wallangarra White Gum) and *Syzygium paniculatum* (Magenta Lilly Pilly) were recorded within the subject site but will not be impacted. An assessment of the Commonwealth Significant Impact Criteria was undertaken for the Grey-headed Flying-fox, *Eucalyptus scoparia* and *Syzygium paniculatum* and concluded that the project would not have a significant impact on these species. As such, a referral to the Commonwealth is not required.

.

Contents

| | |
|--|-----------|
| 1. State Significant Development Application (Updated Masterplan and Stage 1 Works) | 2 |
| 1.1 Introduction..... | 2 |
| 1.2 Background..... | 2 |
| 1.2.1 Need for a Campus Masterplan | 2 |
| 1.2.2 Early Learning Centre | 2 |
| 1.3 The Site | 2 |
| 1.3.1 Legal description and ownership | 3 |
| 1.4 Overview of proposed development | 3 |
| 2. Stage 1: Biodiversity assessment..... | 5 |
| 2.1 Introduction..... | 5 |
| 2.1.1 General description of the subject site..... | 5 |
| 2.1.2 Subject site footprint | 5 |
| 2.1.3 Sources of information used..... | 6 |
| 2.2 Legislative context | 10 |
| 2.3 Landscape features..... | 11 |
| 2.3.1 IBRA regions and subregions | 11 |
| 2.3.2 Mitchell Landscapes..... | 11 |
| 2.3.3 Rivers and streams..... | 11 |
| 2.3.4 Wetlands..... | 11 |
| 2.3.5 Connectivity features..... | 11 |
| 2.3.6 Areas of geological significance and soil hazard features..... | 12 |
| 2.3.7 Site context | 12 |
| 2.4 Native vegetation | 12 |
| 2.4.1 Survey effort | 12 |
| 2.4.2 Plant Community Types present..... | 13 |
| 2.4.3 Vegetation integrity assessment | 16 |
| 2.5 Threatened species..... | 20 |
| 2.5.1 Threatened flora species | 20 |
| 2.5.2 Ecosystem credit species | 23 |
| 2.6 Species credit species | 29 |
| 2.6.1 Targeted surveys..... | 40 |
| 2.6.2 Use of local data | 40 |
| 2.6.3 Expert reports | 40 |
| 3. Stage 2: Impact assessment (biodiversity values)..... | 41 |
| 3.1 Avoiding impacts..... | 41 |

| | |
|--|-----------|
| 3.1.1 Locating a project to avoid and minimise impacts on vegetation and habitat..... | 41 |
| 3.1.2 Designing a project to avoid and minimise impacts on vegetation and habitat..... | 42 |
| 3.1.3 Prescribed biodiversity impacts | 43 |
| 3.2 Assessment of Impacts | 47 |
| 3.2.1 Direct impacts | 47 |
| 3.2.2 Change in vegetation integrity..... | 48 |
| 3.2.3 Indirect impacts | 48 |
| 3.2.4 Prescribed biodiversity impacts..... | 51 |
| 3.2.5 Mitigating and managing impacts..... | 55 |
| 3.2.6 Serious and Irreversible Impacts (SAII) | 58 |
| 3.3 Risk assessment | 61 |
| 3.4 Adaptive management strategy | 63 |
| 3.5 Impact summary | 65 |
| 3.5.1 Serious and Irreversible Impacts (SAII) | 65 |
| 3.5.2 Impacts requiring offsets | 65 |
| 3.5.3 Areas not requiring assessment..... | 65 |
| 3.5.4 Credit summary | 66 |
| 3.6 Consistency with legislation and policy | 66 |
| 3.6.1 <i>Environment Protection and Biodiversity Conservation Act 1999</i> (EPBC Act)..... | 66 |
| 3.6.2 <i>Hornsby Council Local Environmental Plan 2013 (HLEP)</i> | 71 |
| 3.7 Offset options | 73 |
| 4. References | 78 |
| Appendix A: Definitions..... | 80 |
| Appendix B: Vegetation plot data | 83 |
| Appendix C: Photos | 87 |
| Appendix D: Other species recorded | 89 |
| Appendix E: Floristic analysis | 95 |
| Appendix F: Anabat results | 96 |
| Appendix G: Biodiversity credit report | 97 |

List of Figures

| | |
|--|----|
| Figure 1: Masterplan and Stage 1 works..... | 7 |
| Figure 2: Site Map | 8 |
| Figure 3: Location Map..... | 9 |
| Figure 4: Plant Community Types and native vegetation extent..... | 17 |
| Figure 5: Plot locations..... | 18 |
| Figure 6: Threatened Ecological Communities..... | 19 |
| Figure 7: Final project footprint including construction and operation | 64 |
| Figure 8: Serious and Irreversible Impacts..... | 74 |

| | |
|---|----|
| Figure 9: Impacts requiring offset | 75 |
| Figure 10: Areas not requiring assessment | 76 |

List of Tables

| | |
|---|----|
| Table 1: Legal Description | 3 |
| Table 2: Legislative context | 10 |
| Table 3: Mitchell Landscapes | 11 |
| Table 4: Rivers and streams | 11 |
| Table 5: Connectivity features | 12 |
| Table 6: Full floristic and vegetation integrity plots..... | 13 |
| Table 7: Plant Community Types..... | 13 |
| Table 8: Threatened Ecological Communities..... | 14 |
| Table 9: PCT selection justification | 14 |
| Table 10: Vegetation integrity..... | 16 |
| Table 11: Decision making tool for Planted Native Vegetation in accordance with Appendix D of the BAM 2020..... | 21 |
| Table 11: Predicted ecosystem credit species | 23 |
| Table 12: Candidate species credit species..... | 29 |
| Table 13: Locating a project to avoid and minimise impacts on vegetation and habitat | 41 |
| Table 14: Designing a project to avoid and minimise impacts on vegetation and habitat..... | 42 |
| Table 15: Weather conditions during anabat targeted survey | 44 |
| Table 16: Prescribed biodiversity impacts | 45 |
| Table 17: Locating a project to avoid and minimise prescribed biodiversity impacts | 46 |
| Table 18: Designing a project to avoid and minimise prescribed biodiversity impacts..... | 47 |
| Table 19: Direct impacts to native vegetation | 48 |
| Table 20: Direct impacts on threatened ecological communities | 48 |
| Table 21: Change in vegetation integrity | 48 |
| Table 22: Indirect impacts..... | 49 |
| Table 23: Direct impacts on prescribed biodiversity impacts | 51 |
| Table 24: Measures proposed to mitigate and manage impacts..... | 55 |
| Table 25: Candidate Serious and Irreversible Impacts..... | 58 |
| Table 26: Determining whether impacts are serious and irreversible..... | 58 |
| Table 27: Evaluation of an impact on a TEC – Blue Gum High Forest..... | 59 |
| Table 28: Evaluation of an impact on a TEC Sydney Turpentine Ironbark Forest..... | 60 |
| Table 29: Likelihood criteria | 61 |
| Table 30: Consequence criteria..... | 62 |
| Table 31: Risk matrix | 62 |
| Table 32: Risk assessment..... | 62 |
| Table 33: Serious and Irreversible Impacts Summary..... | 65 |
| Table 34: Impacts to native vegetation that require offsets | 65 |
| Table 35: Ecosystem credits required | 66 |
| Table 36: EPBC Act of Significance for <i>Pteropus poliocephalus</i> (Grey-headed Flying-fox) | 67 |

| | |
|---|----|
| Table 37: EPBC Act of Significance for <i>Eucalyptus scoparia</i> | 69 |
| Table 38: EPBC Act of Significance for <i>Syzygium paniculatum</i> | 70 |
| Table 39: Clause 6.4 of the HLEP..... | 72 |
| Table 40: Species matrix (species recorded by plot)..... | 83 |
| Table 41: Vegetation integrity data (Composition, Structure and function) | 85 |

Abbreviations

| Abbreviation | Description |
|--------------|--|
| BAM | Biodiversity Assessment Method |
| BAMC | Biodiversity Assessment Method Credit Calculator |
| BC Act | NSW <i>Biodiversity Conservation Act 2016</i> |
| BDAR | Biodiversity Development Assessment Report |
| BGHF | Blue Gum High Forest |
| CEEC | Critically Endangered Ecological Community |
| DA | Development Application |
| DAWE | Commonwealth Department of Agriculture, Water and Environment (formally Department of Environment and Energy (DoEE)) |
| DotEE | Commonwealth Department of the Environment and Energy |
| DPE | NSW Department of Planning and Environment |
| EEC | Endangered Ecological Community |
| ELA | Eco Logical Australia Pty Ltd |
| EP&A Act | NSW Environmental Planning and Assessment Act 1979 |
| EPBC Act | Commonwealth <i>Environment Protection and Biodiversity Conservation Act 1999</i> |
| FM Act | NSW <i>Fisheries Management Act 1994</i> |
| GIS | Geographic Information System |
| GPS | Global Positioning System |
| HBT | Hollow-bearing Tree |
| HDCP | Hornsby Development Control Plan 2013 |
| HLEP | Hornsby Local Environmental Plan 2013 |
| IBRA | Interim Biogeographic Regionalisation for Australia |
| LEP | Local Environment Plan |
| LGA | Local Government Area |
| LLS | Local Land Service |
| MNES | Matters of Environmental Significance |
| NSW | New South Wales |

| Abbreviation | Description |
|--------------|--|
| NOW | NSW Office of Water |
| OEH | NSW Office of Environment and Heritage |
| PCT | Plant Community Type |
| SEARs | Secretary's Environmental Assessment Requirement |
| SEPP | State Environmental Planning Policy |
| SSD | State Significant Development |
| STIF | Sydney Turpentine Ironbark Forest |
| TEC | Threatened Ecological Community |
| VIS | Vegetation Information System |
| WM Act | NSW <i>Water Management Act 2000</i> |

1. State Significant Development Application (Updated Masterplan and Stage 1 Works)

The following text has been supplied by Ethos Urban to support the BDAR.

1.1 Introduction

This document is to be read in conjunction with the planning report prepared by Ethos Urban and amended consultant documents.

This revised Design Statement, prepared By Allen Jack + Cottier Architects (2020), responds to questions and comments received from the NSW Department of Planning and Environment (DPE) regarding the State Significant Development Application (SSDA) for Loreto Normanhurst Concept Master Plan and detailed design works for a new on-campus student boarding facility, through site road link, new carparking and drop-off/pick-up facilities, landscaping works, and demolition of buildings between the Mary Ward and Givendale dining building.

1.2 Background

1.2.1 Need for a Campus Masterplan

Loreto Normanhurst is an independent, Catholic day and boarding school for girls from Years 5 to 12. The existing school campus was established in 1897 and has evolved in an organic and ad-hoc manner across the span of a 120 years.

A new campus wide planning approach offers the opportunity to strategically review and plan for the campus' future in a sustainable and efficient manner such that the campus' unique aesthetic and ecological values are best preserved. The preparation of a campus wide masterplan is also consistent with the School's 'Loreto Normanhurst 2016 - 2020 Strategic Plan' which identified the need for a broader strategic plan to coordinate renewal and orderly development in a feasible and staged manner.

1.2.2 Early Learning Centre

A separate DA (D/1227/2018) has been submitted to Hornsby Shire Council on 23 November 2018 for an 80 place Early Learning Centre (ELC) building and the DA has been approved (Figure 1). The ELC building is consistent with the overall concept masterplan and was prepared concurrently with the final preferred campus masterplan. However, to meet the School's operational timeframe requirements for the ELC, a separate application was seen to be the best pathway to allow the building to be built, fitout and operational prior to the Masterplan.

1.3 The Site

Loreto Normanhurst is located within the suburb of Normanhurst on Sydney's Upper North Shore approximately 3km south of Hornsby and 25km north of Sydney CBD. The school is located in the local government area of Hornsby Shire Council, approximately 750m south of the Normanhurst Railway Station.

The site comprises the existing campus grounds of the Loreto Normanhurst school at 91 – 93 Pennant Hills Road, Normanhurst. The northern part of the site accommodates much of the school's existing built form, while the rear extent consists of the school's sporting fields, and a portion of largely undeveloped land covered in remnant vegetation.

The campus itself is bound by Pennant Hills Road (to the north), Osborn Road (to the west) and Mount Pleasant Avenue (to the east). Detached dwellings on individual residential lots about the southern boundary of the site.

1.3.1 Legal description and ownership

The campus comprises several allotments, the legal descriptions of which are provided in **Table 1** below. The existing campus has a site area of approximately 13.14 ha. The site in its entirety is owned by the Trustees of the Loreto Property Association.

Table 1: Legal Description

| Address | Lot | Plan |
|-------------------------------|--------------------------|------------|
| 4 Mount Pleasant Avenue | D | DP 366271 |
| 16 Mount Pleasant Avenue | Lot 5 | DP 1218765 |
| | Lot 16 | DP 6612 |
| 30 – 62 Mount Pleasant Avenue | Lots 20 – 23 and 25 – 36 | DP 6612 |
| | Lot 1 | DP 34834 |
| 91 – 93 Pennant Hills Road | Lot 1 | DP 114580 |
| | Lot 3 | DP 1217496 |
| | Lot 1 – Lot 3 | DP 1218765 |
| | Lot B | DP327538 |
| 24 – 28 Mount Pleasant Avenue | Lot 1 | DP 809066 |
| 6 Mount Pleasant Avenue | Lot C | DP 366271 |
| 14 Mount Pleasant Avenue | Lot 4 | DP1218765 |
| 89 Pennant Hills Road | Lot 1 | DP136156 |

1.4 Overview of proposed development

This application sets out a new campus masterplan for the existing school campus that will guide and shape the development of the school campus for the next 30 years. This SSDA resubmission includes detailed plans for the first stage of the concept proposal (Stage 1 works) (Figure 1). Accordingly, consent is sought for the following:

- The Stage 1 amendments including:
 - Boarding house
 - Garden Plaza
 - Osborn Road to Mt Pleasant Avenue link road
 - Interim Osborn Road surface carpark
 - Sports courts and partially underground carpark
 - Tennis courts and second partially underground carpark.

- Masterplan amendments, being:
 - Secondary school upgrade
 - Junior school upgrade
 - Mary Ward building upgrade
 - Gymnasium
 - Gonzaga Barry centre extension
 - All weather playing fields
 - Pedestrian bridge link to Mary Ward wing.

2. Stage 1: Biodiversity assessment

2.1 Introduction

This Biodiversity Development Assessment Report (BDAR) has been prepared by Belinda Failes, an accredited person (BAAS18159) under the NSW *Biodiversity Conservation Act 2016* (BC Act). And peer reviewed by Nicole McVicar (BAAS18077) who is also an accredited person under the BC Act.

A BDAR as part of the SSD application was submitted in 2019 for the proposed Concept Plan (30 year Masterplan) and Stage 1 construction works. Comments from the NSW Department of Planning and Environment (DPE) have been incorporated into the updated Masterplan and Stage 1 works as part of this SSD resubmission.

It is acknowledged that there may be minor amendments to the 30 year Masterplan Concept Plan and will be subject to future Development Applications (DAs). However, the biodiversity impacts of the Concept Plan including Stage 1 works have been considered in this updated BDAR. As such this BDAR will address the biodiversity impacts of the Concept Plan as the proposed footprint.

2.1.1 General description of the subject site

The Masterplan and Stage 1 works are located within Loreto Normanhurst subject site which is approximately 13.14 ha in size, bounded by Pennant Hills Road to the north, Mount Pleasant Avenue to the east, Osborn Road to the west and residential development to the south (Figure 2). The development footprint is 4.34 ha and is zoned R2 Low Density Residential which provides opportunities for residential or educational use with consent approval as defined in the *Hornsby Local Environmental Plan 2013* (HLEP).

The existing development is concentrated within the northern portion of the subject site. A large patch of native vegetation is located in the southern portion, this is separated from the existing educational buildings by a large sporting oval. Planted native trees and one remnant *Eucalyptus pilularis* (Blackbutt) are located in horticultural gardens which are intersperse around the existing buildings and along the eastern and western perimeters which link the scattered vegetation in the north to the vegetation in the south.

There is an intermittent drainage line which flows through the vegetated bushland in the south of subject site and drains into the headwaters to an unnamed Strahler first order stream. This unnamed stream flows south-west and merges with Coups Creek. Coups Creek is surrounded by a large riparian buffer outside of the subject site which is managed by the Sydney Adventist Aged Care Facility. In the broader landscape the subject site is located within an urbanised matrix with scattered canopy trees and to the south there is a large corridor of vegetation along Coups Creek and links to Lane Cove National Park.

This report includes the following base maps, the Concept Plan and Stage 1 works (Figure 1), Site Map (Figure 2) and the Location Map (Figure 3).

2.1.2 Subject site footprint

The development footprint includes the Concept Plan and Stage 1 construction works and are provided in Figure 1. The proposed development will primarily utilise the existing building footprints and open

space where available. The proposed development will involve modification of existing development and construction of several new buildings within the subject site. The Concept Plan includes staged development of the following facilities;

- boarding house
- one surface carpark
- two partially underground carparks and sporting facilities
- a link road between Osborn Road to Mount Pleasant Avenue
- and landscaping.

Scattered canopy trees and some landscaping gardens may be impacted or removed to accommodate the new development. The new Early Learning Centre is shown in this BDAR; however, it has been assessed as a separate Development Application (Figure 1). A detailed description of the project has been provided in the previous section.

2.1.3 Sources of information used

The following data sources were reviewed as part of this report:

- Biodiversity Assessment Methodology Calculator
- BioNet Vegetation Classification System
- BioNet / Atlas of NSW Wildlife 5 km database search (OEH 2018)
- *Environment Protection and Biodiversity Conservation Act 1999* EPBC Act Protected Matters Search Tool 5 km database search (DotEE 2018)
- The Native Vegetation of the Sydney Metropolitan Area (OEH 2016)
- Hornsby Shire Council vegetation mapping (HSC 2017)
- Aerial mapping (SIXMaps)
- Additional GIS datasets including soil, topography, geology and drainage.



Figure 1: Masterplan and Stage 1 works



Figure 2: Site Map




Figure 3: Location Map

2.2 Legislative context

Table 2: Legislative context

| Name | Relevance to the project | Report Section |
|--|---|----------------|
| Commonwealth | | |
| Environmental Protection and Biodiversity Conservation Act 1999 | Matters of National Environmental Significance (MNES) have been identified on or near the subject site. This report assesses impacts to MNES and concludes that the development is not likely to have a significant impact on MNES under the EPBC Act. | 3.6.1 |
| State | | |
| Biodiversity Conservation Act 2016 | The proposed development requires submission of a Biodiversity Development Assessment Report (BDAR) (i.e. this report) under the BC Act. | All |
| Environmental Planning and Assessment Act 1979 (EP&A Act) | The proposed development requires consent under the EP&A Act. | N/A |
| Fisheries Management Act 1994 | The development does not involve impacts to Key Fish Habitat, does not involve harm to marine vegetation, dredging, reclamation or obstruction of fish passage. A permit or consultation under the FM Act is not required. | N/A |
| Local Land Services Amendment Act 2016 (LLS) | Assessment under the LLS Act is not required this development. | N/A |
| Water Management Act 2000 | The project does not involve works on waterfront land or riparian land. As the project is under an SSD a Controlled Activity Approval under s91 of the WM Act is not required. | N/A |
| Planning Instruments | | |
| SEPP Coastal Management 2018 | SEPP Coastal Management 2018 consolidated SEPP 14 Coastal Wetlands, SEPP 26 Littoral Rainforests and SEPP 71 Coastal Protection. The proposed development is not located on land subject to SEPP Coastal Management 2018. | N/A |
| SEPP 2020 – Koala Habitat Protection | The proposed development does not impact on potential or core koala habitat as defined by SEPP. | N/A |
| SEPP (Vegetation in Non-Rural Areas) | This SEPP applies to development that does not require development consent. As this project requires development consent under the EP&A Act, application of the Vegetation SEPP is not required. | N/A |
| Hornsby Local Environmental Plan 2013 (HLEP) | The subject site is zoned R2 Low Density Residential under the HLEP. R2 zones require development consent for the construction of educational facilities. Section 6.4 Terrestrial Biodiversity of the HLEP applies to the bushland vegetation in the southern portion of the subject site. The proposed development impacts to vegetation mapped on the Terrestrial Biodiversity layer. This layer corresponds to the layer on the Biodiversity Values Map (see Figure 2). | 3.6.2 |
| Hornsby Development Control Plan 2013 (HDCP) | The HDCP has been reviewed for additional biodiversity provisions that may relate to the subject and subject site. No additional provisions were identified. | N/A |

2.3 Landscape features

2.3.1 IBRA regions and subregions

The subject site falls within the Sydney Basin Interim Biogeographic Regionalisation for Australia (IBRA) region and Cumberland subregion.

2.3.2 Mitchell Landscapes

The subject site falls within the Pennant Hills Ridges Mitchell Landscapes (DECC 2002) as outlined in Table 2.

Table 3: Mitchell Landscapes

| Mitchell landscape | Description | Area within subject site(ha) |
|----------------------|---|------------------------------|
| Pennant Hills Ridges | Rolling to moderately steep hills on Triassic shales and siltstones. Elevation from 10 to 90m with local relief 60m. Deep red texture-contrast soils on narrow hillcrests, red and brown to yellow texture-contrast soils on slopes becoming slightly harsher in drainage lines. Vegetation typically tall open forest of <i>Eucalyptus saligna</i> and <i>Syncarpia glomulifera</i> . Rainforest elements in protected moist gully heads are also present. | 13.14 |

2.3.3 Rivers and streams

The subject site contains the headwaters to an unnamed 1st order Strahler stream located in the south-western corner of the subject site (Figure 2) as outlined in Table 4. There is evidence of an intermittent drainage line within the bushland to the south of the subject site, which links with the headwaters to the unnamed stream. This is likely to support surface water runoff following heavy rainfall events. At other time of field surveys, this area was dry. There is potential that mapping of the hydroline could be updated to extend the start of the headwaters further north, inside the subject site.

Table 4: Rivers and streams

| River/stream | Order | Riparian buffer |
|--------------|--------------------------------|-----------------|
| Unnamed | 1 st Order Strahler | 10 m |

2.3.4 Wetlands

The subject site does not contain any wetlands.

2.3.5 Connectivity features

The subject site contains the connectivity features outlined in Table 5 and shown in Figure 3. Contiguous connections are present within the subject site, adjoining to vegetation within the subject site and into adjoining land. There are two main links, both located in the south of the subject site. The first occurs in the south-eastern corner where the vegetation within the subject site links with the intact riparian corridor along Coups Creek via a short gap in the vegetation at Mount Pleasant Avenue. Coups Creek riparian corridor is managed by a trust for the Sydney Adventist Aged Care Facility.

The second contiguous connection is present in the south-western corner where a tributary of Coups Creek links with vegetation on council land from Pine Street where it crosses The Comenarra Parkway

and into Lane Cove National Park, approximately 950 m to the south-west of the subject site. There is also a link in the north of the subject site along a tributary of Berowra Creek which flows into Berowra Creek Regional Park.

Connectivity to large tracts of habitat is considered suitable for highly mobile species such as birds and bats. This includes flyways for migratory birds and bat species moving through the landscape. Connectivity is present (with road crossings) for less mobile species such as reptiles and mammals.

Table 5: Connectivity features

| Connectivity feature name | Feature type |
|-------------------------------|------------------------------------|
| Coups Creek riparian corridor | Private land and riparian corridor |
| Lane Cove National Park | National Park |

2.3.6 Areas of geological significance and soil hazard features

The subject site does not contain areas of geological significance and soil hazard features.

2.3.7 Site context

2.3.7.1 Method applied

The site based method has been applied to this development.

2.3.7.2 Percent native vegetation cover in the landscape

The current percent native vegetation cover in the landscape was assessed in a Geographic Information System (GIS) using aerial imagery sourced from SIX Maps using increments of 5%. The percent native vegetation cover within the 1,500 m buffer area is 41 % (392.8 ha).

2.3.7.3 Patch size

Patch size was calculated using available vegetation mapping for all patches of intact native vegetation on and adjoining the subject site. The patch size area is 101 ha.

2.4 Native vegetation

2.4.1 Survey effort

An initial constraints assessment was conducted on 24 and 31 March 2017 to identify the vegetation type and condition of the vegetation community in the southern portion of the subject site for the proposed realignment of the sporting field. Three vegetation plots were undertaken in the bushland in the south using floristic plots (20 x 20m) plots to confirm the vegetation type and condition. These plots were not used in any credit calculations.

The site visits also involved vegetation mapping of the remaining subject site and mapping of habitat features, namely hollow-bearing trees (HBTs).

Additional vegetation survey was undertaken within the subject site by Belinda Failes on 12 October and 5th November 2018 (Figure 5) in accordance with the BC Act and BAM 2017. A total of four full-floristic and vegetation integrity plots (BAM plots) were undertaken in accordance with the BAM (Table 6).

All field data collected in the BAM plots is included in Appendix B:.

Table 6: Full floristic and vegetation integrity plots

| Veg Zone | PCT ID | PCT Name | Ancillary code | Condition | Total area | Impact area (ha) | Plots required | Plots surveyed |
|--------------|--------|-----------------------------------|----------------|-----------|--------------|------------------|----------------|----------------|
| 1 | 1237 | Blue Gum High Forest | Remnant | Moderate | 0.06 | 0.001 | 1 | 0* |
| 2 | 1237 | Blue Gum High Forest | Planted native | Low | 0.99 | 0.35 | 1 | 3** |
| 3 | 1281 | Sydney Turpentine-Ironbark Forest | Remnant | Moderate | 2.66 | 0 | 0 | 0 |
| 4 | 1281 | Sydney Turpentine-Ironbark Forest | Weedy | Low | 1.46 | 0.05 | 1 | 1 |
| 0 | - | Urban Exotics/ Native | - | - | 1.01 | 0.31 | - | - |
| 0 | - | Cleared | - | - | 6.96 | 3.62 | - | - |
| TOTAL | | | | | 13.14 | 4.34 | | |

* Due to the insignificant size of the vegetation zone 1 within the subject site, a plot could not be undertaken.

**Three plots were undertaken for Vegetation zone 2, however, only two plots were used in the credit calculator as one plot was located outside of the subject site area.

2.4.2 Plant Community Types present

Two PCTs were identified in the subject site (Table 7, Figure 4). Both of these PCTs may be listed as a TECs under the BC and/or EPBC Act (Table 8, Figure 6) (refer to Section 2.4.2.2 for further details). The subject site also contains planted native canopy, shrubs and occasionally ground cover species which are native to NSW, however these were not considered locally indigenous to the PCTs. However, under the BAM 2017, planted vegetation native to NSW requires consideration as to the 'best fit' PCT. Based on the soil landscape, elevation, presence of remnant vegetation and remnant regrowth vegetation within the subject site and along the eastern perimeter (i.e. Osbourn Road), it was determined that planted native vegetation 'best-fit' PCT was PCT 1237. Justifications are provided below in Section 2.4.2.1.

Table 7: Plant Community Types

| PCT ID | PCT Name | Vegetation Class | Vegetation Formation | Percent cleared |
|--------|-----------------------------------|---|---|-----------------|
| 1237 | Blue Gum High Forest | North Coast Wet Sclerophyll Forests | Wet Sclerophyll Forests (Shrubby sub-formation) | 90% |
| 1281 | Sydney Turpentine-Ironbark Forest | Northern Hinterland Wet Sclerophyll Forests | Wet Sclerophyll Forests (Grassy sub-formation) | 90% |

Table 8: Threatened Ecological Communities

| PCT ID | BC Act | | | EPBC Act | | |
|--------|----------------|---|-----------|----------------|----------------------------|-----------|
| | Listing status | Name | Area (ha) | Listing status | Name | Area (ha) |
| 1237 | CEEC | Blue Gum High Forest in the Sydney Basin Bioregion | 0.001 | * | Blue Gum High Forest | N/A |
| 1281 | EEC | Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion | 0.05 | CEEC | Turpentine-Ironbark Forest | 0.05 |

CEEC – Critically endangered ecological community; EEC – Endangered ecological community

* Note that planted PCT 1237 (0.35 ha) did not satisfy the requirements for listing under the BC Act or EPBC Act criteria.

Table 9: PCT selection justification

| PCT ID | PCT scientific name | Selection criteria | Justification |
|--------|--|--|--|
| 1237 | Sydney Blue Gum - Blackbutt - Smooth-barked Apple moist shrubby open forest on shale ridges of the Hornsby Plateau, Sydney Basin Bioregion | IBRA region, subregion, soil landscape, elevation and presence of canopy species <i>Eucalyptus saligna</i> and <i>E. pilularis</i> | Remnant patches of vegetation consistent with this PCT have been mapped within the subject site and along the road verge of Osbourn Road. This PCT has been accepted as the best fit PCT for planted native vegetation scattered around the subject site due to the location in the landscape, presence of remnant species within the area and soil profile. |
| 1281 | Sydney Turpentine-Ironbark Forest | IBRA region, subregion, soil landscape, elevation and results of floristic plot analysis including the presence of canopy species <i>Syncarpia glomulifera</i> | This PCT has been accepted as the best fit PCT for remnant vegetation located in the south of the site based on floristic analysis. |

2.4.2.1 PCT selection justification

The boundaries of between PCT 1237 and PCT 1281 were delineated by the transition of two different soil landscapes, change in elevation and the presence of key characteristic canopy species namely presence or absence of *Eucalyptus saligna* (Sydney Blue Gum), *E. pilularis* (Blackbutt) and *Syncarpia glomulifera* (Turpentine). At higher elevations on Glenorie soil landscapes the vegetation was mapped as PCT 1237. Glenorie soil landscape is associated with clay soils of the Wianamatta Group Ashfield Shale (Chapman and Murphy 1989). This vegetation also includes patches of remnant vegetation along Osbourn Road and near Pennant Hills Road. Scattered patches of planted native vegetation within the higher elevations on the same soil landscape were also mapped as part of this PCT 1237, however, they were not considered part of the TEC (see justification in Section 2.4.2.2 below).

PCT 1281 was generally located in the intact native vegetation in the southern portion of the subject site and only a small portion was represented in the subject site. PCT 1281 was located in lower elevations on Hawkesbury soil landscape which is associated with sandier soils. Floristic analysis was conducted to confirm the presence of PCT 1281 and delineate with PCT 1237 (Appendix E). PCT 1237 occurs at the transition between Wianamatta Shale and Hawkesbury Sandstone and corresponds to TEC Sydney Turpentine Ironbark Forest. Justification for the selection of PCTs occurring on the subject site is based on a quantitative analysis of full-floristic plot data (Appendix E) and is provided in Table 9.

2.4.2.2 Threatened Ecological Communities Justification

The BioNet Vegetation Classification lists PCT 1237 as Blue Gum High Forest which is listed as critically endangered ecological community (CEEC) under the BC Act. Patches of certain quality Blue Gum High Forest may be listed under the EPBC Act listed provided they satisfy the following criteria (DotEE 2018a):

- Patch size is greater than 1 ha; AND
 - Canopy cover greater than 10% OR
 - Canopy cover less than 10% and occurs in areas of vegetation in excess of 5 ha.

There was one remnant patch of PCT 1237 located within the subject site. This was represented by a large remnant *Eucalyptus pilularis* located in a landscaped garden in the north of the subject site and is isolated from other remnant vegetation. This patch does not satisfy the criteria for listing under the EPBC Act as the patch size is less than 1 ha.

The remaining patches of PCT 1237 mapped within the subject site represent planted native species including species which are indigenous to PCT 1237 such as *Eucalyptus saligna* and native planted species which are not indigenous to PCT 1237 such as *Ficus macrophylla* (Moreton Bay Fig). A review of the final determination for this Blue Gum High Forest TEC and profile description determined that the native planted vegetation within the subject site does not conform to the TEC. This is due to a lack of regeneration of native vegetation and a low probability of an existing soil seed bank due to the fact that the landscape and soil profile has been substantially altered.

Another small patch of PCT 1237 represented by remnant *Eucalyptus saligna* was mapped along the road verge of Osbourn Road and is located outside of the subject site. Some of the branches were overhanging into the subject site and have therefore been included in the PCT mapping (see Figure 5). There is potential that the road verge patch of PCT 1237 would be considered for listing under the EPBC Act due to its connectivity with the large patch of PCT 1281 in the south of the subject site. However, this patch of road verge PCT 1237 is located outside of the subject site and will not be directly impacted under the proposed development.

The BioNet Vegetation Classification lists PCT 1281 as a component of Sydney Turpentine Ironbark Forest which is listed as endangered under the BC Act and CEEC under the EPBC Act. This PCT was categorised as high and moderate condition vegetation zones based on the presence/absence of weeds. Both conditions of this PCT 1281 are listed as part of the Sydney Turpentine Ironbark Forest listing under the BC Act and the more stringent criteria for listing under the EPBC Act. The criteria for listing under the EPBC Act for Sydney Turpentine Ironbark Forest are provided below (DotEE 2018b):

- The vegetation contains some characteristic components from all structural layers (tree canopy, small tree/shrub midstorey, and understorey).
- Tree canopy cover is greater than 10% and remnant size is greater than one hectare. These areas have the greatest conservation value and their high quality and size makes them most resilient to disturbance.
- However, remnants with tree canopy cover less than 10% are also included in the ecological community, if the fragments are greater than one hectare in size and occur in areas of native vegetation in excess of 5 hectares in area. These areas enhance the potential for

connectivity and viability of the ecological community. They support native flora and fauna species by facilitating gene flow among remnants and buffering against disturbance.

2.4.3 Vegetation integrity assessment

A vegetation integrity assessment using the Credit Calculator (referred to as the BAMC) was undertaken and the results are outlined in Table 10.

The impact areas were entered into the BAMC for each of the vegetation zones. The small impact area for vegetation zone 1 (0.001 ha) was entered into the BAMC, however, due to the small impact size for this zone, the BAMC reverted the scores to zero value and would not allow further assessment for this vegetation zone. Therefore, the assessor added the impact area values (prior to rounding) for zone 1 (0.001 ha) and zone 2 (0.35 ha) together as they are of the same PCT. However, due to the insignificant value it did not result in a change of score for vegetation zone 2 (0.35 ha). Therefore, the vegetation integrity assessment has included the impacts for zone 1, however, the impact values do not change the overall integrity score of the development. No impacts will occur to vegetation zone 3 so no plot data was obtained for this vegetation zone.

Table 10: Vegetation integrity

| Veg Zone | PCT ID | Ancillary code | Condition | Area Impacted (ha) | Composition Condition Score | Structure Condition Score | Function Condition Score | Current vegetation integrity score |
|----------|--------|-----------------------|-----------|--------------------|-----------------------------|---------------------------|--------------------------|------------------------------------|
| 1 | 1237 | Remnant regeneration | Moderate | 0.001* | N/A | N/A | N/A | N/A |
| 2 | 1237 | Planted native canopy | Low | 0.350 | 24.3 | 48.9 | 36.5 | 35.1 |
| 3 | 1281 | Remnant | Moderate | 0 | N/A | N/A | N/A | N/A |
| 4 | 1281 | Weedy | Low | 0.05 | 40.8 | 51.2 | 30 | 39.7 |

* Note – Vegetation Zones 1 was entered into the credit calculator, however, the impact areas are too low that the credit calculator rounded the values to zero. Therefore, the impact hectares for Zone 1 was added to Zone 2, however, the due to the small amounts, this did not change the value of Zone 2.



Figure 4: Plant Community Types and native vegetation extent



Figure 5: Plot locations



Figure 6: Threatened Ecological Communities

2.5 Threatened species

2.5.1 Threatened flora species

Three flora species which have been planted as horticultural varieties which in some locations are listed as threatened flora species under the BC and/or EPBC Act were recorded within the subject site. These species are located outside of their normal distribution and are cultivated varieties and should not be considered as threatened species. These include:

- *Grevillea juniperina* is a horticultural variety and should not be confused with the threatened *Grevillea juniperina* subsp. *juniperina* which is associated with clay soils of western Sydney region and listed as a vulnerable species under the BC Act. The subject site does not comprise appropriate habitat for this threatened species and would therefore not occur naturally.
- *Eucalyptus scoparia* (Wallangarra White Gum) listed as endangered under the BC Act and vulnerable under the EPBC Act. This species has been planted. The threatened species is known from only three locations in NSW near Tenterfield, which is more than 640 km from the subject site.
- *Syzygium paniculatum* (Magenta Cherry), this species is readily available from local nurseries as a horticultural species. This species is also listed as endangered under the BC Act and vulnerable under the EPBC Act. The threatened species is only located in littoral coastal rainforest areas along NSW from Upper Lansdowne to Conjola State Forest. The subject site does not include littoral coastal rainforest environments.

According to the arborist report (Earthscape Horticultural Services 2020) the following threatened species have been assessed and will be retained within the subject site:

- *Eucalyptus scoparia* – identified as Tree 9, 10, 210, 390 and 424
- *Syzygium paniculatum* – identified as Tree 151, 152, 153, 155, 190a and 190b.

The location of these trees are provided in the arborist report (Earthscape Horticultural Services 2020). In accordance with the new BAM 2020 Appendix D (Table 11) the following assessment was conducted for planted threatened species and found that these species have not been established as part of a species recovery, *Saving our Species* project, condition of consent or rehabilitation then these species do not require assessment in accordance with Chapters 4 or 5 of the BAM.

Additionally, Appendix D2 Assessment of planted native vegetation for threatened species habitat was conducted and determined that the planted threatened species does not provide habitat for threatened species credit species. Therefore, consideration of these species beyond Chapters 4 and 5 of the BAM 2020 are not required.

Although that these species are located outside of their natural range of distribution and/or outside of their natural habitat and the fact these species have been clearly planted due to the landscaped setting an assessment of these species in accordance with the EPBC Act has been provided in this BDAR.

Table 11: Decision making tool for Planted Native Vegetation in accordance with Appendix D of the BAM 2020

| Decision making key | Response |
|--|---|
| <p>1) Does the planted native vegetation occur within an area that contains a mosaic of planted and remnant native vegetation and which can be reasonably assigned to a PCT known to occur in the same IBRA subregion as the proposal?</p> <p>i Yes – the planted native vegetation must be allocated to the best-fit PCT and the BAM must be applied.</p> <p>ii No – Go to 2.</p> | No, the planted vegetation does not occur in a mosaic of planted and remnant vegetation. |
| <p>2. Is the planted native vegetation:</p> <p>a. Planted for the purpose of environmental rehabilitation or restoration under an existing conservation obligation listed in BAM Section 11.9(2.), and</p> <p>b. The primary objective was to replace or regenerate a plant community type of a threatened plant species or its habitat?</p> <p>i Yes – the planted native vegetation must be assessed in accordance with Chapters 4 and 5 of the BAM</p> <p>ii No – Go to 3.</p> | <p>No, the vegetation has not been established for rehabilitation or restoration works.</p> <p>No, the primary objective was not conducted to replace or regenerate a PCT as the vegetation consists of non-indigenous species to the area.</p> |
| <p>3. Is the planted / translocated native vegetation individuals of a threatened species or other native species planted/ translocated for the purpose of providing threatened species habitat under one of the following:</p> <p>a. A species recovery project</p> <p>b. <i>Saving our Species</i> project</p> <p>c. Other types of government funded restoration project</p> <p>d. Condition of consent for a development approval that required those species to be planted or translocated for the purpose of providing threatened species habitat</p> <p>e. Legal obligation as part of a condition of ruling of court. This includes regulatory directed or ordered remedial plantings (e.g. Remediation Order for clearing without consent issued under the BC Act or the Native Vegetation Act)</p> <p>f. Ecological rehabilitation to re-establish a PCT or TEC that was, or is carried out under a mine operations plan, or</p> <p>g. Approved vegetation management plan (e.g. as required as part of a Controlled Activity Approval for works on waterfront land under the NSW <i>Water Management Act 2000</i>)?</p> <p>i Yes – the planted native vegetation must be assessed in accordance with Chapters 4 and 5 of the BAM</p> <p>• No – Go to 3.</p> | No, the planted vegetation does not include translocated native vegetation of threatened species. |
| <p>4. Was the planted native vegetation (including individuals of a threatened flora species) undertaken voluntarily for revegetation, environmental rehabilitation or restoration within a legal obligation to secure or provide for management of the native vegetation?</p> <p>i Yes – Go to D.2 Assessment of planted native vegetation for threatened species habitat (the use of Chapters 4 and 5 of the BAM are not required to be applied)</p> <p>• No – Go to 5.</p> | No, the planted threatened species was not conducted for revegetation works or environmental rehabilitation or restoration. The threatened species are located in a highly urbanised environment in landscaped gardens around the school. |
| <p>5. Is the planted native vegetation (including individuals of a threatened flora species) planted for functional, aesthetic, horticultural or plantation forestry</p> | Yes, the planted threatened flora species were planted for |

| Decision making key | Response |
|---|--|
| <p>purposes? This includes examples such as; windbreaks in agricultural landscapes, roadside plantings (including street trees, median stripes, roadside batters), landscaping in parks, gardens and sport fields/complexes, macadamia plantations or teatree farms?</p> <ul style="list-style-type: none"> i Yes – Go to D.2 Assessment of planted native vegetation for threatened species habitat (the use of Chapters 4 and 5 of the BAM are not required to be applied) ii No – Go to 6. | <p>aesthetics in horticultural gardens.</p> <p>An assessment of Appendix D2 has determined that the threatened species do not provide habitat for threatened species credit species. Therefore, no additional consideration is required under the BAM.</p> |
| <p>6. Is the planted native vegetation a species listed as a widely cultivated native species on a list approved by the Secretary of the Department (or an officer authorised by the Secretary)?</p> <ul style="list-style-type: none"> i Yes – Go to D.2 Assessment of planted native vegetation for threatened species habitat (the use of Chapters 4 and 5 of the BAM are not required to be applied) ii No – There may be other types of occurrences of planted native vegetation that do not easily fit into the decision-making key above. | - |

2.5.2 Ecosystem credit species

Ecosystem credit species predicted to occur at the subject site, their associated habitat constraints, geographic limitations and sensitivity to gain class is included in Table 12.

Table 12: Predicted ecosystem credit species

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---------------------------------|----------------------------------|--|---------------------------|--------------------|---------------------|--|
| <i>Anthochaera phrygia</i> | Regent Honeyeater (Foraging) | N/A | High | CE | CE | Included The subject site contains flowering trees suitable for this species. |
| <i>Artamus cyanopterus</i> | Dusky Woodswallow | | Moderate | V | Not Listed | Included The subject site provides suitable foraging habitat for this specie. |
| <i>Callocephalon fimbriatum</i> | Gang-gang Cockatoo (Foraging) | N/A | Moderate | V | Not Listed | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. Additionally, the Gang-gang Cockatoo favours old growth forest/woodland attributes, of which the subject site does not contain. |
| <i>Calyptorhynchus lathami</i> | Glossy Black-Cockatoo (Foraging) | N/A | High | V | Not Listed | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. The subject site was not considered suitable habitat due to disturbance and insufficient presence of foraging habitat. |
| <i>Chthonicola sagittata</i> | Speckled Warbler | N/A | High | V | Not Listed | Excluded Habitat present does not contain suitable habitat features for this species such as abundance of fallen logs. The vegetation within the subject site is substantially degraded. |

| Species | Common Name | | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|-----------------------------------|----------------------|-------|---|---------------------------|--------------------|---------------------|--|
| <i>Daphoenositta chrysoptera</i> | Varied Sittella | | N/A | Moderate | V | Not Listed | Included Habitat present does not contain suitable habitat features for this species such as abundance of fallen logs. The vegetation within the subject site is substantially degraded |
| <i>Dasyurus maculatus</i> | Spotted-tailed Quoll | | N/A | High | V | E | Excluded Habitat features for this species are not present at this site. This species requires habitat features such as maternal den sites, an abundance of food (birds and small mammals) and large areas of relatively intact vegetation to forage in (DECC 2007). |
| <i>Falsistrellus tasmaniensis</i> | Eastern Pipistrelle | False | N/A | High | V | Not Listed | Included Habitat present is substantially degraded however, the subject site still provides foraging habitat for this species. This species was included in this assessment. |
| <i>Glossopsitta pusilla</i> | Little Lorikeet | | N/A | High | V | Not Listed | Included There are eight BioNet records for this species within a 5 km radius of the subject site. This species may utilise the flowering species within the subject site This species was included in this assessment |
| <i>Grantiella picta</i> | Painted Honeyeater | | Other Mistletoes present at a density of greater than five mistletoes per hectare | Moderate | V | V | Excluded Habitat features associated with this species are not present in the subject site. This species is a specialist feeder requiring mistletoe which is absent from the subject site. |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|----------------------------------|--------------------------------------|--|---------------------------|--------------------|---------------------|---|
| <i>Hieraaetus morphnoides</i> | Little Eagle (Foraging) | N/A | Moderate | V | Not Listed | Included Included in this assessment. |
| <i>Hoplocephalus bungaroides</i> | Broad-headed Snake (Foraging) | The south west margins of the sub-region | High | E | V | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. The subject site does not contain sufficient rocky habitat for this species to utilise. No individuals have been recorded within 5km of the subject site. |
| <i>Lathamus discolor</i> | Swift Parrot (Foraging) | N/A | Moderate | E | CE | Included Habitat features associated with this species were identified within the subject site such as <i>Corymbia maculata</i> and <i>Eucalyptus sideroxylon</i> within PCT1237_native Planted and <i>Corymbia. gummifera</i> and <i>Eucalyptus pilularis</i> used for lerps in PCT1237 and PCT 1281. Therefore, this species has been included in this assessment. |
| <i>Lophoictinia isura</i> | Square-tailed Kite (Foraging) | N/A | Moderate | V | Not Listed | Included Included in this assessment. |
| <i>Melanodryas cucullata</i> | Hooded Robin (south-eastern form) | N/A | Moderate | V | Not Listed | Included Included in this assessment. |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---------------------------------------|---|--|---------------------------|--------------------|---------------------|---|
| <i>Melithreptus gularis gularis</i> | Black-chinned Honeyeater (eastern subspecies) | N/A | Moderate | V | Not Listed | Excluded Habitat features associated with this species are not present in the subject site. This species occupies forests or woodlands dominated by box and ironbark eucalypts (especially Mugga Ironbark), which the subject site is not dominated by. No individuals have been recorded within 5km of the subject site. |
| <i>Miniopterus australis</i> | Little Bent-winged Bat (Foraging) | N/A | High | V | Not Listed | Included Included in this assessment |
| <i>Miniopterus orianae oceanensis</i> | Large Bent-winged Bat (Foraging) | N/A | High | V | Not Listed | Included Included in this assessment |
| <i>Micronomus norfolkensis</i> | Eastern Freetail-bat | N/A | High | V | Not Listed | Included Included in this assessment |
| <i>Neophema pulchella</i> | Turquoise Parrot | N/A | High | V | Not Listed | Excluded Habitat features associated with this species are not present in the subject site. The subject site does not contain suitable hollow-bearing trees required for breeding (within woodland) or foraging habitat in open woodland required for this species. No individuals have been recorded within 5km of the subject site. |
| <i>Ninox connivens</i> | Barking Owl (Foraging) | N/A | High | V | Not Listed | Included Included in this assessment. Of PCT 1281_weedy. |
| <i>Ninox strenua</i> | Powerful Owl (Foraging) | N/A | High | V | Not Listed | Included Included in this assessment. |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|-------------------------------|-------------------------------|--|---------------------------|--------------------|---------------------|--|
| <i>Petaurus australis</i> | Yellow-bellied Glider | Hollow-bearing Trees (hollow > 25cm) | High | V | Not Listed | Excluded Habitat features associated with this species (i.e. old growth forests and large hollows) are not present in the subject site. The subject site does not contain suitable hollow-bearing trees required for breeding (within woodland) or foraging habitat in open woodland required for this species. No individuals have been recorded within 5km of the subject site. |
| <i>Petroica boodang</i> | Scarlet Robin | N/A | Moderate | V | Not Listed | Included There is a 2000 BioNet record for this species recorded within the subject site. However, the subject site does not contain breeding habitat (i.e. breeds on ridges, hills and foothills). After breeding, this species disburses to lower elevations. Habitat features associated with this species includes an abundance of logs and fallen timber, these features were not present in the subject site. |
| <i>Petroica phoenicea</i> | Flame Robin | N/A | Moderate | V | Not Listed | Included Included in this assessment |
| <i>Phascolarctos cinereus</i> | Koala (Foraging) | N/A | High | V | V | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. Habitat was not considered suitable due to the high disturbance and limited feed trees. |
| <i>Pteropus poliocephalus</i> | Grey-headed fox (Foraging) | N/A | High | V | V | Included Included in this assessment. |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---------------------------------|-------------------------------|---|---------------------------|--------------------|---------------------|---|
| <i>Ptilinopus superb</i> | Superb Fruit-Dove | N/A | Moderate | V | Not Listed | Included Included in this assessment |
| <i>Saccolaimus flaviventris</i> | Yellow-bellied Sheathtail-bat | N/A | High | V | Not Listed | Included Included in this assessment. |
| <i>Scoteanax rueppellii</i> | Greater Broad-nosed Bat | N/A | High | V | Not Listed | Included Included in this assessment. |
| <i>Tyto novaehollandiae</i> | Masked Owl (Foraging) | N/A | High | V | Not Listed | Included Included in this assessment. |
| <i>Varanus rosenbergi</i> | Rosenberg's Goanna | To northern and south western margins of the sub region | High | V | Not Listed | Excluded Habitat features for this species are not present in the subject site. Critical habitat components such as termite mounds are not present in the subject site. No individuals have been recorded within 5km of the subject site. |

CE = Critically Endangered; E = Endangered; E2 = Endangered Population; V = Vulnerable

2.6 Species credit species

Species credit species predicted to occur in the subject site (i.e. candidate species), their associated habitat constraints, geographic limitations and sensitivity to gain class are shown in Table 13. Habitat assessments were undertaken during the field surveys on 24 and 31 March 2017 and 12 October and 5th November 2018 to determine the likelihood of threatened species occurring within the subject site on an intermittent or permanent basis.

Habitat assessments involved a search of all possible hollow-bearing trees within the subject site, on ground inspection of roof cavities using binoculars for possible entrance for microbats, indirect evidence of fauna foraging such as chew cones or sap trees or roosting habitat in the form of white wash and pellets for owl species.

Table 13: Candidate species credit species

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---|---|--|---------------------------|--------------------|---------------------|---|
| <i>Acacia bynoeana</i> | Bynoe's Wattle | N/A | High | E | V | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Acacia prominens</i> – endangered population | Endangered population Gosford Wattle, Hurstville and Kogarah LGAs | N/A | High | E2 | Not Listed | Excluded The subject site is not located within the Gosford, Hurstville or Kogarah LGAs. This species is <u>not</u> considered a candidate species for this assessment. |
| <i>Acacia pubescens</i> | Downy Wattle | N/A | High | V | V | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Anthochaera phrygia</i> | Regent Honeyeater (Breeding) | As per mapped areas | High | CE | CE | Excluded The subject site is not located within the mapped important areas for this species (15 December 2020). |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---|---|--|---------------------------|--------------------|---------------------|---|
| <i>Burhinus grallarius</i> | Bush Stone-curlew | Fallen/standing dead timber including logs | High | E | Not Listed | Excluded Habitat features for this species are not present in the subject site. Critical habitat components such as fallen or standing dead timber are not present in the subject site. No individuals have been recorded within 5km of the subject site. |
| <i>Caladenia tessellata</i> | Thick Lip Spider Orchid | N/A | Moderate | E | V | Excluded Field surveys of the subject site was conducted during flowering season (September to November) and did not identify this species. This species occurs in grassy sclerophyll woodlands which were not recorded within the subject site. Furthermore, this species is only known from old records in Sydney area. |
| <i>Callocephalon fimbriatum</i> | Gang-gang Cockatoo (Breeding) | N/A | High | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain breeding habitat such as Eucalypt trees with hollows >9cm in diameter and shrubs that are suitable for the species to utilise the site. |
| <i>Callocephalon fimbriatum</i> - endangered population | Gang-gang Cockatoo population in the Hornsby and Ku-ring-gai Local Government Areas | Hornsby and Ku-ring-gai LGAs | High | E | Not Listed | Excluded Although the subject site is located within the Hornsby LGA and there are records for this species within 1.7 km away, the subject site does not contain suitable habitat for this species. |
| <i>Calyptorhynchus lathami</i> | Glossy Black-Cockatoo (Breeding) | N/A | High | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain larger patches of intact |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|--|----------------------|---|---------------------------|--------------------|---------------------|--|
| | | | | | | vegetation or trees with large hollows that are suitable for the species to utilise the site. |
| <i>Camarophyllopsis kearneyi</i> | - | Lane Cove Bushland Park | High | E | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Park (it is located 22 km away to the south-east of the Subject site). This species is unlikely to occur within the Subject site. |
| <i>Cercartetus nanus</i> | Eastern Pygmy-possum | | High | V | Not Listed | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. There is no nesting habitat present or preferred foraging habitat such as <i>Banksia</i> sp. present. No individuals have been recorded within 5km of the subject site. |
| <i>Chalinolobus dwyeri</i> | Large-eared Pied Bat | Cliffs Within 2km of rocky areas containing caves, overhangs, escarpment, outcrops, or crevices, or within 2km of old mines or tunnels | Very High | V | V | Excluded Habitat features associated with this species (caves) are not present in the subject site. There is no suitable breeding habitat such as caves, overhangs, mines or culverts present for the species to utilise the site. |
| <i>Epacris purpurascens</i> var. <i>purpurascens</i> | - | N/A | Moderate | V | Not Listed | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Galium australe</i> | Tangled Bedstraw | N/A | High | E | Not Listed | Excluded |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|--|-------------------------|--|---------------------------|--------------------|---------------------|---|
| | | | | | | The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Grammitis stenophylla</i> | Narrow-leaf Finger Fern | N/A | Moderate | E | Not Listed | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Grevillea parviflora</i> subsp. <i>parviflora</i> | Small-flower Grevillea | N/A | High | V | V | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Grevillea parviflora</i> subsp. <i>supplicans</i> | - | North of the Great Western Highway | High | E | Not Listed | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Gyrostemon thesioides</i> | Gyrostemon thesioides | N/A | High | E | Not Listed | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Heleioporus australiacus</i> | Giant Burrowing Frog | N/A | Moderate | V | V | Excluded Habitat features associated with this species are not present on the subject site. The subject site does not contain suitable waterbodies for this species to utilise the site. |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|--|-------------------------------|--|---------------------------|--------------------|---------------------|---|
| <i>Hibbertia puberula</i> | Hibbertia puberula | N/A | High | E | Not Listed | Excluded Habitat features associated with this species are not present on the subject site. |
| <i>Hibbertia spanantha</i> | Julian's Hibbertia | N/A | N/A | CE | CE | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Hibbertia superans</i> | Hibbertia superans | Other Ridgetops | High | E | Not Listed | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Hieraetus morphnoides</i> | Little Eagle (Breeding) | N/A | Moderate | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain suitable breeding habitat. |
| <i>Hoplocephalus bungaroides</i> | Broad-headed Snake (Breeding) | The south west margins of the region | Very High | E | V | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain suitable breeding habitat. |
| <i>Hygrocybe anomala</i> var. <i>ianthinomarginata</i> | - | Lane Cove Bushland Reserve | High | V | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the subject site). This species is unlikely to occur within the subject site. |
| <i>Hygrocybe aurantipes</i> | - | Lane Cove Bushland Reserve | High | V | Not Listed | Excluded |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|----------------------------------|-------------|--|---------------------------|--------------------|---------------------|---|
| | | | | | | The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the subject site). This species is unlikely to occur within the subject site. |
| <i>Hygrocybe austropratensis</i> | - | Lane Cove Bushland Reserve | High | E | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the subject site). This species is unlikely to occur within the Subject site |
| <i>Hygrocybe collucera</i> | | Lane Cove Bushland Reserve | High | E | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the Subject site). This species is unlikely to occur within the Subject site |
| <i>Hygrocybe griseoramosa</i> | | Lane Cove Bushland Reserve | High | E | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the Subject site). This species is unlikely to occur within the Subject site |
| <i>Hygrocybe lanecovens</i> | | Lane Cove Bushland Reserve | High | E | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the Subject site). This species is unlikely to occur within the Subject site |
| <i>Hygrocybe reesia</i> | | Lane Cove Bushland Reserve | High | V | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the Subject site). This species is unlikely to occur within the Subject site |
| <i>Hygrocybe rubronivea</i> | | Lane Cove Bushland Reserve | High | V | Not Listed | Excluded The subject site is not in within Lane Cove Bushland Reserve (it is located 22 km away to the south-east of the Subject site). This species is unlikely to occur within the Subject site |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|-------------------------------|--|--|---------------------------|--------------------|---------------------|--|
| <i>Lathamus discolor</i> | Swift Parrot (Important foraging areas) | | Moderate | E | CE | Excluded The subject site is not located within mapped important habitat for this species in the BOAMs (accessed 15 December 2020). |
| <i>Litoria aurea</i> | Green and Golden Bell Frog | Semi-permanent/ephemeral wet areas Within 1km of wet areas swamps Within 1km of swamp waterbodies Within 1km of waterbody | High | E | V | Excluded Habitat features associated with this species are not present on the subject site. Although the subject site is located within 1 km of waterbody/ streams, there are no suitable pools, swamps or fringing vegetation within the subject site which may contain suitable habitat for this species |
| <i>Lophoictinia isura</i> | Square-tailed Kite (Breeding) | N/A | Moderate | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain breeding habitat that is suitable for the species to utilise the site. No nests were observed during field surveys. |
| <i>Meridolum corneovirens</i> | Cumberland Plain Land Snail | N/A | High | E | Not Listed | Excluded Habitat features associated with this species are not present in the subject site. This species occurs within Cumberland Plain Woodland and associated shale vegetation communities. The subject site does not support these habitat features. |
| <i>Miniopterus australis</i> | Little Bent-winged Bat (Breeding) | Cave, tunnel, mine, culvert or other structure known or | Very High | V | Not Listed | Excluded |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---------------------------------------|----------------------------------|---|---------------------------|--------------------|---------------------|--|
| | | suspected to be used for breeding | | | | This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain breeding habitat such as caves that are suitable for the species to utilise the site. |
| <i>Miniopterus orianae oceanensis</i> | Large Bent-winged Bat (Breeding) | Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding | Very High | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain breeding habitat such as caves, tunnels, mines or culverts. |
| <i>Myotis macropus</i> | Southern Myotis | Hollow bearing trees Within 200 m of riparian zone other bridges, caves or artificial structures within 200 m of riparian zone | High | V | Not Listed | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. Habitat within the subject site is isolated and disturbed with a higher likelihood of this species using adjoining vegetation in better condition. Additionally, the nearest drainage line (approximately 100 m away from the subject site is only ephemeral in nature. Additionally, no HBTs will be removed in the bushland area. |
| <i>Ninox connivens</i> | Barking Owl (Breeding) | N/A | High | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain suitable breeding habitat. |
| <i>Ninox strenua</i> | Powerful Owl (Breeding) | N/A | High | V | Not Listed | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain suitable breeding habitat. |
| <i>Persoonia hirsuta</i> | Hairy Geebung | N/A | High | E | E | Excluded |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|--|--|--|---------------------------|--------------------|---------------------|---|
| | | | | | | The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Petaurus norfolcensis</i> | Squirrel Glider | N/G | High | V | Not Listed | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. Habitat in the subject site is isolated and disturbed with a higher likelihood of this species using adjoining vegetation in better condition. Additionally, this species has a strong preference for old growth forests which does not include the subject site. Additionally, there are no BioNet records for this species within a 5 km radius of the subject site. |
| <i>Phascolarctos cinereus</i> | Koala (Breeding) | N/A | High | V | V | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. Habitat present is substantially degraded such that this species is unlikely to utilise the site for breeding. |
| <i>Pimelea curviflora</i> var. <i>curviflora</i> | Pimelea curviflora var. <i>curviflora</i> | N/A | High | V | V | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Pomaderris prunifolia</i> – endangered population | Endangered population in Parramatta, Auburn, Strathfield and Bankstown LGA | N/A | High | E | V | Excluded The subject site is not located within the LGA for this endangered population. Furthermore, the presence of this species was not identified (conspicuous species) and it was |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|-------------------------------|--------------------------------------|---|---------------------------|--------------------|---------------------|---|
| | | | | | | determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |
| <i>Pommerhelix duralensis</i> | Dural Woodland Snail | Other Leaf litter and shed bark or within 50m of litter or bark Rocky areas Rocks or within 50m of rocks Fallen/standing dead timber including logs Including logs and bark or within 50m of logs or bark | High | E | E | Excluded Habitat present is substantially degraded such that this species is unlikely to utilise the subject site. Habitat in the subject site is isolated and disturbed with a higher likelihood of this species using adjoining vegetation in better condition. Additionally, this species has specific habitat requirements which were not recorded within the subject site. |
| <i>Pseudophryne australis</i> | Red-crowned Toadlet | N/A | Moderate | V | Not Listed | Excluded Habitat features associated with this species are not present on the subject site. The subject site does not contain suitable drainage lines for this species to utilise the site. |
| <i>Pteropus poliocephalus</i> | Grey-headed Flying-fox (Breeding) | N/A | High | V | V | Excluded This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain any breeding sites that are suitable for the species to utilise. |
| <i>Rhodamnia rubescens</i> | Scrub Turpentine | - | High | CE | Not Listed | Excluded The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site. |

| Species | Common Name | Habitat constraints/ Geographic limitations | Sensitivity to gain class | NSW listing status | EPBC Listing status | Justification if species excluded |
|---|--|--|---------------------------|--------------------|---------------------|--|
| <i>Syzygium paniculatum</i> | Magenta Lilly Pilly | N/A | Moderate | E | V | <p>Excluded</p> <p>The presence of this species was not identified (conspicuous species) and it was determined that the habitat is substantially degraded such that this species is unlikely to utilise the subject site.</p> |
| <i>Tetratheca glandulosa</i> | Tetratheca glandulosa | N/A | High | V | Not Listed | <p>Excluded</p> <p>Habitat features associated with this species (such as ridgetops with shale-sandstone transitional soils) are not present on the subject site. The field surveys were conducted during flowering times (July – November) however, this species and its habitat was not identified within the subject site.</p> |
| <i>Tyto novaehollandiae</i> | Masked Owl (Breeding) | N/A | High | V | Not Listed | <p>Excluded</p> <p>This is a dual credit species, and only a species credit species when specific habitat constraints are present for breeding. The subject site does not contain habitat such as trees with large hollows that are suitable for the species to utilise the site for breeding.</p> |
| <i>Wahlenbergia multicaulis</i> – endangered population | Tadgell's Bluebell in the LGAs of Auburn, Bankstown, Baulkham Hills, Canterbury, Hornsby, Parramatta and Strathfield | Check for updated LGA names | High | E | Not Listed | <p>Excluded</p> <p>This species has two populations recorded in northern Sydney (Thornleigh and Mt Ku-ring-gai), which does not include the subject site area. Habitat features associated with this species includes Hawkesbury soil landscapes, which was not recorded within the Subject site.</p> |

CE = Critically Endangered; E = Endangered; E2 = Endangered Population; V = Vulnerable

2.6.1 Targeted surveys

Due to the high level of modification of vegetation within the subject site and lack of potential habitat, targeted surveys were not conducted for species credit species. Justification for the exclusion of species credit species is provided Table 13.

However, targeted surveys were conducted for potential roosting habitat for threatened and non-threatened microbat species within the roof cavity of one of the residential dwellings to be demolished (Figure 5). Some microbat species are dual credit species which only breeding habitat considered for Species credits. None of the dual credit species are known to breed in man-made structures such as roof cavities. However, under Section 9.2.1 of the BAM, the accessor must take into consideration Prescribed Biodiversity Impacts including any man-made structures which may be roosting habitat for the following threatened microbat species:

- *Saccolaimus flaviventris* (Yellow-bellied Sheath-tail Bat)
- *Falsistrellus tasmaniensis* (Eastern False Pipistrelle)
- *Miniopterus australis* (Little Bent-winged bat)
- *Miniopterus orianae oceanensis* (Large Bent-winged bat).

The methodology and results for the microbat surveys are detailed in the Prescribed Biodiversity Impact Assessment Section 3.1.3.

2.6.2 Use of local data

The use of local data is not proposed.

2.6.3 Expert reports

Expert reports have not been prepared as part of this BDAR.

3. Stage 2: Impact assessment (biodiversity values)

3.1 Avoiding impacts

3.1.1 Locating a project to avoid and minimise impacts on vegetation and habitat

The proposed Concept Plan and Stage 1 development has utilised the existing building footprint and will involve modification to existing building or construction of new development in predominately cleared or fragmented environments. Areas of high biodiversity values have been retained where possible. Justifications on how the development aims to avoid and minimise impacts is outlined in Table 14.

Table 14: Locating a project to avoid and minimise impacts on vegetation and habitat

| Approach | How addressed | Justification |
|--|---|--|
| Locating the project in areas where there are no biodiversity values | The project (i.e. the proposed development footprint) has utilised existing development areas, cleared land and planted gardens to minimise impacts on areas with the highest biodiversity value. Areas of biodiversity values have been retained where possible within the subject site. | <p>Prior to the preparation of the Concept Plan ELA prepared an ecological constraints assessment which identified potential biodiversity values and constraints. This information was fed into the Concept Plan to ensure that the proposed development avoided impacts upon areas of highest biodiversity values, where possible. An updated Masterplan has further refined the impact areas and has excluded impacts to vegetation zone 3 which is listed as a TEC of high biodiversity value.</p> <p>The Masterplan has concentrated the development in the northern section of the subject site to reduce impacts to areas of high biodiversity values. The project has utilised areas with existing development, open space (i.e. the sporting oval) and areas of landscaped plantings and minimised the removal of native vegetation from within the site. While a small amount of native weedy vegetation (PCT 1281) will be removed (0.05 ha) this area is in low condition due to weeds. The remaining vegetation to be removed includes urban landscaped vegetation (0.31 ha) and planted native vegetation PCT 1237 (0.35 ha) which is not considered remnant native vegetation.</p> <p>Areas with higher biodiversity values including remnant native vegetation PCT 1237 and PCT 1281 will be retained within the subject site.</p> <p>The project has ensured that no hollow-bearing trees will be removed.</p> |
| Locating the project in areas where the native vegetation or threatened species habitat is in the poorest condition | The project has been located to utilise areas where native vegetation and threatened species habitat is in the poorest condition. | The project has been located to utilise areas in the north of the subject site comprised of existing buildings, cleared lands, exotic plantings and lower condition vegetation. This placement minimises removal of vegetation from the south of the subject site which contains higher quality remnant vegetation and potential threatened species habitat. |
| Locating the project in areas that avoid habitat for species and vegetation in high threat categories (e.g. an EEC or CEEC), indicated by the | The project has been located to avoid removal of vegetation in high threat categories. | The project has concentrated the impacts in vegetation which are not listed as part of a TEC. The majority of the vegetation within the development footprint (4.34 ha) does not conform to a TEC. The majority of the vegetation retained in the subject site does contain vegetation mapped as part of a TEC. |

| Approach | How addressed | Justification |
|--|--|--|
| biodiversity risk weighting for a species | | <p>The subject site has utilised areas of lower biodiversity value where possible. Areas of lower biodiversity value includes planted vegetation.</p> <p>Under the project, only minor amount of vegetation in high threat categories (EEC or CEEC) will be removed. A small amount (0.001 ha) of PCT 1237 (Blue Gum High Forest) in the form of one single remnant <i>Eucalyptus pilularis</i> has been mapped under the BC Act, will require the trimming of several outer branches for the project. The tree is located 10 m from the proposed buildings and will be retained within the subject site with careful mitigation measures.</p> <p>A small amount of fringing vegetation within zone 3 PCT 1281 (0.003 ha) will be removed to accommodate the new sporting fields. This area is of lower quality due to the edge effects, weed infestation and lower native resilience in the soil seedbank. Much of this edge has been established from revegetation works. This patch is part of a larger patch of mixed condition PCT 1281 which has been listed as a TEC under the BC Act and EPBC Act.</p> |
| Locating the project such that connectivity enabling movement of species and genetic material between areas of adjacent or nearby habitat is maintained | The project has been located to enable connectivity across the local area. | The project has been located to maintain all current connectivity between areas of vegetation. This will enable continued connectivity across the landscape for mobile fauna species and movement of genetic material. |

3.1.2 Designing a project to avoid and minimise impacts on vegetation and habitat

The development has been designed in a way which avoids and minimises impacts as outlined in Table 15.

Table 15: Designing a project to avoid and minimise impacts on vegetation and habitat

| Approach | How addressed | Justification |
|--|---|---|
| Reducing the clearing footprint of the project | The project has been designed to reduce the clearing footprint of the project. | The placement of the subject site footprint has been strategically designed to avoid high biodiversity value areas and utilises mainly cleared or built lands and planted vegetation. |
| Locating ancillary facilities in areas where there are no biodiversity values | Ancillary features have been located in areas where there are no biodiversity values. | <p>Ancillary features will be located at the northern side of the subject site in predominantly cleared areas with limited biodiversity value, avoiding the native vegetation along the southern extent of the subject site. The sporting fields will act as part of the asset protection zone and will reduce the requirement for removal of vegetation within areas of high biodiversity value in the south of the subject site</p> <p>Additionally, temporary ancillary features required during construction (such as stockpiles) will be located in existing cleared</p> |

| Approach | How addressed | Justification |
|---|--|--|
| | | areas such as the sporting fields and will not involve the removal of vegetation. |
| Locating ancillary facilities in areas where the native vegetation or threatened species habitat is in the poorest condition (i.e. areas that have a lower vegetation integrity score) | Ancillary features have been located in areas where native vegetation is in the poorest condition. | Ancillary features will be located at the northern portion of the subject site in predominantly built or cleared lands or in areas where native vegetation has been planted. Native planted vegetation contains a lower vegetation integrity score than the remnant vegetation located in the south of the subject site. |
| Locating ancillary facilities in areas that avoid habitat for species and vegetation in high threat status categories (e.g. an EEC or CEEC) | Ancillary features have been located in areas that avoid habitat for species and vegetation in high threat categories. | The majority of the subject site contains exotic and planted native vegetation which does not support vegetation in high threat categories (e.g. EEC or CEEC). The subject site contains substantial amount of cleared lands which will be utilised for temporary ancillary facilities and will not impact upon high threat category vegetation. |
| Providing structures to enable species and genetic material to move across barriers or hostile gaps | The development has been designed to maintain a vegetated corridor enabling movement of species and genetic material. | The project has been designed to retain quality vegetation in the south of the subject site. The subject site has been designed so that it does not impact on corridors. Existing vegetated corridors will be maintained with connectivity in all directions, allowing for the continued movement of species and genetic material across the landscape. Given that no corridors will be impacted, additional structures are not necessary. |
| Making provision for the demarcation, ecological restoration, rehabilitation and/or ongoing maintenance of retained native vegetation habitat on the subject site. | Vegetation in the east of the subject site will be retained, enhanced and maintained. | Vegetation to be retained in the subject site (to the south of the subject site), will be enhanced and maintained as part of weed removal works. |
| Efforts to avoid and minimise impacts through design must be documented and justified | The project has been designed to reduce the clearing footprint of the project. | The placement of the subject site footprint has been strategically designed to avoid high biodiversity value areas and utilises mainly cleared lands and degraded vegetation. |

3.1.3 Prescribed biodiversity impacts

The list of potential prescribed biodiversity impacts as per the BAM is provided below:

- Occurrences of karst, caves, crevices and cliffs - none occur within the subject site
- Occurrences of rock - no rock outcrops or scattered rocks occur within the subject site
- Occurrences of human made structures and non-native vegetation – **Yes, see section below.**
- Hydrological processes that sustain and interact with the rivers, streams and wetlands – **Yes, the headwaters to a first order Strahler stream is located in the south-west corner of the subject site (Figure 2).**

- Proposed development for a wind farm and use by species as a flyway or migration route - the project does not involve any wind farm development.

The subject site contains both human made structures and non-native vegetation. Additional information regarding consideration of human made structures is provided below. Non-native vegetation was identified and assessed for any potential to provide habitat for threatened flora and fauna species, including presence of HBTs. The subject site contains hydrological processes as seen in Figure 2. The subject site has the prescribed biodiversity impacts as outlined in Table 17.

A literature review was conducted to identify if buildings or structures that could potentially be utilised as a roosting resource by microchiropteran bats (microbats), resources such as relevant literature and BioNet records of the subject site and surrounding landscape were also utilised during the desktop review. Field surveys were conducted to visually determine if the buildings within the subject site contain potential openings, possibly utilised by microbats. Possible threatened microbats surveyed for include:

- *Saccolaimus flaviventris* (Yellow-bellied Sheath-tail Bat)
- *Falsistrellus tasmaniensis* (Eastern False Pipistrelle)
- *Miniopterus australis* (Little Bent-winged bat)
- *Miniopterus orianae oceanensis* (Large Bent-winged bat).

A historic dwelling was identified with two holes which may provide microbat access into the roof cavity. An anabat ultrasonic device was placed inside the roof cavity for three consecutive nights (9 – 11 November 2018) to determine the presence/absence of microbats and the type of species if present. A visual inspection was also conducted with a high-powered torch; however, this method has limitations due to the cryptic nature of microbats to hide in small cavities in the roof. The weather conditions during field surveys were optimal, mild with no rainfall (Table 16). The targeted surveys were conducted within the seasonal survey period for potential microbat species.

The recordings from the anabat device were analysed by ELA's fauna ecologist Rodney Armistead. No microbat species were recorded during the targeted surveys. It is unlikely that microbat species utilise this dwelling for roosting or breeding habitat.

Table 16: Weather conditions during anabat targeted survey

| Date | Rainfall (mm) | Minimum temperature °C | Maximum temperature °C |
|------------------|---------------|------------------------|------------------------|
| 9 November 2018 | 0 | 7.5 | 22.5 |
| 10 November 2018 | 0 | 15.3 | 23.2 |
| 11 November 2018 | 0 | 10.0 | 24.8 |

Sourced from Bureau of Meteorology station number 066124 Parramatta

Table 17: Prescribed biodiversity impacts

| Prescribed biodiversity impact | Description in relation to the subject site | Threatened species or ecological communities effected |
|---|--|--|
| Impacts of development on the habitat of threatened species or ecological communities associated with human made structures. | The subject site contains a number of existing buildings. The majority of the buildings are recently constructed (within the last 20 years) and do not provide potential microbat roosts. There was one historic building present, however, targeted surveys did not record microbat roosting within the roof cavity. | Potential roosting habitat for threatened microbat <i>Saccolaimus flaviventris</i> (Yellow-bellied Sheath-tail Bat) and <i>Falsistrellus tasmaniensis</i> (Eastern False Pipistrelle), <i>Miniopterus australis</i> (Little Bent-winged bat) and <i>Miniopterus schreibersii oceanensis</i> (Eastern Bent-winged bat) although none were recorded during targeted surveys. |
| Impacts of development on the habitat of threatened species or ecological communities associated with non-native vegetation | <p>The subject site contains nectar producing non-native vegetation canopy, in formal gardens which will be removed as part of the proposed development.</p> <p>The subject site contains non-native vegetation for common urban arboreal mammals (possums) which provides foraging opportunities for threatened nocturnal bird species. The development will result in a reduction in the extent of foraging habitat and reduction in availability of their prey items.</p> | <p>Potential foraging habitat for other threatened microbat species above non-native vegetation canopy.</p> <p>Potential foraging habitat for <i>Pteropus poliocephalus</i> (Grey-headed Flying Fox (GHFF)).</p> <p>Potential foraging habitat for <i>Ninox strenua</i> (Powerful Owl).</p> |
| Impacts of development on the connectivity of different areas of habitat of threatened species that facilitates the movement of those species across their range | <p>The proposed development will require the removal of non-native vegetation from within the subject site.</p> <p>The development will result in a minor reduction in the extent of existing non-native vegetation within the subject site which provides stepping stone habitat between urban fragmented patches of vegetation.</p> | <p>Reduction in extent of potential foraging habitat for GHFF.</p> <p>Reduction in extent of potential habitat for Powerful Owl.</p> <p>Reduction in extent of foraging habitat for other threatened microbats.</p> |
| Impacts of development on movement of threatened species that maintains their lifecycle | The proposed development will result in reduction of vegetation within the subject site and marginal loss of connectivity for mobile threatened species. | GHFF, Powerful Owl and microbat species. |
| Impacts of development on water quality, water bodies and hydrological processes that sustain threatened species and threatened ecological communities (including from subsidence or upsidence resulting from underground mining) | The proposed works is located upslope of the 1 st order stream and may result in a decline of water quality. | <p>The 1st order stream does not support water dependent threatened species or water dependent ecological communities.</p> <p>Sydney Turpentine Ironbark Forest TEC is located upslope of the 1st order stream and is <u>not</u> depend upon hydrological flows.</p> |

3.1.3.1 Locating a project to avoid and minimise prescribed biodiversity impacts

The development has been located in a way which avoids and minimises prescribed biodiversity impacts as outlined in Table 18.

Table 18: Locating a project to avoid and minimise prescribed biodiversity impacts

| Approach | How addressed | Justification |
|--|---|--|
| Locating the envelope of surface works to avoid direct impacts on the habitat features | Habitat features including one HBT, foraging habitat for GHFF, Powerful Owl and threatened microbats within the subject site will be removed. | The development has avoided impacts to large tracts of vegetation in the south which includes nectar producing native canopy species for GHFF, foraging habitat for Powerful Owl and microbat species. The development has been located in a way to avoid impact to hollow-bearing trees (HBT). Targeted surveys have been conducted to ensure that the development will not result in the loss of roosting habitat for Yellow-bellied Sheath tailed Bat or other species which utilised man-made structures. |
| Locating the envelope of sub-surface works, both in the horizontal and vertical plane, to avoid and minimise operations beneath the habitat features, e.g. locating long wall panels away from geological features of significance or water dependent plant communities and their supporting aquifers | The development will involve the construction of underground carpark | The underground carparks are located in an existing cleared area which does not contain habitat features and is located more than 200 m from the 1 st order stream. |
| Locating the project to avoid severing or interfering with corridors connecting different areas of habitat, migratory flight paths to important habitat or preferred local movement pathways | The development will involve the removal of some native and exotic vegetation which forms a connective corridor along the eastern perimeter. | Although the development has result in the removal of some native and exotic vegetation along the eastern perimeter, the connectivity will be retained through street verge plantings along the eastern perimeter (Mount Pleasant Avenue). Additionally, the impacts have utilised this area of planted native and exotic vegetation and retained the patch of TEC located along Osbourne Road along the western perimeter instead. The patch of vegetation along the western perimeter will ensure that connectivity is retained. |
| Optimising project layout to minimise interactions with threatened and protected species and ecological communities, e.g. designing turbine layout to allow buffers around features that attract and support aerial species, such as forest edges, | The development has been strategically placed to avoid impacts to areas of high biodiversity value. | The development has utilised the northern portion of the subject site which includes cleared lands and exiting development footprint and vegetation of low biodiversity values and retained areas of high biodiversity values in the south of the subject site |

| Approach | How addressed | Justification |
|---|---|--|
| riparian corridors and wetlands, ridgetops and gullies | | which includes areas of TEC of high quality (STIF). |
| Locating the project to avoid direct impacts on water bodies | The development has been strategically placed in the north of the subject site to avoid direct impacts to waterbodies located in the south. | The subject site will not directly impact upon the 1 st order stream located in the south-western corner of the subject site. |

3.1.3.2 Designing a project to avoid and minimise prescribed biodiversity impacts

The development has been designed in a way which avoids and minimises prescribed biodiversity impacts as outlined in Table 19.

Table 19: Designing a project to avoid and minimise prescribed biodiversity impacts

| Approach | How addressed | Justification |
|---|---|--|
| Design of project elements to minimise interactions with threatened and protected species and ecological communities, e.g. designing turbines to dissuade perching and minimise the diameter of the rotor swept area, designing fencing to prevent animal entry to transport corridors | The development design has retained TEC within the subject site and utilised areas with minimal impacts to biodiversity values. | The development design has utilised existing disturbed areas to minimise interactions with threatened species and minimised impacts to TECs located in the south of the subject site which will be retained under the project. |
| Design of the project to maintain hydrological processes that sustain threatened species and TECs | There are no threatened species and TECS which are depend upon hydrological processes. | The development design is located away from hydrological process and is not anticipated to alter the current hydrological flow regime of the unnamed 1 st order stream located in the south-west corner of the subject site. There are no threatened species and TECs which are dependent upon hydrological processes identified within the subject site or subject site. |
| Design of the project to avoid and minimise downstream impacts on rivers, wetlands and estuaries by control of the quality of water released from the site. | The works are unlikely to alter the current hydrological flow of the unnamed 1 st order stream. | The development design has been conducted so that hydrological flows will be captured on the sporting fields prior to entering the native vegetation located into the south of the subject site and into the 1 st order stream. |

3.2 Assessment of Impacts

3.2.1 Direct impacts

The direct impacts of the development on:

- native vegetation are outlined in Table 20
- threatened ecological communities are outlined in Table 21
- removal of one hollow-bearing tree
- prescribed biodiversity impacts are outlined in Section 3.2.2.

Direct impacts including the final project footprint (construction and operation) are shown on Figure 7.

Table 20: Direct impacts to native vegetation

| PCT ID | PCT Name | Vegetation Class | | | Vegetation Formation | | | Direct impact (ha) |
|--------|-----------------------------------|-------------------------------------|------------|--|---|--|--|--------------------|
| 1237 | Blue Gum High Forest | North Coast Wet Sclerophyll Forests | | | Wet Sclerophyll Forests (Shrubby sub-formation) | | | 0.35 |
| 1281 | Sydney Turpentine-Ironbark Forest | Northern Wet Sclerophyll Forests | Hinterland | | Wet Sclerophyll Forests (Grassy sub-formation) | | | 0.05 |

Table 21: Direct impacts on threatened ecological communities

| PCT ID | BC Act | | | EPBC Act | | |
|--------|----------------|-----------------------------------|--------------------|----------------|----------------------------|--------------------|
| | Listing status | Name | Direct impact (ha) | Listing status | Name | Direct impact (ha) |
| 1237 | CEEC | Blue Gum High Forest | 0.001 | N/A* | N/A* | N/A* |
| 1281 | EEC | Sydney Turpentine-Ironbark Forest | 0.05 | CEEC | Turpentine Ironbark Forest | 0.05 |

* Note the PCT 1237 represented in the subject site did not satisfy listing requirements under the EPBC Act criteria

3.2.2 Change in vegetation integrity

The change in vegetation integrity as a result of the development is outlined in Table 22.

Table 22: Change in vegetation integrity

| Veg Zone | PCT ID | Condition | Area (ha) | Current vegetation integrity score | Future vegetation integrity score | Change in vegetation integrity |
|----------|--------|----------------|-----------|------------------------------------|-----------------------------------|--------------------------------|
| 2 | 1237 | Native planted | 0.35 | 35.1 | 0 | -35.1 |
| 4 | 1281 | Weedy | 0.05 | 39.7 | 0 | -39.7 |

3.2.3 Indirect impacts

The indirect impacts of the development are outlined in Table 23.

Table 23: Indirect impacts

| Indirect impact | Project phase | Nature | Extent | Frequency | Duration | Timing |
|--|--------------------------|--|---|--|---|-------------------------------|
| sedimentation and contaminated and/or nutrient rich run-off | Construction | Runoff during construction works | Confined to subject site with sediment fencing | During heavy rainfall or storm events | During rainfall events | Short-term impacts |
| noise, dust or light spill | Construction | Noise and dust created from machinery (no night works proposed therefore no light spill) | Noise and dust likely to carry beyond subject site boundary | Daily, during construction works | Sporadic throughout construction period | Short-term impacts |
| inadvertent impacts on adjacent habitat or vegetation | Construction | Damage to adjacent habitat or vegetation | Adjacent vegetation | Daily, during construction works | Throughout construction period | Short-term impacts |
| transport of weeds and pathogens from the site to adjacent vegetation | Construction | Spread of weed seed or pathogens | Potential for spread into adjacent habitat | Daily, during construction works | Sporadic throughout construction period | Potentially long-term impacts |
| vehicle strike | Construction / operation | Potential for native fauna to be struck by working machinery and moving vehicles | Within access road and subject site | Daily, during both construction and operational phases. | Throughout life of project | Short-term impacts |
| trampling of threatened flora species | Construction / operation | No threatened flora species present | N/A | N/A | N/A | N/A |
| rubbish dumping | Construction / operation | Illegal dumping by local residents/ construction crews | Potential for rubbish to spread via wind into adjacent vegetation | Potential to occur at any time throughout construction or operational phases | Throughout life of project | Short-term impacts |
| wood collection | Construction / operation | Removal of wood in vegetation located in the southern extent of the subject site | In southern portion of the subject site | Potential to occur at any time throughout construction or operational phases | Throughout life of project | Short-term impacts |

| Indirect impact | Project phase | Nature | Extent | Frequency | Duration | Timing |
|--|--------------------------|---|---|--|--|----------------------------------|
| bush rock removal and disturbance | Construction / operation | Removal of rocks in southern vegetation within the subject site | In vegetation in the southern portion of the subject site | Potential to occur at any time throughout construction or operational phases | Throughout life of project | Short-term impacts |
| increase in predatory species populations | Construction / operation | Potential increase in domestic predatory species due to reduction of vegetation | In vegetation in the southern portion of the subject site | During operational phase | Potential at any point during operation of development | Short-term impacts |
| increase in pest animal populations | Construction / operation | Potential to increase if introduced | In vegetation in the southern portion of the subject site | Potential to occur at any time throughout construction or operational phases | Throughout life of project | Short-term impacts |
| increased risk of fire | Construction / operation | Potential due to presence of vegetation retained in the south of the subject site | In vegetation in the southern portion of the subject site | Potential to occur at any time, although, more likely during dry, windy conditions | Throughout life of project | Short-term and long-term impacts |
| disturbance to specialist breeding and foraging habitat, e.g. beach nesting for shorebirds. | Construction / operation | Runoff during construction works | Confined to subject site with sediment fencing | During heavy rainfall or storm events | During rainfall events | Short-term impacts |

3.2.4 Prescribed biodiversity impacts

An assessment of impacts of the development on prescribed biodiversity impacts is outlined in Table 24 in accordance with Section 9.2.1 of the BAM.

Table 24: Direct impacts on prescribed biodiversity impacts

| BAM Criteria | Justification |
|--|--|
| 9.2.1.3 The assessment of the impacts of the development on the habitat of threatened species or ecological communities associated with human made structures | |
| a) identify the human made structures with potential to be habitat for threatened species or ecological communities | <p>The subject site is located within a highly urbanised area. The development will occur over a 30 year staged development. Stage 1 works will involve the demolition of three residential dwellings for the proposed new boarding house. The remaining Concept Plan will result in the removal of and redesign of a number of multistorey educational buildings which do not contain small gaps suitable for microbat access into the roof cavity.</p> <p>Only one of the residential dwellings contained potential access for microbats; no microbats were identified during targeted surveys or were considered likely to utilise the dwelling as potential roosting habitat.</p> <p>No other human made structures with potential habitat for threatened species or ecological communities were identified in the subject site.</p> |
| b) identify the species and ecological communities likely to use the habitat | <p>The residential dwelling provides potential roosting habitat for a number of threatened microbat species including: <i>Saccolaimus flaviventris</i> (Yellow-bellied Sheath-tail Bat) and <i>Falsistrellus tasmaniensis</i> (Eastern False Pipistrelle), <i>Miniopterus australis</i> (Little Bent-winged bat) and <i>Miniopterus orianae oceanensis</i> (Large Bent-winged bat). These species are known to occasionally roost in buildings. There are BioNet records for these species within a 5 km radius for these species. Targeted surveys did not record evidence of microbat activity within the building.</p> |
| c) describe the nature, extent and duration of short and long-term impacts | <p>The impact involves the permanent removal of three residential dwellings and several multistorey education facilities, of which only one residential dwelling provided potential roosting habitat for microbat species which may utilise it occasionally. The removal of this dwelling may result in a loss of potential roost habitat for microbat species; however, targeted survey did not record evidence of microbat regularly utilising this dwelling. As such these impacts are likely to be minor and alternative roost locations which may occur within the subject site are likely to be used by microbats under these circumstances.</p> |
| d) describe, with reference to relevant literature the importance within the bioregion of the habitat of these species or ecological communities | <p>According to literature documented in Australian Bat (Churchill 2009) the preferred roosting habitat of the following species includes:</p> |

| BAM Criteria | Justification |
|--|--|
| | <ul style="list-style-type: none"> • Yellow-bellied Sheathtail Bat – this species will utilise tree hollows or buildings in small groups. There is potential that this species may utilise the building and tree hollows recorded within the subject site and within the subject site. • Eastern False Pipistrelle – this species primarily roosts in tree trunks in small groups, however it may occasionally utilise wooden buildings. It is unlikely this species would utilise buildings for maternity roosts due to the presence of hollow-bearing trees within the subject site. • Little Bent-winged Bat – this species forms specific maternity roosts in caves. They occasionally utilise buildings in the absence of other alternative roost locations (such as mines, culverts). There is potential that this species may on occasion utilise the residential dwelling as an alternative roost location. • Large Bent-winged Bat – this species primarily roosts in caves, however, it occasionally roosts in human made structures such as buildings. There is potential that this species may on occasion utilise the residential dwelling as an alternative roost location. <p>Targeted surveys did not record evidence of microbat activity within the building.</p> |
| <p>e) predict the consequences of the impacts for the local and bioregional persistence of the suite of threatened species and communities likely to use these areas as habitat, with reference to relevant literature and other published sources of information.</p> | <p>While these species of microbats have been known to utilise human structures for roosting, preferred roosting habitat for these species are non-human made structures (tree hollows or caves). Additionally, only one of the species is likely to utilise buildings more regularly including breeding times, this species is the Yellow-bellied Sheathtailed Bat. The other species of microbats may utilise the residential dwelling on occasion while traversing through the landscape or if other alternative roosting resources are not present. It should be noted that the subject site provides marginal foraging and alternative roosting habitat in the form of buildings for a number of microbat species. The subject site does not contain important habitat for these species.</p> <p>There is potential that the removal of the residential building may impact upon the number of available roosting resources for microbats migrating to breeding or non-breeding habitats such as the two Bent-winged species. There is no available literature which has considered the impacts of removal of human made structures on microbat species.</p> <p>The Priority Action Statement for the Yellow-bellied Sheathtail-bat lists several recommended actions for help in the recovery of this species, those pertaining to retention of roosting habitat focus on the retention of large hollow-bearing trees and retention of vegetated areas. The Priority Action Statement for the Little Bent-winged Bat and Large Bent-winged Bat include further investigation of the wintering roosts for these species which includes tree hollows and undertaking restoration activities to create habitat and connectivity in the landscape. There is no mention of the use of buildings for Bentwing Bat species.</p> <p>The habitat within the subject site is unlikely to be important for any of these microbat species.</p> |

| BAM Criteria | Justification |
|--|---|
| 9.2.1.4 The assessment of the impacts of development on the habitat of threatened species or ecological communities associated with non-native vegetation | |
| a) identify the species and ecological communities likely to use the habitat | Several non-native tree species are present in the subject site which have been planted within residential gardens or are invasive weeds. Non-native species which have been identified as potential foraging species for Grey-headed Flying fox are <i>Syagrus romanzoffiana</i> (Cocos Palm), <i>Liquidambar styraciflua</i> , <i>Ligustrum lucidum</i> (Broad-leaved Privet) and <i>Ligustrum sinense</i> (Small-leaved Privet). Additionally, non-native vegetation may be utilised by arboreal mammals which comprise prey resources for Powerful Owl. |
| (b) describe the nature, extent and duration of short and long-term impact | The proposed development will result in the permanent removal of a small number of non-native trees (listed above) which provide potential foraging habitat for Grey-headed Flying-fox and marginal foraging habitat for the threatened microbat species and foraging habitat for Powerful Owl prey resources. |
| (c) describe, with reference to relevant literature and other reliable published sources of information, the importance within the bioregion of the habitat to these species or ecological communities | <p>These non-native foraging species are in relatively low abundance within the subject site and most species would provide only small amounts of secondary foraging habitat. Flowering resources in the form of native planted <i>Eucalyptus</i>, <i>Melaleuca</i> and <i>Callistemon</i> sp. would more likely be utilised for foraging resources by Grey-headed Flying-fox.</p> <p><i>Syagrus romanzoffiana</i> is known to cause injury and death to flying-foxes due to toxicity of the green fruits and bats becoming entangled in the flower sheaths and frond leaves which can cause death, injury and distress. The removal of this species is therefore likely to be beneficial to the species.</p> |
| (d) predict the consequences of the impacts for the local and bioregional persistence of the suite of threatened species and communities likely to use these areas as habitat, with reference to relevant literature and other published sources of information. | <p>The consequences of the permanent removal of those species listed above for the local and bioregional persistence of the Grey-headed Flying-fox is predicted to be negligible.</p> <p>Several of the non-native species to be removed (<i>Syagrus romanzoffiana</i>, <i>Ligustrum sinense</i>, <i>Ligustrum lucidum</i>) are listed as environmental weeds in the Greater Sydney Regional Strategic Weed Management Plan (2017-2022). There is an abundance of similar habitat within the locality and bioregion, and an abundance of higher quality habitat in the locality and bioregion.</p> |
| 9.2.1.5 The assessment of the impacts of development on the connectivity of different areas of habitat of threatened species that facilitates the movement of those species across their range must: | |
| (a) identify the area/s of connectivity joining different areas of habitat that intersect with the subject land and the areas of habitat that are connected according to Paragraph 4.2.1.3 | The subject site includes predominately disturbed and non-native species. To the south of the development is a large tract of native vegetation retained within the subject site. This area of vegetation connects to riparian corridor along Coups Creek which eventually flows into Lane Cove River and the National Park. To the north, the subject site abuts a major arterial road, Pennant Hills Road, which intersects small patches of urban native/exotic vegetation. 500 m north of the subject site lies vegetative tributaries to Waitara Creek which flows into Berowra Valley Regional Park. |

| BAM Criteria | Justification |
|---|---|
| | <p>The vegetation within the subject site is relatively small compared to the native vegetation retained within the subject site and the vegetation connected in the broader landscape. However, due to the presence of major roads, only highly mobile species are likely to utilise the vegetation within the subject site.</p> |
| (b) identify the species and ecological communities likely to benefit from the connectivity | <p>The species most likely to utilise the connectivity would be Grey-headed Flying-fox, microbat species and disbursal of juvenile Powerful Owl.</p> <p>BGHF and STIF species are likely to benefit from the connectivity within the subject site with adjacent riparian vegetation along Coups River riparian corridor to the south of the subject site.</p> |
| (c) describe the nature, extent and duration of short and long-term impacts | <p>The proposed development will result in the permanent removal of 0.40 ha of native and 0.31 ha of exotic vegetation which forms connecting habitat for highly mobile species. Connectivity will be retained within the subject site and in the adjacent broader locality.</p> |
| (d) describe, with reference to relevant literature and other reliable published sources of information, the importance of the area of connectivity within the bioregion | <p>The connectivity is considered limited except for highly mobile species which easily move across disturbed landscapes. The connecting habitat provides potential foraging habitat for the above listed species, which is part of a fragmented network of urban vegetation within the eastern suburbs. Within the Sydney Basin Bioregion, the removal of 0.40 ha of native and 0.31 ha of exotic vegetation is considered to provide negligible connectivity on a landscape scale. The removal of connecting habitat would not prevent the highly mobile Grey-headed Flying-fox, Powerful Owls or microbats from moving across the landscape in search of foraging resources.</p> <p>The removal of a small amount of connecting habitat from the subject site is unlikely to be of importance to any threatened species within the bioregion considering the availability of connectivity retained within the subject site and immediately adjacent to the subject site.</p> |
| (e) predict the consequences of the impacts for the bioregional persistence of the suite of threatened species and communities currently benefitting from the connectivity with reference to relevant literature and other published sources of information and taking into consideration mobility, abundance, range and other relevant life history factors. | <p>The habitat to be removed forms part of a network or stepping stone habitat in the form of canopy, shrubs and ground layer garden plantings and native species. The vegetation connectivity flows from the south where large tracts of vegetation are retained and abruptly ends in along the northern boundary of the subject site which is bounded by Pennant Hills Road. Only highly mobile species are likely to utilise the stepping stone vegetation from the subject site north. Under the proposal, canopy species will be retained within the subject site to provide connectivity. The proposed development will not result in a loss of connectivity for the highly mobile species likely to utilise it.</p> |

3.2.5 Mitigating and managing impacts

Measures proposed to mitigate and manage impacts at the subject site before, during and after construction are outlined in Table 25.

Table 25: Measures proposed to mitigate and manage impacts

| Measure | Risk before mitigation | Risk after mitigation | Action | Outcome | Timing | Responsibility |
|---|------------------------|-----------------------|--|--|------------------------------------|-----------------------------|
| Displacement of resident fauna | Minor | Negligible | Pre-clearance survey of trees to be removed and identification/location of habitat trees by a suitably qualified ecologist. Supervision by a qualified ecologist/licensed wildlife handler during tree removal in accordance with best practise methods. | Resident fauna relocated in a sensitive manner | Prior to and during clearing works | Project Manager / Ecologist |
| Timing works to avoid critical life cycle events such as breeding or nursing | Minor | Negligible | Avoid clearing works in later winter/spring during breeding/nesting period for birds | Impacts to fauna during nesting/nursing avoided | During clearing works | Project Manager |
| Instigating clearing protocols including pre-clearing surveys, daily surveys and staged clearing, the presence of a trained ecological or licensed wildlife handler during clearing events | Moderate | Minor | Pre-clearance survey of trees to be removed and identification/location of habitat trees by a suitably qualified ecologist. Trees identified for retention should be clearly delineated as a 'No Go' zone with high visibility bunting. Supervision by a qualified ecologist/licensed wildlife handler during tree removal in accordance with best practise methods. Any tree removal is to be undertaken by a suitably qualified and insured arborist. | Any fauna utilising habitat within the subject site will be identified and managed to ensure clearing works minimise the likelihood of injuring resident fauna | During clearing works | Project Manager / Ecologist |
| Installing artificial habitats for fauna in adjacent retained vegetation and habitat or human made structures to replace the habitat resources lost and encourage animals to move from the impacted site, e.g. nest boxes | Minor | Negligible | Any trees removed that have hollows/hollow trunks/fissures should be retained as ground fauna habitat and/or used as replacement hollows and attached to trees within the within the subject site/subject site. If it is impractical to use salvaged hollows as replacement tree hollows, compensatory nest boxes should be installed within vegetation to be retained. | Replacement of habitat features removed | Prior to and during clearing works | Project Manager/ Ecologist |

| Measure | Risk before mitigation | Risk after mitigation | Action | Outcome | Timing | Responsibility |
|---|------------------------|-----------------------|---|---|---|-----------------|
| clearing protocols that identify vegetation to be retained, prevent inadvertent damage and reduce soil disturbance | Moderate | Minor | Vegetation identified for retention should be clearly delineated as a 'No Go' zone with high visibility bunting. No temporary facilities i.e. site offices/toilets/soil stockpiling is to occur within tree protection zone. | Vegetation to be retained outside of the subject site boundary will not be disturbed/impacted | Demarcation of vegetation to be set up prior to any works occurring on site and to remain throughout duration of construction works | Project Manager |
| Sediment barriers or sedimentation ponds to control the quality of water released from the site into the receiving environment | Moderate | Minor | Appropriate controls are to be utilised to manage exposed soil surfaces and stockpiles to prevent sediment discharge into waterways. Soil and erosion measures such as sediment fencing, clean water diversion must be in place prior the commencement of the construction work. | Erosion and sedimentation will be controlled | For the duration of construction works | Project Manager |
| Programming construction activities to avoid impacts; for example, timing construction activities for when migratory species are absent from the site, or when particular species known to or likely to use the habitat on the site are not breeding or nesting | Minor | Negligible | Timing of construction works should be planned to occur outside of the winter/spring breeding season. | impacts to fauna during nesting/nursing avoided | During clearing works | Project Manager |
| Hygiene protocols to prevent the spread of weeds or pathogens between infected areas and uninfected areas | Moderate | Minor | Vehicles, machinery and building refuse should remain only within the subject site and not impinge on the areas of retained native vegetation to be retained in the east. Weed management to be undertaken in retained bushland following construction works. | Spread of weeds prevented | Post-construction | Project Manager |
| Staff training and site briefing to communicate environmental | Minor | Negligible | Construction staff to be briefed prior to work commencing to be made aware of sensitive biodiversity values present and environmental procedures such as: | All staff entering the Subject site are fully aware of all the | To occur for all staff entering/working | Project Manager |

| Measure | Risk before mitigation | Risk after mitigation | Action | Outcome | Timing | Responsibility |
|--|------------------------|-----------------------|--|---|---|-----------------|
| features to be protected and measures to be implemented | | | <ul style="list-style-type: none"> Importance of retained vegetation areas and 'No Go' zones Site environmental procedures (vegetation management, sediment and erosion control, exclusion fencing and noxious weeds) What to do in case of environmental emergency (chemical spills, fire, injured fauna) Key contacts in case of environmental emergency | ecological values present within the Lot and environmental aspects relating to the development and know what to do in case of any environmental emergencies | at the subject site. Site briefings should be updated based on phase of the work and when environmental issues become apparent. | |
| Development control measures to regulate activity in vegetation and habitat adjacent to residential development including controls on rubbish disposal, wood collection, fire management and disturbance to nests and other niche habitats | Minor | Negligible | <p>Strategy to be developed and implemented as part of the residential development may include:</p> <ul style="list-style-type: none"> Signage to indicate areas not to be disturbed i.e. No Go zones Rubbish disposal guidance Prohibition of wood collection Prohibition of bush rock removal Controls on pet ownership such as prohibitions on allowing pets to roam beyond fenced areas | Strategy to protect vegetation and habitat adjacent to development | To be developed to provide awareness to residents of housing development. | Client |
| Making provision for the ecological restoration, rehabilitation and/or ongoing maintenance of retained native vegetation habitat on or adjacent to the subject site | Minor | Negligible | Landscaping in the Subject site is to use locality derived native species and those found within the PCT present. | Areas within the subject site will be landscaped using appropriate species | Throughout construction and following completion of construction activities. | Project Manager |

3.2.6 Serious and Irreversible Impacts (SII)

The development has two candidate entities for Serious and Irreversible Impacts (SII) values as outlined in Table 26. Detailed consideration of whether impacts on candidate species are serious and irreversible is included for BGHF and STIF in Table 28 and Table 29.

Table 26: Candidate Serious and Irreversible Impacts

| Species / Community | Common Name | Principle | Direct impact individuals / area (ha) | Threshold |
|---|-----------------------------------|---------------------|---------------------------------------|---|
| Blue Gum High Forest in the Sydney Basin Bioregion | Blue Gum High Forest | Principals 1, 2 & 4 | 0.001 ha | Listed as 'under development' in BioNet |
| Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion | Sydney Turpentine-Ironbark Forest | Principals 1, 2 & 4 | 0.05 ha | Not yet published |

Table 27: Determining whether impacts are serious and irreversible

| Determining whether impacts are serious and irreversible | Assessment |
|---|---|
| Principle 1 | |
| Does the proposal impact on a species, population or ecological community that is a candidate entity because it is in a rapid rate of decline? | Yes |
| If yes, is the impact in excess of any threshold identified and therefore likely to be serious and irreversible? Note: where candidate entities have no listed threshold, any impact is considered likely to be serious and irreversible | The Thresholds for BGHF and STIF have not been published yet according to the Threatened Biodiversity Data Collection provided in OEH BioNet. |
| Principle 2 | |
| Does the proposal impact on a species that is a candidate entity because it has been identified as having a very small population size? | No. The proposal will not impact upon threatened flora/fauna species which are a candidate entity species because it has been identified as having a small population size. |
| If yes, is the impact in excess of any threshold identified and therefore likely to be serious and irreversible? Note: where candidate entities have no listed threshold, any impact is considered likely to be serious and irreversible | N/A |
| Principle 3 | |
| Does the proposal impact on the habitat of a species or an area of an ecological community that is a candidate entity because it has a very limited geographic distribution? | Yes |
| If yes, is the impact in excess of any threshold identified and therefore likely to be serious and irreversible? Note: where candidate entities have no listed threshold, any impact is considered likely to be serious and irreversible. | The Thresholds for BGHF and STIF have not been published yet according to the Threatened Biodiversity Data Collection provided in OEH BioNet. |
| Principle 4 | |
| | Yes |

| Determining whether impacts are serious and irreversible | Assessment |
|---|---|
| Does the proposal impact on a species, a component of species habitat or an ecological community that is a candidate entity because it is irreplaceable? | |
| If yes, is the impact in excess of any threshold identified and therefore likely to be serious and irreversible? Note: where candidate entities have no listed threshold, any impact is considered likely to be serious and irreversible. | The Thresholds for BGHF and STIF have not been published yet according to the Threatened Biodiversity Data Collection provided in OEH BioNet. |

Table 28: Evaluation of an impact on a TEC – Blue Gum High Forest

| Impact Assessment Provisions | Assessment |
|--|--|
| 1. The area and condition of the TEC to be impacted directly and indirectly by the proposed development | The development will remove 0.001 ha BGHF which is in a degraded condition. The BGHF impacted within the subject site is represented by an isolated, single tree (<i>Eucalyptus pilularis</i>) in a landscaped garden. The works will result in the removal of outer branches to accommodate a new building structure. The tree will be retained within the subject site. |
| 2. The extent and overall condition of the TEC within an area of 1500 metres, and then 5000 metres, surrounding the proposed development footprint. In the case of strategic biodiversity certification projects, the extent and overall condition of the TEC may be assessed across the IBRA sub region | Within the subject site, 0.70 ha of BGHF will be retained. In addition to what has been mapped within the subject site, there is an estimated 61.66 ha of BGHF within an area of 1,500m, in varying condition (from large tracts to small patches) (mapped by SMCMA, 2016). The removal of 0.001 ha of BGHF within the subject site represents 0.001% of the mapped BGHF extent within the 1,500 m radius. Within 5,000 m radius of the subject site, there is an estimated 369.57 ha of BGHF that has been mapped with low disturbance condition. The removal of 0.001 ha of BGHF from within the subject site, represents 0.0002% of the mapped BGHF extent within the 5,000 m radius. |
| 3. An estimate of the extant area and overall condition of the TEC remaining before and after the impact of the proposed development has been taken into consideration | The development will not result in the overall decline of the condition of BGHF after development. |
| 4. The development proposal's impact on: | |
| a. Abiotic factors critical to the long-term survival of the TEC; for example, will the impact lead to a reduction of groundwater levels or substantial alteration of surface water patterns; will it alter natural disturbance regimes that the TEC depends upon, e.g. fire, flooding etc.? | The development will not impact abiotic factors critical to the long-term survival of the TEC. |
| b. Characteristic and functionally important species through impacts such as, but not limited to, inappropriate fire/flooding regimes, removal of under-storey species or harvesting of plants | The development will not impact characteristic and functionally important species outside of the proposed impact area. |
| c. The quality and integrity of an occurrence of the TEC through threats and indirect impacts including, but not limited to, assisting invasive flora and fauna species to become established or causing regular mobilisation of fertilisers, herbicides or other chemicals or pollutants which may harm or inhibit growth of species in the TEC | The development proposal has the potential to impact upon the health of the isolated <i>Eucalyptus pilularis</i> which will remain within the subject site. The development proposal has potential to assist in the spread of invasive species into the patch of BGHF that will be retained within the subject site. These potential impacts will be controlled during the construction phase and long-term maintenance of the |

| Impact Assessment Provisions | Assessment |
|---|--|
| | Subject site. These works will retain the quality and integrity of the isolated patch of BGHF. |
| 5. Direct or indirect fragmentation and isolation of an area of the TEC | The development will not result in an increase in the direct or indirect fragmentation or isolation of any areas of BGHF. The impacts occur on an already isolated fragment patch of BGHF. Furthermore, the works will involve pruning of vegetation mapped as BGHF rather than removal of vegetation. |
| 6. The measures proposed to contribute to the recovery of the TEC in the IBRA subregion. | In its current form, the proposed development does not contribute to the recovery of this TEC in the IBRA subregion. |

Table 29: Evaluation of an impact on a TEC Sydney Turpentine Ironbark Forest

| Impact Assessment Provisions | Assessment |
|---|---|
| 1. the area and condition of the TEC to be impacted directly and indirectly by the proposed development | The development will remove 0.05 ha STIF which is in a degraded, weedy condition. The STIF impacted within the subject site is represented by a dominate layer of exotic weeds in the midstorey and ground layer which will be removed for the playing fields. |
| 2. the extent and overall condition of the TEC within an area of 1500 metres, and then 5000 metres, surrounding the proposed development footprint. In the case of strategic biodiversity certification projects, the extent and overall condition of the TEC may be assessed across the IBRA sub region | Within the subject site, 4.07 ha of STIF will be retained. In addition to what has been mapped within the subject site, there is an estimated 75.87 ha of STIF within an area of 1,500m, in varying condition (from large tracts to small patches) (mapped by SMCMA, 2016). The removal of 0.05 ha of STIF within the subject site represents 0.07 % of the mapped STIF extent within the 1,500 m radius. Within 5,000 m radius of the subject site, there is an estimated 361.71 ha of STIF that has been mapped with low disturbance condition. The removal of 0.05 ha of STIF from within the subject site, represents 0.01 % of the mapped STIF extent within the 5,000 m radius. |
| 3. An estimate of the extant area and overall condition of the TEC remaining before and after the impact of the proposed development has been taken into consideration | The development will not result in the overall decline of the condition of STIF after development. |
| 4. the development proposal's impact on: | |
| a. abiotic factors critical to the long-term survival of the TEC; for example, will the impact lead to a reduction of groundwater levels or substantial alteration of surface water patterns; will it alter natural disturbance regimes that the TEC depends upon, e.g. fire, flooding etc.? | The development proposal will not impact abiotic factors critical to the long-term survival of the TEC as the proposal will only result in the removal of exotic weeds and will not impact upon native vegetation of the installation of the sporting field. |
| b. characteristic and functionally important species through impacts such as, but not limited to, inappropriate fire/flooding regimes, removal of under-storey species or harvesting of plants | The development will not impact characteristic and functionally important species outside of the proposed impact area. |
| c. the quality and integrity of an occurrence of the TEC through threats and indirect impacts including, but not limited to, assisting invasive flora and fauna species to become established or causing regular mobilisation of | The development proposal is located within an area of significant weed infestation which will be removed during the proposed works. The proposed development works have potential to remove or trample regeneration of native |

| Impact Assessment Provisions | Assessment |
|---|---|
| fertilisers, herbicides or other chemicals or pollutants which may harm or inhibit growth of species in the TEC | vegetation following weed removal. The proposed development works have potential to result in the introduction of new weed plumes into the subject site. These potential impacts will be controlled during the construction phase and long-term maintenance of the subject site. These works will retain the quality and integrity of the isolated patch of STIF. |
| 5. direct or indirect fragmentation and isolation of an area of the TEC | The development will not result in an increase in the direct or indirect fragmentation or isolation of any areas of STIF. The impacts occur within a large patch of STIF which is contiguous with other patches of STIF. The proposed development will not result in isolation or fragment patch of STIF within the subject site. |
| 6. the measures proposed to contribute to the recovery of the TEC in the IBRA subregion. | In its current form, the proposed development does not contribute to the recovery of this TEC in the IBRA subregion. |

3.3 Risk assessment

A risk assessment has been undertaken for any residual impacts likely to remain after the mitigation measures (Section Table 25) have been applied. Likelihood criteria, consequence criteria and the risk matrix are provided in Table 30, Table 31 and Table 32 respectively. The risk assessment is provided in Table 33.

Table 30: Likelihood criteria

| Likelihood criteria | Description |
|--|---|
| Almost certain (Common) | Will occur, or is of a continuous nature, or the likelihood is unknown. There is likely to be an event at least once a year or greater (up to ten times per year). It often occurs in similar environments. The event is expected to occur in most circumstances. |
| Likely (Has occurred in recent history) | There is likely to be an event on average everyone to five years. Likely to have been a similar incident occurring in similar environments. The event will probably occur in most circumstances. |
| Possible (Could happen, has occurred in the past, but not common) | The event could occur. There is likely to be an event on average every five to twenty years. |
| Unlikely (Not likely or uncommon) | The event could occur but is not expected. A rare occurrence (once per one hundred years). |
| Remote (Rare or practically impossible) | The event may occur only in exceptional circumstances. Very rare occurrence (once per one thousand years). Unlikely that it has occurred elsewhere; and, if it has occurred, it is regarded as unique. |

Table 31: Consequence criteria

| Consequence category | Description |
|---|---|
| Critical (Severe, widespread long-term effect) | Destruction of sensitive environmental features. Severe impact on ecosystem. Impacts are irreversible and/or widespread. Regulatory and high-level government intervention/action. Community outrage expected. Prosecution likely. |
| Major (Wider spread, moderate to long term effect) | Long-term impact of regional significance on sensitive environmental features (e.g. wetlands). Likely to result in regulatory intervention/action. Environmental harm either temporary or permanent, requiring immediate attention. Community outrage possible. Prosecution possible. |
| Moderate (Localised, short-term to moderate effect) | Short term impact on sensitive environmental features. Triggers regulatory investigation. Significant changes that may be rehabilitated with difficulty. Repeated public concern. |
| Minor (Localised short-term effect) | Impact on fauna, flora and/or habitat but no negative effects on ecosystem. Easily rehabilitated. Requires immediate regulator notification. |
| Negligible (Minimal impact or no lasting effect) | Negligible impact on fauna/flora, habitat, aquatic ecosystem or water resources. Impacts are local, temporary and reversible. Incident reporting according to routine protocols. |

Table 32: Risk matrix

| Consequence | Likelihood | | | | |
|-------------------|----------------|-----------|----------|----------|----------|
| | Almost certain | Likely | Possible | Unlikely | Remote |
| Critical | Very High | Very High | High | High | Medium |
| Major | Very High | High | High | Medium | Medium |
| Moderate | High | Medium | Medium | Medium | Low |
| Minor | Medium | Medium | Low | Low | Very Low |
| Negligible | Medium | Low | Low | Very Low | Very Low |

Table 33: Risk assessment

| Potential impact | Project phase | Risk (pre-mitigation) | Risk (post mitigation) |
|--|--------------------------|-----------------------|------------------------|
| Vegetation clearing | Construction / operation | Medium | Low |
| sedimentation and contaminated and/or nutrient rich run-off | Construction | Medium | Low |
| noise, dust or light spill | Construction | Low | Very Low |
| inadvertent impacts on adjacent habitat or vegetation | Construction | Medium | Low |
| transport of weeds and pathogens from the site to adjacent vegetation | Construction | Medium | Low |

| Potential impact | Project phase | Risk (pre-mitigation) | Risk (post mitigation) |
|---|-----------------------------|-----------------------|------------------------|
| vehicle strike | Construction / operation | Low | Very Low |
| trampling of threatened flora species | Construction / operation | Low | Very Low |
| rubbish dumping | Construction / operation | Low | Very Low |
| wood collection | Construction / operation | Low | Very Low |
| bush rock removal and disturbance | Construction / operation | Medium | Low |
| increase in predatory species populations | Construction / operation | Low | Very Low |
| increase in pest animal populations | Construction / operation | Low | Very low |
| increased risk of fire | Construction / operation | Medium | Low |
| disturbance to specialist breeding and foraging habitat, e.g. beach nesting for shorebirds. | Construction / operation | Medium | Low |
| sedimentation and contaminated and/or nutrient rich run-off | Construction | Low | Very Low |

3.4 Adaptive management strategy

This section is required for those impacts that are infrequent, cumulative or difficult to predict. Impacts associated with the proposed development have been considered and addressed **Section 3.2.5** and no further impacts are required to be addressed.



Figure 7: Final project footprint including construction and operation

3.5 Impact summary

Following implementation of the BAM and the BAMC, the following impacts have been determined.

3.5.1 Serious and Irreversible Impacts (SAIL)

The development has candidate Serious and Irreversible Impacts (SAIL) values as outlined in Table 26 and shown on Figure 8. Detailed consideration of whether impacts on candidate species are serious and irreversible is included in Table 27 and on TECs are included in Table 28 and Table 29.

The development includes two candidates for Serious and Irreversible Impacts (SAIL), BGHF and STIF. The threshold for both of these SAILs has yet to be published by OEH, it cannot be determined with certainty if the proposed development will have a SAIL. Only a small amount of (0.001 ha) of BGHF will be removed, this equates to the removal of the outer branches of a remnant *Eucalyptus pilularis* present within the subject site. Additionally, 0.05 ha of STIF in weedy low condition will be impacted.

Table 34: Serious and Irreversible Impacts Summary

| Species Community | Common Name | Principle | Direct impact individuals / area (ha) | Summary |
|--|-----------------------------------|---------------------|---------------------------------------|--|
| Blue Gum High (BGHF) | Blue Gum High Forest | Principals 1, 2 & 4 | 0.001 | The thresholds for BGHF have not been published by OEH. The proposed development is unlikely to result in a SAIL on BGHF |
| Sydney Turpentine-Ironbark Forest (STIF) | Sydney Turpentine-Ironbark Forest | Principals 1, 2 & 4 | 0.05 | The thresholds for STIF have not been published by OEH. The impacts of the proposed development are unlikely to result in a SAIL on STIF |

3.5.2 Impacts requiring offsets

The impacts of the development requiring offset for native vegetation are outlined in Table 35 and shown on Figure 9. .

Table 35: Impacts to native vegetation that require offsets

| Veg Zone | PCT ID | PCT Name | TEC | Vegetation Class | Vegetation Formation | Direct impact (ha) |
|----------|--------|-----------------------------------|-----------|---|---|--------------------|
| 2 | 1237 | Blue Gum High Forest | Not a TEC | North Coast Wet Sclerophyll Forests | Wet Sclerophyll Forests (Shrubby sub-formation) | 0.35 |
| 4 | 1281 | Sydney Turpentine-Ironbark Forest | TEC | Northern Hinterland Wet Sclerophyll Forests | Wet Sclerophyll Forests (Grassy sub-formation) | 0.05 |

3.5.3 Areas not requiring assessment

Areas not requiring assessment include existing buildings, recreational areas (sporting fields), carparks, paths and exotic vegetation. The subject site contained build/cleared area (3.63 ha) or exotic vegetation (0.31 ha) including areas classified as 'Urban Exotics' and shown in Figure 4. These areas were not

consistent with any listed PCT, nor did they contain any threatened species, hence further assessment under the BAM was not required. Areas not requiring assessment are shown on Figure 10.

3.5.4 Credit summary

The number of ecosystem credits required for the development are outlined in Table 36. A total of seven (7) ecosystem credits are required for impacts to PCT 1237 and one (1) ecosystem credits for PCT 1281. No candidate species credit species or likely habitat was recorded within the subject site; hence no species credits are required to offset the development. A biodiversity credit report is included in Appendix G:

Table 36: Ecosystem credits required

| PCT ID | PCT Name | Credit class | Vegetation Formation | Direct impact (ha) | Credits required |
|--------|-----------------------------------|---|---|--------------------|------------------|
| 1237 | Blue Gum High Forest | North Coast Wet Sclerophyll Forests \geq 90% cleared group (including Tier 1 or higher threat status) | Wet Sclerophyll Forests (Shrubby sub-formation) | 0.35 | 7 |
| 1281 | Sydney Turpentine-Ironbark Forest | Not a TEC Like for like PCTs 1183, 1281, 1284 in Cumberland, Burragorang, Pittwater, Sydney Cataract, Wollemi or Yengo IBRA sub-region | Wet Sclerophyll Forests (Grassy sub-formation) | 0.05 | 1 |

3.6 Consistency with legislation and policy

Additional matters relating to impacts on flora and fauna which are not covered by the BC Act must also be addressed for the proposed development. Potential “Matters of National Environmental Significance” (MNES) in accordance with the EPBC Act have been addressed in Section 3.6.1. Matters relating to Hornsby Council planning instruments have been addressed in Section 3.6.2.

3.6.1 Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)

The EPBC Act establishes a process for assessing the environmental impact of activities and developments where “Matters of National Environmental Significance” (MNES) may be affected. Under the Act, any action which “has, will have, or is likely to have a significant impact on a matter of MNES” is defined as a “controlled action”, and requires approval from the Commonwealth Department of the Environment (DotE), which is responsible for administering the EPBC Act (DotE 2014).

The process includes conducting an Assessment of Significance for listed threatened species and ecological communities that represent a matter of MNES that will be impacted as a result of the proposed action. Significant impact guidelines (DotE 2014) that outline a number of criteria have been developed by the Commonwealth, to provide assistance in conducting the Assessment of Significance and help decide whether or not a referral to the Commonwealth is required.

A habitat assessment and Likelihood of Occurrence was completed and three MNES (*Pteropus poliocephalus* (Grey-headed Flying-fox), *Eucalyptus scoparia* and *Syzygium paniculatum*) was assessed under the act (Table 37).

Pteropus poliocephalus (Grey-headed Flying-fox)

The Grey-headed Flying-fox (GHFF) is listed as a Vulnerable species under the EPBC Act.

This species utilises a wide variety of habitats (including disturbed areas) for foraging and have been recorded travelling long distances on feeding forays. Fruits and flowering plants of a wide variety of species are the main food source. The species roosts in large 'camps' of up to 200 000 individuals. Camps are usually formed close to water and along gullies, however, the species has been known to form camps in urban areas (DECCW 2009).

The Gordon Grey-headed Flying-fox camp is known from the locality to be within 8 km of the subject site (OEH 2017b). The vegetation within the subject site provides potential foraging habitat. It is considered likely that this species would use the site on occasion for foraging purposes. According to the National Flying-fox Monitoring Program, no GHFF camps currently occur or have been recorded within the subject site (DotE 2018).

Table 37: EPBC Act of Significance for *Pteropus poliocephalus* (Grey-headed Flying-fox)

| Criterion | Assessment |
|--|--|
| Criterion a: lead to a long-term decrease in the size of an important population of a species | <p>The Matters of National Environmental Significance Impact Guidelines 1.1 (Commonwealth of Australia, 2013) defines an important population as a population that is necessary for a species' long-term survival and recovery. This may include populations identified as such in recovery plans, and/or that are:</p> <ul style="list-style-type: none"> • Key source populations either for breeding or dispersal • Populations that are necessary for maintaining genetic diversity, and/or • Populations that are near the limit of the species range <p>No important populations have been recorded within the subject site. The site does not support key source populations for breeding or dispersal, populations necessary for maintaining genetic diversity, or populations near the limit of the species range. According to the National Flying-fox Monitoring Program, no GHFF camps currently occur or have ever been recorded within the subject site (DotE 2018). The nearest active GHFF camp occurs approximately 8 km to the south-east of the subject site, within Gordon (DotE 2018).</p> |
| Criterion b: reduce the area of occupancy of an important population | No important populations have been recorded within the subject site. Therefore, the proposed works would not reduce the area of occupancy of an important population. |
| Criterion c: fragment an existing important population into two or more populations | No important populations have been recorded within the subject site. The potential foraging habitat to be removed is marginal relative to adjacent potential habitat within the region. Whilst the potential foraging habitat may contribute as a 'stepping stone' for this highly mobile species to other more substantial foraging habitat sites, this function is unlikely to be significantly inhibited by the proposed works. Furthermore, this species has been recorded in urban environments and is likely to continue to forage adjacent to the site and across the broader locality. |
| Criterion d: adversely affect habitat critical to the survival of a species | <p>Less than half of the potential foraging habitat in canopy trees within the subject site will be removed by the proposal.</p> <p>These individual trees represent a negligible amount of potential foraging resources in the locality. Potential foraging habitat will persist in close proximity to the subject site, within the remaining subject site and in large stands of high quality intact native vegetation in Coups Creek riparian corridor adjacent to the Subject site and in the Lane Cove River National Park (approximately 2 km NE from the Subject site). Given that this</p> |

| Criterion | Assessment |
|---|---|
| | species is highly mobile (traveling up to 50 km to forage), it is considered unlikely that the works would adversely affect habitat critical to the survival of this species |
| e: disrupt the breeding cycle of an important population | According to the National Flying-fox Monitoring Program, no GHFF camps currently occur or have ever been recorded within the subject site (DotE 2018). The nearest active GHFF camp occurs approximately 8 km to the north-east of the subject site, within Gordon (DotE 2018). Thus, no important population of GHFF occurs within the subject site, and the proposed works is unlikely to disrupt the breeding cycle of an important population. |
| Criterion f: Adversely affect habitat critical to the survival of a species; modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline | The potential foraging habitat to be removed is marginal and of low quality. Given the small amount of potential foraging habitat to be removed, that potential foraging habitat will persist adjacent to the subject site and across the locality, and that this species is highly mobile, it is unlikely that the habitat to be removed would cause the species to decline. Furthermore, according to the National Flying-fox Monitoring Program, no GHFF camps currently occur or have ever been recorded within the subject site (DotE 2018). The nearest active GHFF camp occurs approximately 8 km to the north-east of the Subject site, within Gordon (DotE 2018). Therefore, no known GHFF roosting camps for this species will be impacted by the proposed works. |
| Criterion g: Result in invasive species that are harmful to a vulnerable species becoming established in the vulnerable species' habitat | The proposed works will not result in the establishment of an invasive species that is harmful to GHFF. |
| Criterion h: Introduce disease that may cause the species to decline | The proposed works will not result in the introduction of a disease that is harmful to the GHFF. |
| Criterion i: Interfere substantially with the recovery of the species | Considering the above factors, the proposed works will not interfere substantially with the recovery of the species. |
| Conclusion | In consideration of the above, the proposed works are considered unlikely to have a significant impact on the GHFF. |

Eucalyptus scoparia (Wallangarra White Gum)

Eucalyptus scoparia (Wallangarra White Gum) is listed as endangered under the BC Act and vulnerable under the EPBC Act. This species has been planted as landscaping trees and was not recorded within remnant or part of a native vegetation patch. This threatened species is known from only three locations in NSW near Tenterfield, which is more than 640 km from the development site. The development site is not connected to the known geographic distribution of this species. *Eucalyptus scoparia* occurs in open eucalypt forests and heath, typically at high altitudes. The development site does not represent suitable habitat for this species. Five specimens were recorded within the development site (Tree 9, 10, 310, 390 and 424) by arborist (Earthscapes 2020).

Table 38: EPBC Act of Significance for *Eucalyptus scoparia*

| Criterion | Assessment |
|---|---|
| Criterion a: lead to a long-term decrease in the size of an important population of a species | Five <i>Eucalyptus scoparia</i> were recorded within the subject site from the arborist report and will be retained within the subject site. This species has a highly restricted distribution which does not include the subject site. The <i>Eucalyptus scoparia</i> species profile has recognised that this species also occurs as a planted specimen. It is unlikely the specimens recorded within the subject site represent an important population as they are not indigenous to the local area, is not associated with PCT 1231 or 1281 and are unlikely to have germinated from any soil seedbank. As the development does not occur within the geographic distribution for this species the works will not impact upon an important population of this species. As the proposed development will not directly impact upon this species the proposed works are unlikely to result in a long-term decrease in the size of an important population. |
| Criterion b: reduce the area of occupancy of an important population | The subject site contains five scattered <i>Eucalyptus scoparia</i> which will be retained within the subject site. The subject site is not located within their normal range of distribution. As the subject site is located 640 km from potential habitat for this species, it is unlikely that these trees occur naturally. The subject site is not located within an area which would represent an important population for this species. As such the proposed works will not reduce the area of occupancy of an important population for <i>Eucalyptus scoparia</i> . |
| Criterion c: fragment an existing important population into two or more populations | The development will not result in the fragmentation of an existing important population into two or more population as the subject site is not located within an area which this species occurs naturally. The five <i>Eucalyptus scoparia</i> identified within the subject site are likely to be planted specimens with unknown origin of genetic material. These specimens are not considered part of an important population as they are located outside of their natural geographic distribution, are of unknown seed stock and are not considered native to the PCT. The works will not result in fragmentation of important populations for this species. |
| Criterion d: adversely affect habitat critical to the survival of a species | Habitat critical to the survival of this species is likely to include grassy woodlands on infertile soils on slopes and ridges. The subject site is located more than 640 km from this species geographic distribution and does not contain suitable habitat or potential soil seed bank for this species. The development does not contain habitat critical to the survival of this species. |
| Criterion e: disrupt the breeding cycle of an important population | As the development will not remove potential habitat for this species, it is unlikely that the works would disrupt the breeding cycle (i.e. soil seed bank) of this species in an important population. |
| Criterion f: Adversely affect habitat critical to the survival of a species; modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline | No habitat critical to the survival of <i>Eucalyptus scoparia</i> would be removed, or disturbed, under the proposed works. The proposed action would therefore be unlikely to modify, destroy, remove, or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline. |
| Criterion g: Result in invasive species that are harmful to a vulnerable species becoming established in the vulnerable species' habitat | The proposal would not result in invasive species, such as weeds, that would be harmful. It is unlikely that the proposed action will result in a large increase in the number of weeds due to the current disturbed nature of the site. |
| Criterion h: Introduce disease that may cause the species to decline | The proposed development is unlikely to introduce disease that may cause the species to decline. |

| Criterion | Assessment |
|--|--|
| Criterion i: Interfere substantially with the recovery of the species | As recovery of this species aims to protect suitable habitat for this species which is restricted to its specific geographic distribution. As the proposed works are not located within the geographic distribution for this species, the proposed works are unlikely to interfere substantially with the recovery of this species. |
| Conclusion | The five <i>Eucalyptus scoparia</i> specimens located within the subject site are planted stock from an unknown origin of seed. The subject site is located more than 640 km from the known distribution of this species. The proposed works will not result in removal of a known or potential important population. No important populations would be isolated or fragmented and the life cycle (seed bank) of this species is not likely to be affected. Therefore, the action is not likely to have a significant impact on this species and a Referral is not required. |

Syzygium paniculatum (Magenta Lilly Pilly)

This species is listed as vulnerable under the EPBC Act. It occurs from Upper Landsowne to Conjola State Forest. It is restricted mainly to remnant stands of littoral (coastal) rainforest.

Table 39: EPBC Act of Significance for *Syzygium paniculatum*

| Criterion | Assessment |
|--|---|
| Criterion a: lead to a long-term decrease in the size of an important population of a species | An important population is defined as a population that is necessary for a species' long-term survival and recovery. Seven small <i>Syzygium paniculatum</i> specimens were recorded within the subject site located within landscaped garden which is disconnected from suitable habitat for this species. No <i>Syzygium paniculatum</i> will be removed for the proposed development. Consequently, it is considered that the proposed development will not lead to a long-term decrease in the size of an important population of the species. |
| Criterion b: reduce the area of occupancy of an important population | This species typically grows in littoral coastal rainforests of NSW; however, it has been identified as a species in association with PCT 1281. The <i>Syzygium paniculatum</i> within the subject site is not located within littoral rainforest or within PCT 1281. The specimens are located within a highly landscaped environment. The specimens are immature and located within a linear garden. There is potential that this species has been planted outside of its natural area of distribution. Therefore, it is unlikely to form part of an important population. Consequently, it is considered that the proposed development will not reduce the area of occupancy of an important population of this species. |
| Criterion c: fragment an existing important population into two or more populations | The development site contains seven immature <i>Syzygium paniculatum</i> specimens, of which none will be removed for the proposed redevelopment. The specimens within the development site are unlikely to be considered an important or viable population given that these specimens may have been planted and do not form part of an important population (i.e. they are not located in their natural habitat). Consequently, it is considered that the proposed development will not fragment an existing important population. |
| Criterion d: adversely affect habitat critical to the survival of a species | The <i>Syzygium paniculatum</i> were recorded within a landscaped garden and is therefore it is not considered to be important or critical to the survival of the species. Consequently, it is considered that the proposed development will not adversely affect habitat critical to the survival of this species. |
| Criterion e: disrupt the breeding cycle of an important population | Not applicable. |

| Criterion | Assessment |
|---|--|
| Criterion f: Adversely affect habitat critical to the survival of a species; modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline | No <i>Syzygium paniculatum</i> are proposed to be removed. This species was identified outside its natural littoral coastal rainforest habitat in a highly landscaped urbanised environment. It is considered unlikely that the development site will modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline. |
| Criterion g: Result in invasive species that are harmful to a vulnerable species becoming established in the vulnerable species' habitat | The site is currently in a disturbed and modified condition and does not represent known habitat for this threatened species. Consequently, the proposed development is unlikely to result in the establishment of an invasive species that is harmful to this species. |
| Criterion h: Introduce disease that may cause the species to decline | It is considered unlikely that the proposed action would introduce disease that may cause the decline of <i>Syzygium paniculatum</i> . |
| Criterion i: Interfere substantially with the recovery of the species | <p>The 2012 <i>Syzygium paniculatum</i> Recovery Plan aims to</p> <ul style="list-style-type: none"> ensuring a coordinated and efficient approach to the implementation of recovery efforts establishing the full extent of the distribution of Magenta Lilly Pilly increasing the understanding of Magenta Lilly Pilly biology and ecology minimising the decline of Magenta Lilly Pilly through in situ habitat protection and management reducing impacts of Myrtle Rust on Magenta Lilly Pilly and its habitat maintaining a representative ex situ collection of Magenta Lilly Pilly raising awareness of the conservation significance of Magenta Lilly Pilly and involving the broader community in the recovery program <p>The proposed would not interfere substantially with the recovery of this species.</p> |
| Conclusion | <p>No. The proposed action is unlikely to have a significant impact on the <i>Syzygium paniculatum</i> for the following reasons:</p> <ul style="list-style-type: none"> No individuals are to be removed The specimens are likely to have been planted and do not form part of an important population The development site is located outside of its known habitat (littoral coastal rainforests or mapped PCT 1281 areas). |

3.6.2 Hornsby Council Local Environmental Plan 2013 (HLEP)

The HLEP has identified that the subject site contains land identified as 'Biodiversity' as mapped by the HLEP Terrestrial Biodiversity Map (Figure 2).

Clause 6.4 of the HLEP provides provisions for the protection of lands identified as 'Terrestrial Biodiversity'. Development within this area must avoid any significant adverse environmental impact or minimise or mitigate the impact on the biodiversity. This section describes the consistency of the proposed development with clause 6.4. Clause 6.4 (3a) of the HLEP states that Council must consider any potential adverse impacts from the proposed development on the matters listed in Table 40.

Table 40: Clause 6.4 of the HLEP

| Condition | Response | Environmental impact |
|---|--|----------------------|
| a) whether the development is likely to have: | | |
| (i) any adverse impacts on the condition, ecological value and significance of the fauna and flora on the land; and | The proposal will impact on approximately 0.05 ha of the land mapped as 'Biodiversity' which has been identified as a TEC (STIF) in weedy condition. The proposal will not result in a significant impact on any threatened flora or fauna species listed under environmental legislation. Although 0.05 ha of land mapped as 'Biodiversity' will be impacted, an additional 4.12 ha of vegetation will be retained within the subject site. The vast majority (>99%) of the area mapped as 'Biodiversity' will be retained within the subject site. The majority of the STIF vegetation to be retained within the Biodiversity layer is in better condition than that proposed to be impacted. Additionally, the vegetation removed comprise only weeds, with existing native species retained. Therefore, it is expected the proposal will impact positively on the TEC due to the removal of weeds. | Positive |
| (ii) any adverse impact on the importance of the vegetation on the land to the habitat and survival of native fauna; and | The vegetation within the subject site is connected to a large tract of vegetation to the south which eventually extends into Lane Cove National Park. The 0.05 ha of vegetation mapped as 'Biodiversity' is currently highly infested with weeds and has marginal potential to be habitat for threatened species. The proposal will not impact on the vegetation that is likely to be significant for threatened flora and fauna species. The 0.05 ha of land mapped as 'Biodiversity' does not contain any hollow bearing trees and it is unlikely to significantly impact habitat for threatened species. The proposal will improve the condition of the existing vegetation, therefore having a positive impact on threatened flora and fauna within the surrounding area. | Positive |
| (iii) any potential to fragment, disturb or diminish the biodiversity structure, function and composition of the land; and | The vegetation within the subject site is connected to a large tract of vegetation to the south which eventually extends into Lane Cove National Park. The proposed works will not result in a loss of biodiversity structure or function or composition of the land such that the vegetation condition is reduced or species assemblage is diminished. Furthermore, the proposal will not fragment the vegetation. | Positive |
| (iv) any adverse impacts on the habitat elements providing connectivity on the land; and | The vegetation within the subject site and subject site is well connected to a much larger vegetation patch within the surrounding landscape which is also connected to a number of national parks and the vegetation associated with Berowra Valley Regional Park and Lane Cove National Park. The proposal will not fragment the vegetation and will not significantly impact on the vegetation's ability to act as a habitat corridor. | Neutral |
| (b) any proposed measures to avoid, minimise or mitigate the impacts of the development. | The proposed development has utilised existing development footprint, cleared areas, planted vegetation and low condition TEC and retained areas of higher quality TEC. | Neutral |

3.7 Offset options

There are a number of options that can be utilised to offset the required ecosystem credits. These include retiring matching biodiversity credits either through establishing a Biodiversity Stewardship Agreement (offset) on land owned by Loreto Normanhurst (i.e. to the onsite bushland), through purchasing matching credits on the open market, making a payment to the Biodiversity Conservation Trust, or funding biodiversity actions for individual species or communities. However, this last option has some limitations. Due to the small scale of the project, it is likely that making a payment to the Biodiversity Conservation Trust will be the preferred option to retire credits for this redevelopment.



Figure 8: Serious and Irreversible Impacts



Figure 9: Impacts requiring offset



Figure 10: Areas not requiring assessment



Plate 1: Impacts not requiring assessment

4. References

- Allen Jack and Cottier (AJ+C) 2020. Revised SSD application – concept and envelopes – revisions responding to DPE comments and questions
- Chapman, G.A and Murphy, C.L. 1989. *Soil Landscapes of the Sydney 1:100 000 sheet*. Soil Conservation Service of NSW, Sydney.
- Churchill, S. 2009. *Australian Bats*. Allen & Unwin. 2nd Edition.
- Department of the Environment and Energy (DotEE) 2018a. *Blue Gum High Forest of the Sydney Basin Bioregion*. Australian Government. Accessed online: <http://environment.gov.au/biodiversity/threatened/conservation-advice/blue-gum-high-forest-sydney-region> (Accessed 16 November 2018).
- Department of the Environment and Energy (DotEE) 2018b. *Turpentine-Ironbark Forest of the Sydney Basin Bioregion*. Australian Government. Accessed online: <http://www.environment.gov.au/biodiversity/threatened/conservation-advice/turpentine-ironbark-forest-sydney-basin-bioregion> (Accessed 16 November 2018).
- Department of Environment and Climate Change. 2002, 'Descriptions for NSW (Mitchell) Landscapes Version 2'. Sourced 3 October 2018 from: <http://www.environment.nsw.gov.au/resources/conservation/landscapesdescriptions.pdf>
- Department of Environment, Climate Change and Water NSW (DECCW) 2009. Draft National Recovery Plan for the Grey-headed Flying-fox *Pteropus poliocephalus*. Prepared by Dr Peggy Eby. Department of Environment, Climate Change and Water NSW, Sydney.
- Department of the Environment and Energy (DoEE) 2018c. National Flying-fox monitoring viewer. Australian Government. Available: <http://www.environment.gov.au/webgis-framework/apps/ffc-wide/ffc-wide.jsf> (Accessed: 3 October 2018)
- Department of the Environment and Energy (DoEE) 2018d. Protected Matters Search Tool [online]. Available: <http://www.environment.gov.au/epbc/protect/index.html> (Accessed: October 2018).
- Department of the Environment and Energy (DoEE) 2018e. Species Profile and Threats Database. Available <http://www.environment.gov.au/cgi-bin/sprat/public/sprat.pl>.
- Earthscape Horticultural services. 2020. Arboricultural impact assessment report – concept masterplan and detailed stage 1 works Loreto Normanhurst 91-93 Pennant Hills Road Normanhurst
- Hornsby Shire Council 2010. Native vegetation communities map - Pennant Hills, ENV Smith and Smith Vegetation. [online] <http://www.hornsby.nsw.gov.au/media/documents/environment-and-waste/bushland-and-biodiversity/native-vegetation-communities/Native-Vegetation-Communities-Map-Pennant-Hills.pdf> [Accessed November 2018].

Office of Environment and Heritage 2016. NSW Guide of Surveying Threatened Plants. Available: <http://www.environment.nsw.gov.au/resources/threatenedspecies/160129-threatened-plants-survey-guide.pdf>

Office of Environment and Heritage 2016. The Native Vegetation of the Sydney Metropolitan Area. Volume 2: Vegetation Community Profiles. Version 3.0. NSW Office of Environment and Heritage, Sydney.

Office of Environment and Heritage (OEH). 2018a. Threatened Species Database (5 km radius search). OEH Sydney, NSW. (Data viewed November 2018).

Office of Environment and Heritage (OEH) 2018b. Threatened Species Profiles. Available: <http://www.environment.nsw.gov.au/threatenedspeciesapp/profile.aspx?>

Appendix A: Definitions

| Terminology | Definition |
|--------------------------------|---|
| Biodiversity credit report | The report produced by the Credit Calculator that sets out the number and class of biodiversity credits required to offset the remaining adverse impacts on biodiversity values at a subject site, or on land to be biodiversity certified, or that sets out the number and class of biodiversity credits that are created at a biodiversity stewardship site. |
| BioNet Atlas | The BioNet Atlas (formerly known as the NSW Wildlife Atlas) is the OEH database of flora and fauna records. The Atlas contains records of plants, mammals, birds, reptiles, amphibians, some fungi, some invertebrates (such as insects and snails) and some fish |
| Broad condition state: | Areas of the same PCT that are in relatively homogenous condition. Broad condition is used for stratifying areas of the same PCT into a vegetation zone for the purpose of determining the vegetation integrity score. |
| Connectivity | The measure of the degree to which an area(s) of native vegetation is linked with other areas of vegetation. |
| Credit Calculator | The computer program that provides decision support to assessors and proponents by applying the BAM, and which calculates the number and class of biodiversity credits required to offset the impacts of a development or created at a biodiversity stewardship site. |
| Development | Has the same meaning as development at section 4 of the EP&A Act, or an activity in Part 5 of the EP&A Act. It also includes development as defined in section 115T of the EP&A Act. |
| Development footprint | The area of land that is directly impacted on by a proposed development, including access roads, and areas used to store construction materials. |
| Subject site | An area of land that is subject to a proposed development that is under the EP&A Act. |
| Ecosystem credits | A measurement of the value of EECs, CEECs and threatened species habitat for species that can be reliably predicted to occur with a PCT. Ecosystem credits measure the loss in biodiversity values at a subject site and the gain in biodiversity values at a biodiversity stewardship site. |
| High threat exotic plant cover | Plant cover composed of vascular plants not native to Australia that if not controlled will invade and outcompete native plant species. |
| Hollow bearing tree | A living or dead tree that has at least one hollow. A tree is considered to contain a hollow if: (a) the entrance can be seen; (b) the minimum entrance width is at least 5 cm; (c) the hollow appears to have depth (i.e. you cannot see solid wood beyond the entrance); (d) the hollow is at least 1 m above the ground. Trees must be examined from all angles. |
| Important wetland | A wetland that is listed in the Directory of Important Wetlands of Australia (DIWA) and SEPP 14 Coastal Wetlands |
| Linear shaped development | Development that is generally narrow in width and extends across the landscape for a distance greater than 3.5 kilometres in length |
| Local population | The population that occurs in the study area. In cases where multiple populations occur in the study area or a population occupies part of the study area, impacts on each subpopulation must be assessed separately. |
| Local wetland | Any wetland that is not identified as an important wetland (refer to definition of Important wetland). |
| Mitchell landscape | Landscapes with relatively homogeneous geomorphology, soils and broad vegetation types, mapped at a scale of 1:250,000. |

| Terminology | Definition |
|---|--|
| Multiple fragmentation impact development | Developments such as wind farms and coal seam gas extraction that require multiple extraction points (wells) or turbines and a network of associated development including roads, tracks, gathering systems/flow lines, transmission lines |
| Operational Manual | The Operational Manual published from time to time by OEH, which is a guide to assist assessors when using the BAM |
| Patch size | An area of intact native vegetation that: a) occurs on the subject site or biodiversity stewardship site, and b) includes native vegetation that has a gap of less than 100 m from the next area of native vegetation (or ≤ 30 m for non-woody ecosystems). Patch size may extend onto adjoining land that is not part of the subject site or stewardship site. |
| Proponent | A person who intends to apply for consent to carry out development or for approval for an activity. |
| Reference sites | The relatively unmodified sites that are assessed to obtain local benchmark information when benchmarks in the Vegetation Benchmarks Database are too broad or otherwise incorrect for the PCT and/or local situation. Benchmarks can also be obtained from published sources. |
| Regeneration | The proportion of over-storey species characteristic of the PCT that are naturally regenerating and have a diameter at breast height < 5 cm within a vegetation zone. |
| Remaining impact | An impact on biodiversity values after all reasonable measures have been taken to avoid and minimise the impacts of development. Under the BAM, an offset requirement is calculated for the remaining impacts on biodiversity values. |
| Retirement of credits | The purchase and retirement of biodiversity credits from an already-established biobank site or a biodiversity stewardship site secured by a biodiversity stewardship agreement. |
| Riparian buffer | Riparian buffers applied to water bodies in accordance with the BAM |
| Sensitive biodiversity values land map | Development within an area identified on the map requires assessment using the BAM. |
| Site attributes | The matters assessed to determine vegetation integrity. They include: native plant species richness, native over-storey cover, native mid-storey cover, native ground cover (grasses), native ground cover (shrubs), native ground cover (other), exotic plant cover (as a percentage of total ground and mid-storey cover), number of trees with hollows, proportion of over-storey species occurring as regeneration, and total length of fallen logs. |
| Site-based development | a development other than a linear shaped development, or a multiple fragmentation impact development |
| Species credits | The class of biodiversity credits created or required for the impact on threatened species that cannot be reliably predicted to use an area of land based on habitat surrogates. Species that require species credits are listed in the Threatened Biodiversity Data Collection. |
| Subject land | Is land to which the BAM is applied in Stage 1 to assess the biodiversity values of the land. It includes land that may be a subject site, clearing site, proposed for biodiversity certification or land that is proposed for a biodiversity stewardship agreement. |
| Threatened Biodiversity Data Collection | Part of the BioNet database, published by OEH and accessible from the BioNet website. |
| Threatened species | Critically Endangered, Endangered or Vulnerable threatened species as defined by Schedule 1 of the BC Act, or any additional threatened species listed under Part 13 of the EPBC Act as Critically Endangered, Endangered or Vulnerable. |

| Terminology | Definition |
|--------------------------------|---|
| Vegetation Benchmarks Database | A database of benchmarks for vegetation classes and some PCTs. The Vegetation Benchmarks Database is published by OEH and is part of the BioNet Vegetation Classification. |
| Vegetation zone | A relatively homogenous area of native vegetation on a subject site, land to be biodiversity certified or a biodiversity stewardship site that is the same PCT and broad condition state. |
| Wetland | An area of land that is wet by surface water or ground water, or both, for long enough periods that the plants and animals in it are adapted to, and depend on, moist conditions for at least part of their life cycle. Wetlands may exhibit wet and dry phases and may be wet permanently, cyclically or intermittently with fresh, brackish or saline water |
| Woody native vegetation | Native vegetation that contains an over-storey and/or mid-storey that predominantly consists of trees and/or shrubs |

Appendix B: Vegetation plot data

Table 41: Species matrix (species recorded by plot)

| Stratum | Form | Species name | Exotic (*) | High Threat Weed* | Cover (%) Plot 1 | Cover (%) Plot 2 | Cover (%) Plot 4 | Cover (%) Plot 5** |
|---------|------------------------|--------------------------------|------------|-------------------|------------------|------------------|------------------|--------------------|
| G | Shrub (SG) | <i>Acacia falcata</i> | | 0 | 0.1 | | | |
| G | Shrub (SG) | <i>Acacia linifolia</i> | | 0 | 1 | | | |
| M | Tree (TG) | <i>Acacia parramattensis</i> | | 0 | 8 | | | |
| G | 0 | <i>Agapanthus spp.</i> | * | 0 | 20 | 20 | | |
| M | Tree (TG) | <i>Allocasuarina torulosa</i> | | 0 | 5 | | | |
| U | Tree (TG) | <i>Angophora costata</i> | | 0 | | | 5 | 10 |
| U | Tree (TG) | <i>Angophora floribunda</i> | | 0 | | | 1 | |
| G | 0 | <i>Anigozanthos sp.</i> | * | 0 | 0.1 | | | |
| G | 0 | <i>Aristea ecklonii</i> | * | 0 | | | | 0.1 |
| G | 0 | <i>Asparagus aethiopicus</i> | * | 1 | 10 | 0.1 | 0.1 | |
| G | 0 | <i>Bidens pilosa</i> | * | 1 | 0.1 | | | |
| M | Shrub (SG) | <i>Breynia oblongifolia</i> | | 0 | | | 0.1 | |
| M | Shrub (SG) | <i>Callistemon sp.</i> | | 0 | | 10 | | 5 |
| M | Shrub (SG) | <i>Callistemon sp.</i> | | 0 | 10 | | | |
| G | Forb (FG) | <i>Centella asiatica</i> | | 0 | | | 0.1 | |
| U | 0 | <i>Cinnamomum camphora</i> | * | 1 | | | 5 | |
| U | Tree (TG) | <i>Corymbia eximia</i> | | 0 | 10 | | | |
| U | Tree (TG) | <i>Corymbia maculata</i> | | 0 | 15 | | | |
| M | 0 | <i>Cotoneaster spp.</i> | * | 1 | | | 0.1 | |
| G | Grass & grasslike (GG) | <i>Cynodon dactylon</i> | | 0 | 0.1 | 0.5 | | |
| G | Forb (FG) | <i>Dianella caerulea</i> | | 0 | | | 0.1 | |
| G | Forb (FG) | <i>Dianella sp.</i> | | 0 | 0.1 | 0.1 | | |
| G | 0 | <i>Ehrharta erecta</i> | * | 1 | 5 | 0.1 | 0.1 | 11 |
| U | Shrub (SG) | <i>Elaeocarpus reticulatus</i> | | 0 | | | 1 | |
| G | Grass & grasslike (GG) | <i>Entolasia marginata</i> | | 0 | 0.1 | | 1 | |
| U | Tree (TG) | <i>Eucalyptus punctata</i> | | 0 | | | 10 | 20 |
| U | Tree (TG) | <i>Eucalyptus resinifera</i> | | 0 | | | 5 | |

| Stratum | Form | Species name | Exotic (*) | High Threat Weed* | Cover (%) Plot 1 | Cover (%) Plot 2 | Cover (%) Plot 4 | Cover (%) Plot 5** |
|---------|------------------------|------------------------------------|------------|-------------------|------------------|------------------|------------------|--------------------|
| U | Tree (TG) | <i>Eucalyptus saligna</i> | | 0 | 10 | | 15 | 15 |
| G | 0 | <i>Euphorbia cyathophora</i> | * | 0 | | | | 0.1 |
| G | Other (OG) | <i>Eustrephus latifolius</i> | | 0 | | | 0.1 | |
| U | Tree (TG) | <i>Ficus macrophylla</i> | | 0 | | 20 | | 15 |
| U | Tree (TG) | <i>Ficus rubiginosa</i> | | 0 | | 20 | | |
| M | 0 | <i>Genista monspessulana</i> | * | 1 | 0.1 | 0.1 | | |
| G | Other (OG) | <i>Glycine clandestina</i> | | 0 | | | 0.1 | |
| M | Shrub (SG) | <i>Grevillea sp.</i> | | 0 | 1 | | | |
| M | Shrub (SG) | <i>Grevillea juniperina</i> | | 0 | 11 | | | |
| M | Tree (TG) | <i>Grevillea robusta</i> | | 0 | 1 | | | |
| G | 0 | <i>Hedera helix</i> | * | 1 | 0.1 | | | |
| G | 0 | <i>Hyparrhenia hirta</i> | * | 1 | 0.1 | | | |
| M | 0 | <i>Jacaranda mimosifolia</i> | * | 0 | 1 | 15 | | |
| M | 0 | <i>Lantana camara</i> | * | 1 | | | 5 | |
| G | Grass & grasslike (GG) | <i>Lepidosperma laterale</i> | | 0 | | | 5 | |
| M | Shrub (SG) | <i>Leptospermum polygalifolium</i> | | 0 | 5 | | | |
| M | 0 | <i>Ligustrum lucidum</i> | * | 1 | | | 1 | |
| M | 0 | <i>Ligustrum sinense</i> | * | 1 | | | 30 | |
| M | 0 | <i>Lilium formosanum</i> | * | 0 | | | 0.1 | |
| M | Shrub (SG) | <i>Lissanthe strigosa</i> | | 0 | | | 5 | |
| G | Grass & grasslike (GG) | <i>Lomandra longifolia</i> | | 0 | 0.1 | 0.1 | | |
| G | 0 | <i>Lonicera japonica</i> | * | 1 | | | 2 | |
| U | Tree (TG) | <i>Lophostemon confertus</i> | | 0 | | 15 | | |
| M | Shrub (SG) | <i>Melaleuca styphelioides</i> | | 0 | 2 | | | |
| G | Other (OG) | <i>Morinda jasminoides</i> | | 0 | | | 0.1 | |
| M | 0 | <i>Morus alba</i> | * | 0 | 1 | 1 | | |
| M | 0 | <i>Ochna serrulata</i> | * | 1 | 0.5 | | 0.1 | |
| G | 0 | <i>Olea europaea</i> | * | 1 | 0.1 | | 0.1 | |
| M | 0 | <i>Olea europaea</i> | * | 1 | | | 0.1 | |
| G | Grass & grasslike (GG) | <i>Opismenus aemulus</i> | | 0 | 0.01 | | 2 | |

| Stratum | Form | Species name | Exotic (*) | High Threat Weed* | Cover (%) Plot 1 | Cover (%) Plot 2 | Cover (%) Plot 4 | Cover (%) Plot 5** |
|---------|------------|----------------------------------|------------|-------------------|------------------|------------------|------------------|--------------------|
| G | Forb (FG) | <i>Oxalis spp.</i> | | 0 | | | | 0.1 |
| G | 0 | <i>Pennisetum clandestinum</i> | * | 1 | 0.1 | | | |
| M | 0 | <i>Phoenix spp.</i> | * | 0 | 0.1 | 0.1 | | |
| M | 0 | <i>Phyllostachys aurea</i> | * | 1 | | 0.1 | | |
| G | 0 | <i>Phyllostachys aurea</i> | * | 1 | 0.1 | | | |
| M | Shrub (SG) | <i>Pittosporum undulatum</i> | | 0 | 0.1 | | 5 | |
| G | 0 | <i>Plantago lanceolata</i> | * | 0 | 0.1 | | | |
| G | Forb (FG) | <i>Pratia purpurascens</i> | | 0 | | | 0.1 | |
| G | Forb (FG) | <i>Pseuderanthemum variabile</i> | | 0 | | | 0.1 | |
| G | Fern (EG) | <i>Pteridium esculentum</i> | | 0 | | | 5 | |
| G | Shrub (SG) | <i>Rubus sp.</i> | * | 0 | | | 1 | |
| M | 0 | <i>Sida rhombifolia</i> | * | 0 | 0.1 | | | |
| G | 0 | <i>Sonchus asper</i> | * | 0 | 0.1 | | | |
| G | 0 | <i>Stenotaphrum secundatum</i> | * | 1 | | 1 | | |
| G | 0 | <i>Strelitzia spp.</i> | * | 0 | | | 0.1 | |
| U | Tree (TG) | <i>Syncarpia glomulifera</i> | | 0 | | | 20 | 10 |
| G | Forb (FG) | <i>Veronica plebeia</i> | | 0 | | 0.1 | | |

Tree (TG), Shrub (SG), Grass & Grasslike (GG), Forb (FG), Fern (EG), Other (OG)

* 0 = not a High Threat Weed. 1= High Threat Weed

** note Plot 5 was not used in the credit calculator

Table 42: Vegetation integrity data (Composition, Structure and function)

| Plot location data | | | | | | | |
|--------------------|------|-----------------|-----------|------|----------|-----------|---------|
| Plot no. | PCT | Vegetation Zone | Condition | Zone | Eastings | Northings | Bearing |
| 1 | 1237 | 2 | Degraded | 56 | 323829 | 6266693 | 140 |
| 2 | 1237 | 2 | Degraded | 56 | 323824 | 6266693 | 0 |
| 4 | 1281 | 4 | Degraded | 56 | 0323675 | 6266287 | 200 |
| 5** | 1237 | 2 | Degraded | 56 | 0323662 | 6266580 | 30 |

* Note, plot 3 was not used.

** note plot 5 was not entered into the credit calculator

Composition (number of species)

| Plot no. | Tree | Shrub | Grass | Forb | Fern | Other |
|----------|------|-------|-------|------|------|-------|
| 1 | 7 | 8 | 4 | 1 | 0 | 0 |
| 2 | 3 | 1 | 2 | 2 | 0 | 0 |
| 4 | 6 | 5 | 3 | 4 | 1 | 3 |
| 5* | 5 | 1 | 0 | 1 | 0 | 0 |

* note this plot was not used in the credit calculator

Structure (Total cover)

| Plot no. | Tree | Shrub | Grass | Forb | Fern | Other |
|----------|------|-------|-------|------|------|-------|
| 1 | 49 | 30 | <1 | <1 | 0 | 0 |
| 2 | 55 | 10 | 1 | <1 | 0 | 0 |
| 4 | 56 | 12 | 8 | <1 | 5 | 0 |
| 5 | 70 | 5 | 0 | <1 | 0 | 0 |

Function

| Plot no. | Large Trees | Hollow trees | Litter Cover | Length Fallen Logs | Tree Stem 5-9 | Tree Stem 10-19 | Tree Stem 20-29 | Tree Stem 30-49 | Tree Stem 50-79 | Tree Regen | High Threat Weed Cover |
|----------|-------------|--------------|--------------|--------------------|---------------|-----------------|-----------------|-----------------|-----------------|------------|------------------------|
| 1 | 0 | 0 | 65 | 0 | 0 | 1 | 1 | 1 | 1 | 0 | 16 |
| 2 | 2 | 1 | 60 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 1 |
| 4 | 0 | 0 | 84 | 0 | 1 | 1 | 1 | 1 | 1 | 0 | 44 |
| 5* | 0 | 0 | 80 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 11 |

Appendix C: Photos



Plate 2: Plot 1: Left start. Right end.



Plate 3: Plot 2: Left start. Right end



Plate 4: Plot 3 Left start, Right end



Plate 5: Plot 4 Left start, Right end



Appendix D: Other species recorded

| Botanic Name | Common Name | High Threat Exotics | WoNS | Exotic/ Native | Study area | Bushland |
|---------------------------------|-----------------------|---------------------|------|----------------|------------|----------|
| <i>Acacia baileyana</i> | Cootamundra wattle | | | E | x | |
| <i>Acacia decurrens</i> | Black Wattle | | | N | x | |
| <i>Acacia implexa</i> | | | | N | x | |
| <i>Acacia linifolia</i> | White Wattle | | | N | x | |
| <i>Acacia myrtifolia</i> | Red-stemmed Wattle | | | N | | X |
| <i>Acacia parramattensis</i> | Parramatta Wattle | | | N | x | |
| <i>Acacia ulicifolia</i> | Prickly Moses | | | N | | X |
| <i>Acer negundo</i> | | Other | | E | x | |
| <i>Adiantum aethiopicum</i> | Common Maidenhair | | | N | | X |
| <i>Agapanthus praecox</i> | African Lily | | | E | x | |
| <i>Allocasuarina torulosa</i> | Forest Oak | | | N | x | X |
| <i>Angophora costata</i> | Sydney Red Gum | | | N | | X |
| <i>Angophora floribunda</i> | Apple | | | N | | X |
| <i>Anredera cordifolia</i> | Madeira Vine | State - AP | | E | x | |
| <i>Araucaria heterophylla</i> | Norfolk Island Pine | | | N-PI | x | |
| <i>Araujia sericifera</i> | Moth vine, Moth plant | Other | | | x | |
| <i>Aristida vagans</i> | Threeawn Speargrass | | | N | | X |
| <i>Arundo donax</i> | Giant Reed | Regional - AP | | E | | X |
| <i>Asparagus aethiopicus</i> | Asparagus fern | State - AP | Y | E | x | X |
| <i>Asparagus asparagoides</i> | Bridal Creeper | State - AP | Y | E | | X |
| <i>Banksia oblongifolia</i> | Fern-leaved Banksia | | | N-PI | | X |
| <i>Bidens pilosa</i> | Cobblers Pegs | | | E | x | |
| <i>Blechnum cartilagineum</i> | Gristle Fern | | | N | | X |
| <i>Brachychiton acerifolius</i> | Flame Tree | | | N-PI | x | |
| <i>Breynia oblongifolia</i> | Coffee Bush | | | N | x | X |
| <i>Bromus catharticus</i> | Prairie Grass | | | E | x | |

| Botanic Name | Common Name | High Threat Exotics | WoNS | Exotic/ Native | Study area | Bushland |
|--------------------------------|-----------------------|---------------------|------|----------------|------------|----------|
| <i>Bursaria spinosa</i> | Blackthorn | | | N - PL | | X |
| <i>Callistemon</i> sp. | | | | N-PI | x | X |
| <i>Camellia</i> sp. | | | | E | x | |
| <i>Cassytha pubescens</i> | | | | N | | X |
| <i>Casuarina glauca</i> | Swamp Oak | | | N-PI | x | |
| <i>Cenchrus clandestinus</i> | Kikuyu | Other | | E | x | |
| <i>Centella asiatica</i> | Indian Pennywort | | | N | x | X |
| <i>Cinnamomum camphora</i> | Camphor Laurel | Other | | E | x | X |
| <i>Clematis aristata</i> | Old Man's Beard | | | N | | X |
| <i>Conyza bonariensis</i> | Flaxleaf Fleabane | | | E | x | X |
| <i>Corymbia maculata</i> | Spotted Gum | | | N-PI | x | |
| <i>Cryptostylis</i> sp. | | | | N | | X |
| <i>Cupressus</i> sp. | | | | E | x | |
| <i>Cyathea australis</i> | Black Tree-fern | | | N | x | |
| <i>Cyathochaeta diandra</i> | | | | N | | X |
| <i>Cynodon dactylon</i> | Couch | | | E | x | |
| <i>Desmodium varians</i> | | | | N | | X |
| <i>Dianella caerulea</i> | | | | N | x | X |
| <i>Dichondra repens</i> | Kidney Weed | | | N | | X |
| <i>Dodonaea triquetra</i> | Large-leaf Hop-bush | | | N | x | |
| <i>Doryanthes excelsa</i> | Gynea Lily | | | N-PI | x | |
| <i>Echinopogon caespitosus</i> | Bushy Hedgehog-grass | | | N | | X |
| <i>Ehrharta erecta</i> | Panic Veldtgrass | | | E | | X |
| <i>Elaeocarpus reticulatus</i> | Blueberry Ash | | | N | x | X |
| <i>Entolasia marginata</i> | Bordered Panic | | | N | x | X |
| <i>Entolasia stricta</i> | Wiry Panic | | | N | | X |
| <i>Eragrostis curvula</i> | African Lovegrass | Other | | E | x | |
| <i>Erythrina</i> sp. | Coral Tree | Other | | E | x | |
| <i>Eucalyptus grandis</i> | Flooded Gum | | | N-PI | x | |
| <i>Eucalyptus fibrosa</i> | Broad-leaved Ironbark | | | N | | X |
| <i>Eucalyptus haemastoma</i> | Scribbly Gum | | | N-PI | x | |
| <i>Eucalyptus paniculata</i> | Grey Ironbark | | | N | | X |

| Botanic Name | Common Name | High Threat Exotics | WoNS | Exotic/ Native | Study area | Bushland |
|---------------------------------|----------------------------|---------------------|------|----------------|------------|----------|
| <i>Eucalyptus pilularis</i> | Blackbutt | | | N | x | X |
| <i>Eucalyptus punctata</i> | Grey Gum | | | N | x | X |
| <i>Eucalyptus resinifera</i> | Red Mahogany | | | N | | X |
| <i>Eucalyptus saligna</i> | Sydney blue gum | | | N | x | X |
| <i>Eucalyptus scias</i> | Large-fruited Red Mahogany | | | N | x | |
| <i>Eucalyptus scoparia</i> | White Gum | | | N-PI | x | |
| <i>Eucalyptus sideroxylon</i> | Mugga Ironbark | | | N- PI | x | |
| <i>Eustrephus latifolius</i> | Wombat Berry | | | N | | X |
| <i>Ficus macrophylla</i> | Moreton Bay Fig | | | N- PI | x | |
| <i>Ficus rubiginosa</i> | Port Jackson Fig | | | N- PI | x | |
| <i>Gahnia aspera</i> | | | | N | | X |
| <i>Genista monspessulana</i> | Montpellier Broom | State - AP | | E | x | |
| <i>Geranium homeanum</i> | | | | N | | X |
| <i>Gleditsia triacanthos</i> | Honey Locust | Other | | E | x | |
| <i>Glycine microphylla</i> | | | | N | | X |
| <i>Glycine tabacina</i> | | | | N | | X |
| <i>Gonocarpus tetragynus</i> | | | | N | | X |
| <i>Grevillea juniperina</i> | | | | N-PI | x | |
| <i>Hakea</i> sp. | | | | N-PI | x | |
| <i>Hardenbergia violacea</i> | Purple Coral Pea | | | N | | X |
| <i>Hibbertia aspera</i> | Rough Guinea Flower | | | N | | X |
| <i>Hibbertia dentata</i> | Trailing Guinea Flower | | | N | | X |
| <i>Homolanthus populifolius</i> | Bleeding Heart | | | N | | X |
| <i>Hypochaeris radicata</i> | Catsear | | | E | x | |
| <i>Imperata cylindrica</i> | Blady Grass | | | N | x | |
| <i>Indigofera australis</i> | Australian Indigo | | | N-PI | | X |
| <i>Ipomoea indica</i> | Morning Glory | Other | | E | | X |
| <i>Jacaranda mimosifolia</i> | | | | E | x | |
| <i>Kunzea ambigua</i> | Tick Bush | | | N - PI | | X |
| <i>Lagerstroemia archeriana</i> | Crepe Myrtle | | | E | x | |

| Botanic Name | Common Name | High Threat Exotics | WoNS | Exotic/ Native | Study area | Bushland |
|------------------------------------|-------------------------|---------------------|------|----------------|------------|----------|
| <i>Lagunaria patersonia</i> | Norfolk Island Hibiscus | | | E | x | |
| <i>Lantana camara</i> | Lantana | State AP | - Y | E | | X |
| <i>Lasiopetalum ferrugineum</i> | | | | N | | X |
| <i>Lepidosperma laterale</i> | | | | N | | X |
| <i>Leptospermum polygalifolium</i> | | | | N | | X |
| <i>Leucopogon juniperinus</i> | Prickly Beard-heath | | | N | | X |
| <i>Ligustrum lucidum</i> | Large-leaved Privet | Other | | E | | X |
| <i>Ligustrum sinense</i> | Small-leaved Privet | Other | | E | | X |
| <i>Lilium formosanum</i> | | Other | | E | | X |
| <i>Lindsaea linearis</i> | Screw Fern | | | N | | X |
| <i>Liquidambar styraciflua</i> | Liquidambar | | | E | x | |
| <i>Lissanthe strigosa</i> | Native Peach | | | N | | X |
| <i>Livistona australis</i> | Cabbage Tree Palm | | | N-PI | x | |
| <i>Lomandra longifolia</i> | Spiny-headed Mat-rush | | | N-PI | x | |
| <i>Lomandra multiflora</i> | | | | N | | X |
| <i>Lomandra obliqua</i> | | | | N | | X |
| <i>Lonicera japonica</i> | Japanese Honeysuckle | Other | | N | | X |
| <i>Lophostemon confertus</i> | Brush Box | | | N-PI | x | |
| <i>Malus sp.</i> | Apple | | | E | x | |
| <i>Mangifera indica</i> | Mango Tree | | | E | x | |
| <i>Melaleuca styphelioides</i> | Prickly-leaved Tree | Tea | | N-PI | x | |
| <i>Micrantheum ericoides</i> | | | | N | | X |
| <i>Microlaena stipoides</i> | Weeping Grass | | | N | | X |
| <i>Monstera deliciosa</i> | Fruit Salad Plant | | | E | | X |
| <i>Morinda jasminoides</i> | Sweet Morinda | | | N | | X |
| <i>Myrsine variabilis</i> | | | | N | | X |
| <i>Notelaea longifolia</i> | Mock Olive | | | N | | X |
| <i>Ochna serrulata</i> | Ochna | Other | | E | X | X |

| Botanic Name | Common Name | High Threat Exotics | WoNS | Exotic/ Native | Study area | Bushland |
|--|-----------------------|---------------------|------|----------------|------------|----------|
| <i>Olea europaea</i> subsp. <i>cuspidata</i> | African Olive | Regional - C | | E | | X |
| <i>Opercularia hispida</i> | | | | N | | X |
| <i>Oplismenus aemulus</i> | Australian Grass | Basket | | N | X | X |
| <i>Oplismenus imbecillis</i> | Creeping Grass | Beard | | N | | X |
| <i>Oxalis perennans</i> | | | | N | | X |
| <i>Ozothamnus diosmifolius</i> | White Dogwood | | | N | | X |
| <i>Pandorea pandorana</i> | Wonga Wonga Vine | | | N | | X |
| <i>Parsonsia straminea</i> | Common Silkpod | | | N | | X |
| <i>Paspalum dilatatum</i> | Paspalum | | | E | X | |
| <i>Passiflora</i> sp. | | | | E | | X |
| <i>Persoonia linearis</i> | Narrow-leaved Geebung | | | N | | X |
| <i>Phyllanthus hirtellus</i> | Thyme Spurge | | | N | | X |
| <i>Pittosporum undulatum</i> | | | | N | | X |
| <i>Plantago debilis</i> | | | | N | | X |
| <i>Plantago lanceolata</i> | Lamb's Tongues | | | E | X | |
| <i>Platylobium formosum</i> | Handsome Flat Pea | | | N | | X |
| <i>Plectranthus parviflorus</i> | Cockspur Flower | | | N | | X |
| <i>Poa affinis</i> | | | | N | | X |
| <i>Polyscias sambucifolia</i> | Elderberry Panax | | | N | | X |
| <i>Pratia purpurascens</i> | Whiteroot | | | N | | X |
| <i>Prunus</i> sp. | Plum Tree | | | E | X | |
| <i>Pseuderanthemum variable</i> | Pastel Flower | | | N | | X |
| <i>Pteridium esculentum</i> | Bracken | | | N | | X |
| <i>Rhododendron</i> sp. | | | | E | X | |
| <i>Robinia pseudocacia</i> | Golden Robinia | | | E | X | |
| <i>Rubus fruticosus</i> species aggregate | | State - AP | Y | E | X | X |
| <i>Rubus parvifolius</i> | Native Raspberry | | | N | | X |
| <i>Senecio madagascariensis</i> | Fireweed | State - AP | | | X | |
| <i>Senna pendula</i> | Cassia, Senna | Other | | | X | |
| <i>Sida rhombifolia</i> | | | | E | X | |
| <i>Smilax glycyphylla</i> | Sweet Sarsaparilla | | | N | | X |
| <i>Solanum aviculare</i> | Kangaroo Apple | | | N-PI | X | |
| <i>Solanum mauritianum</i> | Wild Tobacco | Other | | E | | X |
| <i>Strelitzia nicolai</i> | Bird of Paradise | | | E | X | |

| Botanic Name | Common Name | High Threat Exotics | WoNS | Exotic/ Native | Study area | Bushland |
|---------------------------------|----------------|---------------------|------|----------------|------------|----------|
| <i>Syagrus romanoffianum</i> | Cocos Palm | Other | | E | X | |
| <i>Syncarpia glomulifera</i> | Turpentine | | | N | | X |
| <i>Syzygium paniculatum</i> | Magenta Cherry | | | N-PI | X | |
| <i>Tagetes minuta</i> | Stinking Roger | | | E | X | |
| <i>Themeda australis</i> | Kangaroo Grass | | | N | | X |
| <i>Tradescantia fluminensis</i> | Trad | Other | | E | | X |
| <i>Veronica plebeia</i> | | | | N | | X |
| <i>Xanthorrhoea</i> sp. | | | | N | | X |
| <i>Zieria smithii</i> | Sandfly Zieria | | | N | | X |

E – Exotic; N – Native. N-PI = Native Planted.

| Family | Common Name | Scientific Name | Observation |
|---------------|--------------------------|---------------------------------|-------------|
| Artamidae | Australian Magpie | <i>Gymnorhina tibicen</i> | Observed |
| Artamidae | Grey Butcherbird | <i>Cracticus torquatus</i> | Observed |
| Artamidae | Pied Currawong | <i>Strepera graculina</i> | Heard |
| Cacatuidae | Sulphur-crested Cockatoo | <i>Cacatua galerita</i> | Observed |
| Corcoracidae | White-winged Chough | <i>Corcorax melanorhamphos</i> | Heard |
| Corvidae | Australian Raven | <i>Corvus coronoides</i> | Observed |
| Estrildidae | Red-browed Finch | <i>Neochmia temporalis</i> | Observed |
| Eupetidae | Eastern Whipbird | <i>Psophodes olivaceus</i> | Heard |
| Halcyonidae | Laughing Kookaburra | <i>Dacelo novaeguineae</i> | Observed |
| Meliphagidae | Lewin's Honeyeater | <i>Meliphaga lewinii</i> | Heard |
| Meliphagidae | Noisy Miner | <i>Manorina melanocephala</i> | Observed |
| Monarchidae | Magpie-lark | <i>Grallina cyanoleuca</i> | Observed |
| Psittacidae | Australian King Parrot | <i>Alisterus scapularis</i> | Heard |
| Psittacidae | Eastern Rosella | <i>Platycercus eximius</i> | Observed |
| Psittacidae | Rainbow Lorikeet | <i>Trichoglossus haematodus</i> | Observed |
| Psittaculidae | Musk Lorikeet | <i>Glossopsitta concinna</i> | Observed |

Appendix E: Floristic analysis

| This sheet Version 07/06/2016 - Input data OEH 2013 | | GLD1 | GLD2 | GLD3 | GLD4 | GLD5 | GLD6 | GLD7 | GLD8 | GLD9 | GLD10 | GLD11 | GLD12 | GLD13 | GLD14 | GLD15 | GLD16 | GLD17 | GLD18 | GLD19 | GLD20 | GLD21 | GLD22 | GLD23 | GLD24 | GLD25 | GLD26 | GLD27 | GLD28 | GLD29 | GLD30 | GLD31 | GLD32 | GLD33 | GLD34 | GLD35 | GLD36 | | | | | | | |
|--|-----------|------|------|------|------|------|------|------|------|------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|------|-----|-----|-----|-----|-----|---|
| Community Code | | GLD1 | GLD2 | GLD3 | GLD4 | GLD5 | GLD6 | GLD7 | GLD8 | GLD9 | GLD10 | GLD11 | GLD12 | GLD13 | GLD14 | GLD15 | GLD16 | GLD17 | GLD18 | GLD19 | GLD20 | GLD21 | GLD22 | GLD23 | GLD24 | GLD25 | GLD26 | GLD27 | GLD28 | GLD29 | GLD30 | GLD31 | GLD32 | GLD33 | GLD34 | GLD35 | GLD36 | | | | | | | |
| PCT code | | 1204 | 897 | 830 | 850 | 849 | 1395 | 815 | 771 | 664 | 1061 | 772 | 1143 | 1143 | 882 | 881 | 882 | 1810 | 906 | 888 | 888 | 877 | 888 | 910 | 910 | 920 | 1126 | 1913 | 1237 | 1841 | 888 | 694 | 694 | 1085 | 1253 | 1847 | 1281 | 1214 | 888 | 888 | 888 | 888 | 888 | |
| Positive species count | | 3 | 13 | 33 | 43 | 118 | 72 | 20 | 13 | 29 | 32 | 11 | 21 | 17 | 132 | 7 | 65 | 11 | 51 | 16 | 55 | 26 | 35 | 40 | 40 | 2 | 8 | 9 | 60 | 74 | 30 | 38 | 50 | 60 | 36 | 24 | 69 | 62 | 43 | 38 | 32 | 80 | 36 | |
| Species in the Diagnostic list (OEH) | | 4 | 54 | 58 | 82 | 123 | 121 | 77 | 48 | 99 | 81 | 21 | 91 | 99 | 163 | 43 | 117 | 20 | 70 | 70 | 32 | 42 | 66 | 70 | 123 | 3 | 12 | 9 | 106 | 116 | 62 | 133 | 109 | 132 | 97 | 172 | 110 | 117 | 59 | 82 | 121 | 80 | 78 | |
| Benchmark species count (HNI) | | 2 | 20 | 28 | 29 | 29 | 36 | 18 | 19 | | | 4 | | | 34 | 16 | 34 | | | 19 | 33 | 32 | 25 | 27 | 27 | 1 | 4 | | 40 | | 25 | | 40 | 42 | | 39 | 42 | 30 | 35 | 35 | 35 | 33 | | |
| Benchmark species count (ME) | | 2 | 11 | 31 | 31 | 31 | 31 | | 12 | 27 | 27 | 22 | 17 | 17 | 34 | 22 | 34 | 12 | 26 | 23 | 31 | 31 | 23 | 23 | 1 | 1 | -1 | 32 | 32 | | 32 | 32 | 40 | | 35 | 35 | | 32 | 35 | 35 | 35 | | | |
| Total native species | | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | 23 | | |
| Total diagnostic species | | 0 | 2 | 4 | 0 | 3 | 5 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 3 | 4 | 2 | 1 | 5 | 3 | 0 | 0 | 0 | 12 | 10 | 4 | 4 | 8 | 8 | 4 | 1 | 14 | 10 | 3 | 7 | 6 | 9 | 1 | |
| Required minimum total native species | | 13 | | | | 32 | 41 | 23 | 14 | 24 | 27 | 5 | 24 | 16 | 42 | | 40 | | 24 | 16 | 27 | | | | | | | 34 | 33 | 17 | 33 | 35 | 39 | 32 | | 35 | 42 | | 34 | 33 | | 28 | | |
| Required minimum +ve diagnostic species | Achieved? | Yes | | | | No | No | Yes | Yes | No | No | Yes | No | Yes | No | No | No | No | No | Yes | No | No | No | No | No | No | No | No | No | No | No | Yes | No | No | No | No | No | No | No | No | No | No | No | |
| Required minimum +ve diagnostic species | Achieved? | No | | | | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | No | | |
| No. by which required minimum +ve diagnostic spp. is exceeded | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ratio of actual : required +ve diagnostic species | | 33% | | | | 12% | 19% | 0% | 25% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 6% | 24% | | | | | | | | | | | | | | | | | | | | | | | | | |
| Ratio of +ve diagnostic species : total native species | | 9% | | | | 13% | 22% | 0% | 4% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 0% | 4% | 13% | 17% | | | | | | | | | | | | | | | | | | | | | | | | |
| Standardised ratio comparison (+ve x (req/total) : total native) | | 0.83 | | | | 1.18 | 1.77 | 0.00 | 0.26 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.41 | 0.47 | 1.45 | | | | | | | | | | | | | | | | | | | | | | | | |
| Native Species | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | | |
| Angophora costata | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Angophora floribunda | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Breytia oblongifolia | | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Centella asiatica | | 0 | 1 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Dianella caerulea | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Elaeocarpus reticulatus | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Entolasia marginata | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Eucalyptus punctata | | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Eucalyptus resinifera | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Eucalyptus saligna | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Eustrephus latifolius | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Glycine clandestina | | 0 | 0 | 1 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Lepidosperma laterale | | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Lissanthe strigosa | | 0 | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Morinda jasminoides | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Opilismenus aemulus | | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Pittosporum undulatum | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Pratia purpurascens | | 0 | 1 | 0 | 0 | 1 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Pseuderanthemum variabile | | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Pteridium esculentum | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Rubus sp. | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Strelitzia spp. | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |
| Syncarpia glomulifera | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | |

Analysis of floristic data from plot 4 PCT 1281 was conducted to determine the best-fit PCT for the vegetation located in the southern portion of the subject site. This spreadsheet shows that PCT 1281 represented the best-fit PCT for this area. PCT 1237 was also considered the next best fit, however, this PCT was not chosen for the southern vegetation (vegetation zones 3 and 4) due to landscape and topography factors.

Appendix F: Anabat results

There were no microbat records on the anabat from 9th – 12th November 2018. Therefore, there is no data to show here.

Appendix G: Biodiversity credit report



BAM Credit Summary Report

Proposal Details

| | | |
|--------------------------------|---------------------------------|-------------------------|
| Assessment Id | Proposal Name | BAM data last updated * |
| 00013237/BAAS18159/18/00013238 | Loreto Concept Plan and Stage 1 | 07/12/2020 |
| Assessor Name | Report Created | BAM Data version * |
| Belinda Jane Failes | 15/12/2020 | 34 |
| Assessor Number | BAM Case Status | Date Finalised |
| BAAS18159 | Open | To be finalised |
| Assessment Revision | Assessment Type | |
| 3 | Major Projects | |

* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.

Ecosystem credits for plant communities types (PCT), ecological communities & threatened species habitat

| Zone | Vegetation zone name | TEC name | Current Vegetation integrity score | Change in Vegetation integrity (loss / gain) | Area (ha) | BC Act Listing status | EPBC Act listing status | Species sensitivity to gain class (for BRW) | Biodiversity risk weighting | Potential SAIL | Ecosystem credits |
|-----------------------------|----------------------|-----------|------------------------------------|--|-----------|-----------------------|-------------------------|---|-----------------------------|-----------------|-------------------|
| Blue Gum high forest | | | | | | | | | | | |
| 2 | 1237_native_planted | Not a TEC | 34.1 | 34.1 | 0.35 | | | High Sensitivity to Potential Gain | 2.50 | | 7 |
| | | | | | | | | | | Subtotal | 7 |

Assessment Id
00013237/BAAS18159/18/00013238

Proposal Name
Loreto Concept Plan and Stage 1

Page 1 of 2



BAM Credit Summary Report

| Sydney Turpentine - Ironbark forest | | | | | | | | | | | |
|-------------------------------------|------------|---|------|------|------|--|--|------------------------------------|----------|------|---|
| 1 | 1281_Weedy | Sydney Turpentine-Ironbark Forest in the Sydney Basin Bioregion | 39.7 | 39.7 | 0.05 | Critically Endangered Ecological Community | | High Sensitivity to Potential Gain | 2.50 | TRUE | 1 |
| | | | | | | | | | Subtotal | | 1 |
| | | | | | | | | | Total | | 8 |

Species credits for threatened species

| Vegetation zone name | Habitat condition (Vegetation Integrity) | Change in habitat condition | Area (ha)/Count (no. individuals) | BC Act Listing status | EPBC Act listing status | Biodiversity risk weighting | Potential SAI | Species credits |
|----------------------|--|-----------------------------|-----------------------------------|-----------------------|-------------------------|-----------------------------|---------------|-----------------|
|----------------------|--|-----------------------------|-----------------------------------|-----------------------|-------------------------|-----------------------------|---------------|-----------------|

| | | |
|--------------------------------|---------------------------------|-------------|
| Assessment Id | Proposal Name | Page 2 of 2 |
| 00013237/BAAS18159/18/00013238 | Loreto Concept Plan and Stage 1 | |

