

Climate Change, Wildfires, Heatwaves and Health Impacts in Australia



Nicolas Borchers Arriagada, David M. J. S. Bowman, Andrew J. Palmer and Fay H. Johnston

Abstract Heat-related extreme events, such as wildfires and heatwaves, have historically imposed a burden on Australian society, and according to rigorous and robust scientific literature, it is expected that there will be an increase in frequency, intensity and duration of these types of natural hazards. Within Australia, wildfires and heatwaves are currently responsible for more than 60% of all direct fatalities related to natural hazards, and it is highly likely that this is an underestimation of all deaths as some health impacts are not routinely quantified (e.g. premature death related to air pollution from wildfire smoke exposure). Deaths attributable to heatwaves and fire smoke pollution are more commonly due to exacerbations of pre-existing health conditions, than to specific direct impacts such as heat stroke. Some groups, such as the elderly, infants and those with pre-existing conditions, tend to be more vulnerable to these impacts. Furthermore, evidence suggests that there are synergistic additional impacts when exposed to high temperature and air pollution and that probably health impacts are considerably underestimated in the case of some specific groups such as those with occupational chronic exposure to fire smoke. To avoid increases in public health effects, society at all levels needs to increase its adaptive capacity. Measures need to be taken from a planning and management perspective through to community response at a local level, adequately focusing resources to include vulnerable sectors and population groups.

Keywords Climate change · Bushfires · Wildfires · Heatwaves · Health · Australia

N. Borchers Arriagada (✉) · A. J. Palmer · F. H. Johnston
Menzies Institute for Medical Research, University of Tasmania, Private Bag 23, Hobart, TAS
7000, Australia
e-mail: nicolas.borchers@utas.edu.au

N. Borchers Arriagada
New South Wales Bushfire Risk Management Research Hub, University of Tasmania, Hobart,
TAS, Australia

D. M. J. S. Bowman
School of Natural Sciences, University of Tasmania, Private Bag 55, Hobart, TAS 7001, Australia

© Springer Nature Switzerland AG 2020

R. Akhtar (ed.), *Extreme Weather Events and Human Health*,
https://doi.org/10.1007/978-3-030-23773-8_8

1 Wildfires and Heatwaves and Climate Change: Global Context

Wildfires and heatwaves are important natural hazards that will be exacerbated by climate change. Fire is a process that is complex by nature and is highly variable in how it behaves in space and time; it needs to be ignited, have available fuel to burn and be set in particular weather and topographic conditions that will allow it to successfully spread (Hughes and Alexander 2017). Wildfires, in contrast to prescribed burns, are less predictable processes, with high uncertainty in their behaviour and impacts due to variations in how they start, the fuel being burnt and the existing meteorological conditions that may or may not be beneficial to their spreading.

For heatwaves (or heat events), there is no quantitative universal definition, and different metrics to identify these have been used. According to the World Meteorological Organization and World Health Organization, heatwaves are defined as: “periods of unusually hot and dry or hot and humid weather that have a subtle onset and cessation, a duration of at least two to three days and a discernible impact on human activities. During such periods of hot weather, not only do daytime temperatures reach high values but nocturnal temperatures and humidity levels may also rise well beyond their long-term mean” (McGregor et al. 2015, p. 1). A key feature is a marked departure from typical background weather conditions for the affected geographical region (Nairn and Fawcett 2013). This means, for example, that a heatwave in Tasmania may be considered as moderate weather conditions in another place such as Adelaide, and therefore, the definition of a heatwave is relative to its location.

A wide body of scientific literature supports the fact that global warming will produce more frequent and intense extreme weather events worldwide, and there is very high confidence that these shifting patterns, including those caused by changing temperature and occurrence of heatwaves and fires, have an impact on human health (Bowman et al. 2017; Mora et al. 2017b; Smith et al. 2014). In the Intergovernmental Panel on Climate Change (IPCC) Fifth Assessment Report (AR5), Smith et al. (2014, p. 713) state that there is a “greater risk of injury, disease, and death due to more intensive heatwaves and fires” with some level of climate change.

During a forest fire, the process of combustion produces great amounts of smoke, consequently being an important contributor to air pollution in urban and rural settings. Air pollution is one of the largest threats to humans, one of the main causes of death in the world, and there is no evidence of a safe threshold level of exposure below which no negative effects occur (WHO 2013). It is estimated that outdoor and indoor air pollution (urban and rural sources) is responsible for around 7 million premature annual deaths (1 out of 8 total deaths), mostly from cardiovascular diseases, and from the development of respiratory diseases such as acute respiratory infections (ARIs) and chronic obstructive pulmonary diseases (COPDs) (WHO 2014a). In fact, a study published in 2012 in the *Lancet* found that for the year 2010, household air pollution from solid fuels was the third leading risk factor for global disease burden (Lim et al. 2012). Air pollution produced by landscape fires is not

very different in the way it impacts health, and one of the main differences would be related to exposure. With urban air pollution people usually suffer from chronic exposure, while with landscape fire smoke people might be exposed for short periods of time to very high levels of air pollution. It is estimated that landscape fire smoke is responsible for 340,000 annual deaths globally (Johnston et al. 2012), and for a high number of morbidity effects such as increased respiratory hospital admissions and chronic diseases, asthma attacks, emergency department visits and reduced productivity, among others (Adetona et al. 2016; Black et al. 2017; Duran 2014; Kochi et al. 2010; Liu et al. 2015; Reid et al. 2016; Youssouf et al. 2014). Mortality estimates may have a great variation related to climatic conditions, ranging from 262,000 deaths globally with strong La Nina conditions to 532,000 with strong El Nino conditions (Johnston et al. 2012), and may potentially be underestimated as long-term effects have not been quantified. In fact, landscape fires could well be one of the most relevant emission sources, because of the uncertainty in its nature and its capacity to negatively influence air pollution reduction efforts, producing average increases of annual PM_{2.5} concentration (Larsen et al. 2018) and producing greater impacts in public health (Ford et al. 2018).

According to the World Health Organization (2014b) and assuming no adaptation to climate change, by the year 2030, heat will be responsible for 92,207 (95% confidence intervals: 64,458–121,464) additional deaths globally for people aged 65 and over. This value will rise to 255,486 (191,816–364,002) for the year 2050. Worldwide, a significant amount of people (30%) are currently exposed to extreme heat conditions for at least 20 days per year, and it is expected that by the year 2100, exposure will increase to about 48% in a conservative climate change scenario (Mora et al. 2017b), while severe warming could produce increasingly intolerable conditions (Sherwood and Huber 2010).

These two types of extreme events and their impacts are not necessarily disconnected, and there are often strong associations between them. A heatwave can occur in isolation, but it will increase the risk and likelihood of a severe wildfire (Cardil et al. 2014; Johnston et al. 2011a; Shaposhnikov et al. 2014).

Impacts of extreme weather events related to increased temperatures translate into multiple impacts on human well-being. The objective of this chapter is to integrate research and findings from both government and non-government reports and assessments and peer-reviewed scientific articles to present an overview of the growing health and well-being implications the Australian community will face with increasing risk of wildfires and heatwaves driven by change in climatic conditions. The following sections will present in more detail the health impacts produced by exposure to wildfires and heatwaves, will discuss the conjoined effects of high temperatures and wildfires and will highlight the need to increase adaptation capacities throughout Australian society.

2 Wildfires and Health Impacts in Australia

Wildfires (known as bushfires in Australia) and their consequences have always been a part of the Australian landscape (Hughes and Alexander 2017), and their projected increase in frequency and intensity is of great concern when the effects on human well-being (and health specifically) are considered.

Impacts of wildfires on human well-being are diverse by nature and include:

- Loss of houses and other buildings (Ashe et al. 2009; Blanchi et al. 2014)
- Property damage and destruction (Ashe et al. 2009; Blanchi et al. 2014)
- Threats to water security (Johnston 2009)
- Loss of agricultural crops (Stephenson et al. 2013)
- Fatalities (Blanchi et al. 2014; Haynes et al. 2010)
- Injuries (Cameron et al. 2009)
- Loss of livestock (Whittaker et al. 2012)
- Loss of biodiversity (Stephenson 2010)
- Air pollution emissions and its consequences in human health (Adetona et al. 2016; Cascio 2018; Hyde et al. 2017; Liu et al. 2015; Navarro et al. 2018; Reid et al. 2016)
- Mental health problems (e.g. Post-Traumatic Stress Disorder) (Gibbs et al. 2013; Hughes and Alexander 2017; Reid et al. 2016; Stephenson 2010).

The occurrence of wildfires has a very high impact on Australian society, with a recorded yearly average of 100 fatalities, 3,000 direct injuries and 83 lost homes per year, with an estimated cost of AUD \$12,000 million or 1.3% of GDP for 2005 (Ashe et al. 2009). Blanchi et al. (2014) developed and analysed a data set regarding wildfire with loss of life for the period between 1901 and 2011. Complementing this analysis with official fatality figures reported by the Australian Disaster Resilience Knowledge Hub (AIDR 2018), it is estimated that only 7 fire events were responsible for 71% of all civilian fatalities (Table 1) for the period studied.

Although these numbers seem relevant by themselves, impacts of wildfires go further, negatively impacting the quality of the air that we breathe and imposing substantial health burdens on society. But in general, the study of the effects that fire smoke has on human health tends to be challenging (compared to general urban air pollution) for a myriad of reasons that include (1) the unpredictability of episodes, (2) disperse and heterogeneous spatial and temporal distribution of smoke impacts compared to those of other sources, (3) exposure affecting both urban and rural areas with differing levels of population density, (4) higher peak concentrations of PM_{2.5}, (5) lack of availability of monitoring data in rural areas affected by fires, (6) exposure of small populations not representative of broader community and (7) short duration of episodes (even on large populations) that don't allow for detection of significant changes (Johnston et al. 2011b; Salimi et al. 2017). Also, the study of health impacts due to exposure to fire smoke has had less attention than those related to urban air pollution, but there is still a considerable amount of literature available, and it has been found that health outcomes are significantly diverse in the nature and size of populations affected as shown in Fig. 1.

Table 1 Major fire events in Australia according to the number of direct civilian fatalities

#	Date of fire	Bushfire	Location	Number of civilian fatalities	Number of homes destroyed
1	Feb/1926–Mar/1926	1926 bushfires	Gippsland, Victoria	60	550
2	Jan/1939	Black Friday bushfires	Victoria	71	1,000
3	14/Jan–14/Feb 1944	1944 bushfires	Victoria	15–20	>500
4	7/Feb/1967	Tasmanian “Black Tuesday” bushfires	Hobart, Tasmania	62	1,400
5	Jan/1969	1969 bushfires	Lara, Victoria	23	230
6	Feb/1983	Ash Wednesday bushfires	South Australia	75	2,000
7	Feb/2009	Black Saturday bushfires	Victoria	173	2,029

Adapted from Blanchi et al. (2014) and AIDR (2018)




Fig. 1 Total public health impacts of wildfire smoke or PM_{2.5}. Adapted from Cascio (2018)

There is robust evidence for an increase in respiratory morbidity and all-cause mortality, and mixed findings in the case of both cardiovascular morbidity and cause-specific mortality due to short-term exposure to fire smoke (Black et al. 2017; Cascio 2018; Reisen et al. 2015; Youssouf et al. 2014). Exacerbation of respiratory health conditions is likely to be the most common immediate impact (Black et al. 2017). Long-term health effects due to exposure to fire smoke should also be taken into account, as some studies have shown that a considerable amount of population may be exposed to an average annual increase of more than $0.75 \mu\text{g}/\text{m}^3$ of $\text{PM}_{2.5}$ (Cascio 2018; Rappold et al. 2017), in which case public health impacts in terms of morbidity and mortality will likely be considerable.

Epidemiological studies following Australian bushfire smoke events have found positive associations for multiple effects, especially respiratory (Table 2), while results for cardiovascular morbidity and mortality in general have been more variable. In 2010, Morgan et al. (2010) studied fires occurring in Sydney between 1994 and 2002 and found no consistency in associations between wildfire particulate matter (PM_{10}) with cardiovascular admissions and mortality. Later in 2011, Johnston et al. (2011a) analysed fires between 1997 and 2004 and found that smoke events could produce a 5% increase in non-accidental and cardiovascular mortality. Evidence regarding cardiovascular impacts has been strengthened, as Dennekamp et al. (2015) and Haikerwal et al. (2015) have found positive associations between exposure to smoke from the 2006 to 2007 wildfires in Victoria and out-of-hospital cardiac arrest (OHCA) and ischaemic heart disease (IHD). Results seen in Australian studies are consistent with those done elsewhere, where evidence is robust around positive associations between fire smoke and respiratory morbidity, evidence is increasing regarding positive associations between smoke and respiratory infections and all-cause mortality, but there are still important gaps in research with respect to effects on cardiovascular morbidity, specific causes of mortality and other endpoints such as birth outcomes and mental health (Reid et al. 2016).

Similar to the effects of air pollution in general, certain population groups tend to be more vulnerable to these outcomes, such as pregnant women, the elderly, infants, smokers, firefighters and people with pre-existing respiratory or cardiovascular conditions (Hughes and Alexander 2017; Rappold et al. 2017; Youssouf et al. 2014). Australian indigenous people might also be affected more strongly than the rest of the population (Hanigan et al. 2008). Johnston et al. (2007) found effects to be higher in Indigenous people compared to the rest of the population: associations for COPD and asthma were more than double, and they appeared to have a higher risk of being admitted for cardio-respiratory effects. This likely is a reflection of how Indigenous people are at higher risk from environmental hazards such as air pollution (Johnston et al. 2007). Some areas of Australia tend to be at higher risk of suffering bushfires and the largest effects will be experienced in high-density urban areas, where concentrated populations are potentially exposed to fire smoke.

Table 2 Examples of positive associations between fire smoke PM and health effects—Australian studies

Health outcome	Place	Year/period/episode	Reference
Increased respiratory and cardiovascular emergency ambulance dispatches (EAD)	Sydney Greater Metropolitan Region	2004–2015	Salimi et al. (2017)
Increased emergency department (ED) attendances for asthma	Victoria	1 Dec 2006–31 Jan 2007	Haikerwal et al. (2016)
Increased out-of-hospital cardiac arrest (OHCA)	Melbourne	July 2006–June 2007	Dennekamp et al. (2015)
Increased out-of-hospital cardiac arrest (OHCA) and ischaemic heart disease (IHD)	Victoria	1 Dec 2006–31 Jan 2007	Haikerwal et al. (2015)
Increased non-accidental and cardiovascular mortality	Sydney	January 1994–June 2007	Johnston et al. (2011a)
Increased ED attendances for all non-trauma conditions, respiratory conditions, asthma, and chronic obstructive pulmonary disease (COPD)	Sydney	1996–2007	Johnston et al. (2014)
Increased admissions for respiratory infections for Indigenous people	Darwin	Fire seasons between 1996 and 2005 (1 April–30 November)	Hanigan et al. (2008)

3 Heatwaves and Health Impacts in the Australian Context

Australians have been heavily impacted from extreme heat events throughout history. It is a country that is highly vulnerable to changing climatic conditions, and it is expected that the number of hot days, warm nights and heatwaves will increase by the end of the twenty-first century (Cowan et al. 2014). Australian society will experience various types of impacts, such as:

- increased human morbidity and mortality (Cheng et al. 2018; Lindstrom et al. 2013; McGregor et al. 2015; Mora et al. 2017b; Tong et al. 2014; Turner et al. 2013; Wilson et al. 2013)
- increased risk for outdoor activities (Nairn and Fawcett 2013)
- stress for people, animals and plants (Sherwood and Huber 2010)
- increased bushfire risk (Cardil et al. 2014)
- damage to crop and vegetation (Wreford and Neil Adger 2010)
- increased demand for resources (e.g. water, energy, etc.) (Zuo et al. 2015)
- stress on infrastructure such as roads and buildings (McEvoy et al. 2012)
- negative impacts in tourism (Nairn and Fawcett 2013).

Heatwaves will be responsible, in the case of Australasia (Australia and New Zealand), for an estimated additional 217 (132–345) yearly deaths by the year 2030 and for 605 (434–980) by 2050 (WHO 2014b). However, these numbers may increase considerably when in the presence of an extreme heat event. For example, a major heatwave struck Europe during the summer of 2003 and was responsible for over 70,000 deaths (Robine et al. 2008). In Australia, the highest number of recorded deaths from natural hazards are attributable to heatwaves, and even with their relative importance in terms of fatalities, these have not been studied with similar intensity as other natural hazards such as floods and storms (Coates et al. 2014; Lee 2014). Only 11 events (out of 350) are responsible for more than 30% of all recorded heatwave-related deaths for the period 1844–2011 (Table 3). Victoria, New South Wales and South Australia have the highest number of fatalities, with most of these occurring during summer months.

It is probable that many earlier studies underestimated the real impact of heatwaves on Australians, as only uncommon but specific heat-related diseases such as heat stroke were attributed to the heat event (Coates et al. 2014). However, epidemiological studies increasingly use statistical methods to attribute excess hospital admissions, regardless of the specific medical diagnosis, to particular events such as heatwaves, and the pathophysiological processes that lead from elevated body temperature to death through direct toxicity and exacerbations of cardiovascular, renal and metabolic conditions are well characterised (Mora et al. 2017a).

Australian regions are prone to high temperatures, hot summer and extreme heat, and the association of these conditions with adverse health impacts has been well studied (Bi et al. 2011), with a more intense focus on mortality rather than morbidity effects (Dalip et al. 2015; Loughnan et al. 2010) and with less available studies for the southern hemisphere including Australia (Loughnan et al. 2010). Health

Table 3 Significant heat events in Australia, 1844–2011 (Coates et al. 2014)

#	Date of event	Area affected	Total heat-associated deaths
1	January–February 1879	NSW, Vic	22
2	October 1895–January 1896	WA, SA, Vic, Qld, NSW	435
3	January 1906	NSW, SA	28
4	January 1908	Vic, SA, NSW	213
5	January 1939	Vic, SA, NSW	420
6	January 1940	Qld, NSW	65
7	February 1955	Perth (WA)	30
8	January–February 1959	Melbourne (Vic)	145
9	January 1960	Greater Sydney (NSW)	25
10	January 2000	Southeast Qld	22
11	January–February 2009	Vic, SA	432
Total deaths			1837

impacts include dehydration, heat strokes, cardiovascular morbidity effects, renal diseases and death. Recent studies (Table 4) have addressed the associations between heat events and multiple health effects in various cities around Australia and have found increased adverse impacts that range from increased use of health services to increased total mortality.

The very young, the elderly (especially those 75+) and people with pre-existing medical conditions are especially vulnerable, while more socially advantaged groups (in terms of socioeconomic status) tend to have lower impacts (Cheng et al. 2018; Coates et al. 2014; Dalip et al. 2015; Xiao et al. 2017). Other population characteristics such as living in urban or rural areas (Jegasothy et al. 2017), gender (Xiao et al. 2017) and geographical location (Tong et al. 2014) may define the level of vulnerability and adaptability that exists towards heat-related health impacts. Cowan et al. (2014) estimate that changes in frequency and duration of heat events will affect mainly northern tropical regions of Australia, while higher maximum temperatures will be felt in southern Australia.

Table 4 Examples of positive associations between heatwaves and health effects—Australian studies

Health outcome	Place	Year/period/episode	References
Increased health service utilisation	New South Wales	2005–2015	Jegasothy et al. (2017)
Increased mortality in the elderly	Brisbane, Melbourne, Sydney	1988–2009	Tong et al. (2015)
Increased emergency department visits	Brisbane	Summer seasons (December–February) from 2000 to 2012	Toloo et al. (2014)
Increased emergency hospital admissions for renal diseases in children (0–14 years)	Brisbane	1 Jan 1996–31 Dec 2005	Wang et al. (2014)
Increased all-cause, cardiovascular and respiratory mortality and hospital admissions for heat-related injuries, dehydration, and other fluid disorders	Sydney	Mortality (1997–2007) Hospital admissions (1997–2010)	Wilson et al. (2013)
Significantly increased total hospital admissions, emergency department presentations, general medical admissions, and rise in number of deaths	Melbourne	2009 heatwave (Jan/Feb 2009)	Lindstrom et al. (2013)
Increased daily mortality, total emergency department presentations, renal-related emergency department presentations	Perth	1994–2008	Williams et al. (2012)
Increased ambulance call-outs, renal morbidity for elders, heat-related admissions, ischaemic heart disease in the 15–64 age group, total mortality particularly in the 15–64 age group	Adelaide	Summers of 2008 and 2009	Nitschke et al. (2011)

4 Wildfire and Heatwaves: Relative and Combined Effects

Severe fire and heat events have been recorded and will likely increase in the future, producing important environmental, social and economic consequences (Bowman et al. 2017). For Australia, 2017 was its third hottest year on record, including highest historical maximum temperatures during winter and very low rainfall, shaping a perfect scenario for heat-related extreme weather events such as wildfires and heatwaves to develop (Steffen et al. 2018).

Heatwaves and wildfires were responsible for more than 65% of the 8,256 direct fatalities attributed to natural hazards in Australia for the period between 1900 and 2011 (Fig. 2). Heatwaves alone accounted for more fatalities than all other natural hazards (>50% of deaths), while wildfires were responsible for more than 10% of these deaths (Coates et al. 2014), with most wildfire fatalities (>65%) concentrated in the State of Victoria (Blanchi et al. 2014). As we discussed above in the case of wildfires though, it is highly likely that the magnitude of these impacts would increase considerably if indirect deaths caused by exposure to fire smoke were included. Increases in air pollution can have a large public health impact if extensive populations are affected. For example, Horsley et al. (2018) estimated that 183 days of exposure to landscape fire smoke were responsible for 197 premature deaths, 436 cardiovascular hospitalisations and 787 respiratory hospitalisations in a period of 13 years (2001–2013) in Sydney alone. This represents almost 23% of all direct fatalities recorded for Australia in a 111-year period.

As mentioned earlier, days of high temperature (heat events) have a strong influence on increasing the likelihood of large wildfires, especially when combined with high wind. Conditions for fires become highly favourable, and as moisture content decreases the probability for ignition increases, and fires can burn more intensely and rapidly (Cardil et al. 2014). Additionally, there may even be interactions in mortality effects produced by exposure to heatwaves and fire smoke (Katsouyanni et al.




Fig. 2 Total fatalities from bushfires and extreme heat compared with other Australian natural hazards between 1900 and 2011. Adapted from Coates et al. (2014)

1993; Shaposhnikov et al. 2014). In 1987, a major heatwave impacted Greece, and high levels of smoke and ozone were present during the same period. Katsouyanni et al. (1993) found statistically significant interactions between high temperatures ($>30^{\circ}\text{C}$) and sulphur dioxide (SO_2), and statistically suggestive interactions for ozone and smoke with high temperatures. In 2010, Moscow experienced a disastrous heatwave with high levels of air pollution simultaneously, with 44 days of continuous heat during major heat event and PM_{10} exceeding $300\text{ }\mu\text{g}/\text{m}^3$ on multiple days. During this period, there was an estimated near 11,000 excess deaths, almost doubling the expected deaths when considering both phenomena as if they were independent (Shaposhnikov et al. 2014). According to the authors, the interactions between pollution from fire smoke and high temperatures were responsible for more than 2000 of these deaths. Different studies have been undertaken in Australian cities, and there is suggestion for a synergistic effect between high temperature and air pollution (PM_{10} , SO_2 and ozone) and that this impact (excess mortality) is additional to that of the sum of effects from each exposure variable (Dean and Green 2018).

Furthermore, there are fewer studies that relate the effects of occupational exposure (e.g. firefighters) to fire smoke (Adetona et al. 2016). These types of professionals are part of the population segments that are particularly more vulnerable to health effects associated with fire smoke exposure (Youssouf et al. 2014). The exposure to smoke that firefighters experience is very different to that of the rest of the population, and they work very close to the actual sources of emission and are exposed to high concentration levels of particulate matter and other pollutants (Adetona et al. 2016). Some of the effects experienced by firefighters include the increase of pulmonary and systemic inflammation and a decrease in lung function (Youssouf et al. 2014). A study carried on in Australia assessed exposure of firefighters to smoke during prescribed burns and wildfires and found that a considerable proportion of the firefighters had high to very high levels of exposure, and this would result in an increased likelihood of having acute and chronic health effects (Reisen et al. 2011).

It is very likely that the total public health impacts of fire smoke exposure have consistently been underestimated. There appears to an opportunity and need to adequately assess the effects produced by a conjoined exposure to fire smoke and heatwaves, as the synergistic impacts might be higher than the sum of its parts. Also, it seems relevant to consider impacts due to occupational exposure and find ways in which this could be reduced.

5 Adaptation to Heat-Related Extreme Weather Events

As climate change shapes the future of Australia with increasing heat-related extreme weather events, there will be a higher demand for resources, capacities and planning that should be available to approach these issues. There is an urgent need to act regarding future potential natural hazards produced by climate change (Steffen et al. 2018). Wildfires and heatwaves will become a norm for Australian society, and it is necessary that authorities and decision-makers focus adequate amount and type

of resources on reducing risks (Clayton et al. 2014; Cleland et al. 2011; Nairn and Fawcett 2013). Actions such as increasing effectiveness of communication channels and levels with the community (Fish et al. 2017), education programs aimed at changing behaviour (Richardson et al. 2012; Walker and Salt 2006), improving emergency response and incorporating wildfire and heatwave events into the planning and management of our built environment and social processes are necessary to improving the adaptive capacity that systems will have in an ever-changing environment (Tong et al. 2014). Resources must be focused on increasing adaptation to climate change, improving and increasing preparedness for bushfire disasters (Johnston 2009) and extreme heat events, including improvement of public health services, identification of vulnerable sectors and population groups and development of early warning systems (Lowe et al. 2016; Smith et al. 2014).

6 Conclusions

As global warming imposes important changes in climate and weather, with increasing temperatures and levels of drought, Australians will experience a higher risk of suffering from extreme wildfires and intense heatwaves. The many impacts will probably be distributed unequally throughout society. Infants, the elderly, labourers and people with pre-existing conditions, lower socioeconomic status or belonging to indigenous groups may be at higher risk of suffering from smoke and heat-related illnesses, while those living at the urban–rural interface are at higher risk of being directly affected by wildfires. Geographically, population exposure and density differ around Australia, and studies suggest that the Southeast of Australia has higher risk of mortality and disease from wildfire and smoke-related impacts while South Australia presents the highest heat-related mortality rates, although most fatalities have concentrated in Victoria, NSW and South Australia. There are some important aspects that have had relatively limited consideration, such as the possible synergistic adverse health impacts of simultaneous exposure to fire smoke and heatwaves, and the more extreme personal exposures of professionals who work outside during heatwaves and bushfire air pollution events. Necessary steps should be taken at federal, state and local levels to ensure safety of communities and to minimise the loss of resources, lives and well-being, especially focusing on the sectors of society that are more vulnerable due to their physical locations, social and economic disadvantage or higher-risk occupational groups.

References

- Adetona, O., Reinhardt, T. E., Domitrovich, J., Broyles, G., Adetona, A. M., Kleinman, M. T. ... Naeher, L. P. (2016). Review of the health effects of wildland fire smoke on wildland firefighters and the public. *Inhalation Toxicology*. <https://doi.org/10.3109/08958378.2016.1145771>.

- AIDR. (2018). *Australian Disaster Resilience Knowledge Hub: Disasters, Fire—Bushfire*. Melbourne: Australian Institute for Disaster Resilience. Retrieved from <https://knowledge.aidr.org.au/search?keywords=&collection=1316&categories=Fire-Bushfire&from=&to=®ion=&type=&page=1&results=100>.
- Ashe, B., McAneney, K. J., & Pitman, A. J. (2009). Total cost of fire in Australia. *Journal of Risk Research*. <https://doi.org/10.1080/13669870802648528>.
- Bi, P., Williams, S., Loughnan, M., Lloyd, G., Hansen, A., Kjellstrom, T. ... Saniotis, A. (2011). The effects of extreme heat on human mortality and morbidity in Australia: Implications for public health. *Asia-Pacific Journal of Public Health/Asia-Pacific Academic Consortium for Public Health*. <https://doi.org/10.1177/1010539510391644>.
- Black, C., Tesfaigzi, Y., Bassein, J. A., & Miller, L. A. (2017). Wildfire smoke exposure and human health: Significant gaps in research for a growing public health issue. *Environmental Toxicology and Pharmacology*, 55, 186–195. <https://doi.org/10.1016/J.ETAP.2017.08.022>.
- Blanchi, R., Leonard, J., Haynes, K., Opie, K., James, M., & de Oliveira, F. D. (2014). Environmental circumstances surrounding bushfire fatalities in Australia 1901–2011. *Environmental Science and Policy*. <https://doi.org/10.1016/j.envsci.2013.09.013>.
- Bowman, D. M. J. S., Williamson, G. J., Abatzoglou, J. T., Kolden, C. A., Cochrane, M. A., & Smith, A. M. S. (2017). Human exposure and sensitivity to globally extreme wildfire events. *Nature Ecology & Evolution*, 1(58), 1–6. <https://doi.org/10.1038/s41559-016-0058>.
- Cameron, P. A., Mitra, B., Fitzgerald, M., Scheinkestel, C. D., Stripp, A., Batey, C. ... Cleland, H. (2009). Black Saturday: The immediate impact of the February 2009 bushfires in Victoria, Australia. *Medical Journal of Australia*. <https://doi.org/10.5694/j.1326-5377.2009.tb02666.x>.
- Cardil, A., Salis, M., Spano, D., Delogu, G., & Terrén, D. M. (2014). Large wildland fires and extreme temperatures in Sardinia (Italy). *IForest*. <https://doi.org/10.3832/ifor1090-007>.
- Cascio, W. E. (2018). Wildland fire smoke and human health. *Science of the Total Environment*. <https://doi.org/10.1016/j.scitotenv.2017.12.086>.
- Cheng, J., Xu, Z., Bambrick, H., Su, H., Tong, S., & Hu, W. (2018). Heatwave and elderly mortality: An evaluation of death burden and health costs considering short-term mortality displacement. *Environment International*. <https://doi.org/10.1016/j.envint.2018.03.041>.
- Clayton, H., Mylek, M. R., Schirmer, J., Cary, G. J., & Dovers, S. R. (2014). Exploring the use of economic evaluation in Australian wildland fire management decision-making. *International Journal of Wildland Fire*. <https://doi.org/10.1071/WF13140>.
- Cleland, H. J., Proud, D., Spinks, A., & Wasiak, J. (2011). Multidisciplinary team response to a mass burn casualty event: Outcomes and implications. *Medical Journal of Australia*. <https://doi.org/10.5694/j.1326-5377.2011.tb03110.x>.
- Coates, L., Haynes, K., O'Brien, J., McAneney, J., & De Oliveira, F. D. (2014). Exploring 167 years of vulnerability: An examination of extreme heat events in Australia 1844–2010. *Environmental Science and Policy*. <https://doi.org/10.1016/j.envsci.2014.05.003>.
- Cowan, T., Purich, A., Perkins, S., Pezza, A., Boschat, G., & Sadler, K. (2014). More frequent, longer, and hotter heat waves for Australia in the twenty-first century. *Journal of Climate*. <https://doi.org/10.1175/JCLI-D-14-00092.1>.
- Dalip, J., Phillips, G. A., Jelinek, G. A., & Weiland, T. J. (2015). Can the elderly handle the heat? A retrospective case-control study of the impact of heat waves on older patients attending an inner city Australian emergency department. *Asia-Pacific Journal of Public Health*. <https://doi.org/10.1177/1010539512466428>.
- Dean, A., & Green, D. (2018). Climate change, air pollution and human health in Sydney, Australia: A review of the literature. *Environmental Research Letters*, 13(5), 53003. Retrieved from <http://stacks.iop.org/1748-9326/13/i=5/a=053003>.
- Dennekamp, M., Straney, L. D., Erbas, B., Abramson, M. J., Keywood, M., Smith, K. ... Tonkin, A. M. (2015). Forest fire smoke exposures and out-of-hospital cardiac arrests in Melbourne, Australia: A case-crossover study. *Environmental Health Perspectives*. <https://doi.org/10.1289/ehp.1408436>.

- Duran, S. (2014). Evidence review: Wildfire smoke and public health risk. Vancouver, British Columbia: BC Centre for Disease Control. Retrieved from http://www.bccdc.ca/resource-gallery/Documents/Guidelines%and%Forms/Guidelines%and%Manuals/Health-Environment/WFSG_EvidenceReview_WildfireSmoke_FINAL_v3_edstrs.pdf.
- Fish, J. A., Peters, M. D. J., Ramsey, I., Sharplin, G., Corsini, N., & Eckert, M. (2017). Effectiveness of public health messaging and communication channels during smoke events: A rapid systematic review. *Journal of Environmental Management*. <https://doi.org/10.1016/j.jenvman.2017.02.012>.
- Ford, B., Val Martin, M., Zelasky, S. E., Fischer, E. V., Anenberg, S. C., Heald, C. L., & Pierce, J. R. (2018). Future fire impacts on smoke concentrations, visibility, and health in the contiguous United States. *GeoHealth*. <https://doi.org/10.1029/2018GH000144>.
- Gibbs, L., Waters, E., Bryant, R. A., Pattison, P., Lusher, D., Harms, L. ... Forbes, D. (2013). Beyond bushfires: Community, resilience and recovery—A longitudinal mixed method study of the medium to long term impacts of bushfires on mental health and social connectedness. *BMC Public Health*. <https://doi.org/10.1186/1471-2458-13-1036>.
- Haikerwal, A., Akram, M., Del Monaco, A., Smith, K., Sim, M. R., Meyer, M. ... Dennekamp, M. (2015). Impact of fine particulate matter (PM_{2.5}) exposure during wildfires on cardiovascular health outcomes. *Journal of the American Heart Association*. <https://doi.org/10.1161/JAHA.114.001653>.
- Haikerwal, A., Akram, M., Sim, M. R., Meyer, M., Abramson, M. J., & Dennekamp, M. (2016). Fine particulate matter (PM_{2.5}) exposure during a prolonged wildfire period and emergency department visits for asthma. *Respirology*. <https://doi.org/10.1111/resp.12613>.
- Hanigan, I. C., Johnston, F. H., & Morgan, G. G. (2008). Vegetation fire smoke, indigenous status and cardio-respiratory hospital admissions in Darwin, Australia, 1996–2005: A time-series study. *Environmental Health: A Global Access Science Source*. <https://doi.org/10.1186/1476-069X-7-42>.
- Haynes, K., Handmer, J., McAneney, J., Tibbits, A., & Coates, L. (2010). Australian bushfire fatalities 1900–2008: Exploring trends in relation to the “Prepare, stay and defend or leave early” policy. *Environmental Science and Policy*. <https://doi.org/10.1016/j.envsci.2010.03.002>.
- Horsley, J. A., Broome, R. A., Johnston, F. H., Cope, M., & Morgan, G. G. (2018). Health burden associated with fire smoke in Sydney, 2001–2013. *The Medical Journal of Australia*, 208(7), 309–310.
- Hughes, L., & Alexander, D. (2017). *Climate change and the Victorian bushfire threat: Update 2017*. Sydney: Climate Council of Australia.
- Hyde, J. C., Yedinak, K. M., Talhelm, A. F., Smith, A. M. S., Bowman, D. M. J. S., Johnston, F. H. ... Tinkham, W. T. (2017). Air quality policy and fire management responses addressing smoke from wildland fires in the United States and Australia. *International Journal of Wildland Fire*, 26(5), 347–363. Retrieved from <https://doi.org/10.1071/WF16154>.
- Jegasothy, E., McGuire, R., Nairn, J., Fawcett, R., & Scalley, B. (2017). Extreme climatic conditions and health service utilisation across rural and metropolitan New South Wales. *International Journal of Biometeorology*. <https://doi.org/10.1007/s00484-017-1313-5>.
- Johnston, F. H. (2009). Bushfires and human health in a changing environment. *Australian Family Physician*.
- Johnston, F. H., Bailie, R. S., Pilotto, L. S., & Hanigan, I. C. (2007). Ambient biomass smoke and cardio-respiratory hospital admissions in Darwin, Australia. *BMC Public Health*. <https://doi.org/10.1186/1471-2458-7-240>.
- Johnston, F., Hanigan, I., Henderson, S., Morgan, G., & Bowman, D. (2011a). Extreme air pollution events from bushfires and dust storms and their association with mortality in Sydney, Australia 1994–2007. *Environmental Research*. <https://doi.org/10.1016/j.envres.2011.05.007>.
- Johnston, F. H., Hanigan, I. C., Henderson, S. B., Morgan, G. G., Portner, T., Williamson, G. J., & Bowman, D. M. J. S. (2011b). Creating an integrated historical record of extreme particulate air pollution events in Australian cities from 1994 to 2007. *Journal of the Air and Waste Management Association*. <https://doi.org/10.3155/1047-3289.61.4.390>.

- Johnston, F. H., Henderson, S. B., Chen, Y., Randerson, J. T., Marlier, M., DeFries, R. S. ... Brauer, M. (2012). Estimated global mortality attributable to smoke from landscape fires. *Environmental Health Perspectives*, 120(5), 695–701. <https://doi.org/10.1289/ehp.1104422>.
- Johnston, F. H., Purdie, S., Jalaludin, B., Martin, K. L., Henderson, S. B., & Morgan, G. G. (2014). Air pollution events from forest fires and emergency department attendances in Sydney, Australia 1996–2007: A case-crossover analysis. *Environmental Health: A Global Access Science Source*. <https://doi.org/10.1186/1476-069X-13-105>.
- Katsouyanni, K., Pantazopoulou, A., Touloumi, G., Tselepidaki, L., Moustiris, K., Asimakopoulos, D. ... Trichopoulos, D. (1993). Evidence for interaction between air pollution and high temperature in the causation of excess mortality. *Archives of Environmental Health*. <https://doi.org/10.1080/00039896.1993.9940365>.
- Kochi, I., Donovan, G. H., Champ, P. A., & Loomis, J. B. (2010). The economic cost of adverse health effects from wildfire-smoke exposure : A review. *International Journal of Wildland Fire*, 19, 803–817. Retrieved from <http://dx.doi.org/10.1071/WF09077>.
- Larsen, A. E., Reich, B. J., Ruminiski, M., & Rappold, A. G. (2018). Impacts of fire smoke plumes on regional air quality, 2006–2013. *Journal of Exposure Science and Environmental Epidemiology*. <https://doi.org/10.1038/s41370-017-0013-x>.
- Lee, W. V. (2014). Historical global analysis of occurrences and human casualty of extreme temperature events (ETEs). *Natural Hazards*. <https://doi.org/10.1007/s11069-013-0884-7>.
- Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H. ... Ezzati, M. (2012). A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the Global Burden of Disease Study 2010. *The Lancet*. [https://doi.org/10.1016/S0140-6736\(12\)61766-8](https://doi.org/10.1016/S0140-6736(12)61766-8).
- Lindstrom, S. J., Nagalingam, V., & Newnham, H. H. (2013). Impact of the 2009 Melbourne heatwave on a major public hospital. *Internal Medicine Journal*. <https://doi.org/10.1111/imj.12275>.
- Liu, J. C., Pereira, G., Uhl, S. A., Bravo, M. A., & Bell, M. L. (2015). A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environmental Research*. <https://doi.org/10.1016/j.envres.2014.10.015>.
- Loughnan, M. E., Nicholls, N., & Tapper, N. J. (2010). The effects of summer temperature, age and socioeconomic circumstance on Acute Myocardial Infarction admissions in Melbourne, Australia. *International Journal of Health Geographics*. <https://doi.org/10.1186/1476-072X-9-41>.
- Lowe, R., García-Díez, M., Ballester, J., Creswick, J., Robine, J. M., Herrmann, F. R., & Rodó, X. (2016). Evaluation of an early-warning system for heat wave-related mortality in Europe: Implications for sub-seasonal to seasonal forecasting and climate services. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph13020206>.
- McEvoy, D., Ahmed, I., & Mullett, J. (2012). The impact of the 2009 heat wave on Melbourne's critical infrastructure. *Local Environment*. <https://doi.org/10.1080/13549839.2012.678320>.
- McGregor, G. R., Bessemoulin, P., Ebi, K. L., & Menne, B. (2015). *Heatwaves and health: Guidance on warning-system development*. Geneva, Switzerland.
- Mora, C., Counsell, C. W. W., Bielecki, C. R., & Louis, L. V. (2017a). Twenty-seven ways a heat wave can kill you: Deadly heat in the era of climate change. *Circulation: Cardiovascular Quality and Outcomes*. <https://doi.org/10.1161/CIRCOUTCOMES.117.004233>.
- Mora, C., Dousset, B., Caldwell, I. R., Powell, F. E., Geronimo, R. C., Bielecki, C. R. ... Trauer-nicht, C. (2017b). Global risk of deadly heat. *Nature Climate Change*. <https://doi.org/10.1038/nclimate3322>.
- Morgan, G., Sheppeard, V., Khalaj, B., Ayyar, A., Lincoln, D., Jalaludin, B. ... Lumley, T. (2010). Effects of bushfire smoke on daily mortality and hospital admissions in Sydney, Australia. *Epidemiology*. <https://doi.org/10.1097/EDE.0b013e3181c15d5a>.
- Nairn, J., & Fawcett, R. (2013). *Defining heatwaves: Heatwave defined as a heat-impact event servicing all community and business sectors in Australia*. CAWCR Technical Report 551.5250994.

- Navarro, K., Schweizer, D., Balmes, J., & Cisneros, R. (2018). A review of community smoke exposure from wildfire compared to prescribed fire in the United States. *Atmosphere*. <https://doi.org/10.3390/atmos9050185>.
- Nitschke, M., Tucker, G. R., Hansen, A. L., Williams, S., Zhang, Y., & Bi, P. (2011). Impact of two recent extreme heat episodes on morbidity and mortality in Adelaide, South Australia: A case-series analysis. *Environmental Health: A Global Access Science Source*. <https://doi.org/10.1186/1476-069X-10-42>.
- Rappold, A. G., Reyes, J., Pouliot, G., Cascio, W. E., & Diaz-Sanchez, D. (2017). Community Vulnerability to Health Impacts of Wildland Fire Smoke Exposure. *Environmental Science and Technology*. <https://doi.org/10.1021/acs.est.6b06200>.
- Reid, C. E., Brauer, M., Johnston, F. H., Jerrett, M., Balmes, J. R., & Elliott, C. T. (2016). Critical review of health impacts of wildfire smoke exposure. *Environmental Health Perspectives*, 124(9), 1334–1343. <https://doi.org/10.1289/ehp.1409277>.
- Reisen, F., Duran, S. M., Flannigan, M., Elliott, C., & Rideout, K. (2015). Wildfire smoke and public health risk. *International Journal of Wildland Fire*. <https://doi.org/10.1071/WF15034>.
- Reisen, F., Hansen, D., & Meyer, (Mick) P. (2011). Exposure to bushfire smoke during prescribed burns and wildfires: Firefighters' exposure risks and options. *Environment International*. <https://doi.org/10.1016/j.envint.2010.09.005>.
- Richardson, L. A., Champ, P. A., & Loomis, J. B. (2012). The hidden cost of wildfires: Economic valuation of health effects of wildfire smoke exposure in Southern California. *Journal of Forest Economics*. <https://doi.org/10.1016/j.jfe.2011.05.002>.
- Robine, J. M., Cheung, S. L. K., Le Roy, S., Van Oyen, H., Griffiths, C., Michel, J. P., & Herrmann, F. R. (2008). Death toll exceeded 70,000 in Europe during the summer of 2003. *Comptes Rendus—Biologies*. <https://doi.org/10.1016/j.crv.2007.12.001>.
- Salimi, F., Henderson, S. B., Morgan, G. G., Jalaludin, B., & Johnston, F. H. (2017). Ambient particulate matter, landscape fire smoke, and emergency ambulance dispatches in Sydney, Australia. *Environment International*. <https://doi.org/10.1016/j.envint.2016.11.018>.
- Shaposhnikov, D., Revich, B., Bellander, T., Bedada, G. B., Bottai, M., Kharkova, T. ... Pershagen, G. (2014). Mortality related to air pollution with the Moscow heat wave and wildfire of 2010. *Epidemiology*. <https://doi.org/10.1097/EDE.0000000000000090>.
- Sherwood, S., & Huber, M. (2010). An adaptability limit to climate change due to heat stress. *Pnas*. <https://doi.org/10.1073/pnas.0913352107/-/DCSupplemental>. www.pnas.org/cgi/doi/10.1073/pnas.0913352107.
- Smith, K. R., Woodward, A., Campbell-Lendrum, D., Chadee, D., Honda, Y., Liu, Q. ... Genova, R. (2014). *Human health: Impacts, adaptation, and co-benefits*. Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change [Field CB, Barros VR, Dokken DJ, Mach KJ, Ma. <https://doi.org/10.1017/CBO9781107415379.016>.
- Steffen, W., Rice, M., & Alexander, D. (2018). *2017: Another record-breaking year for heat and extreme weather*. Sydney: Climate Council of Australia.
- Stephenson, C. (2010). *A literature review on the economic, social and environmental impacts of severe bushfires in south-eastern Australia*. (V. D. of S. and Environment & E. M. Australia, Eds.). Melbourne: Victorian Government Department of Sustainability and Environment.
- Stephenson, C., Handmer, J., & Betts, R. (2013). Estimating the economic, social and environmental impacts of wildfires in Australia. *Environmental Hazards*. <https://doi.org/10.1080/17477891.2012.703490>.
- Toloo, G. S., Yu, W., Aitken, P., FitzGerald, G., & Tong, S. (2014). The impact of heatwaves on emergency department visits in Brisbane, Australia: A time series study. *Critical Care (London, England)*. <https://doi.org/10.1186/cc13826>.
- Tong, S., FitzGerald, G., Wang, X. Y., Aitken, P., Tippet, V., Chen, D. ... Guo, Y. (2015). Exploration of the health risk-based definition for heatwave: A multi-city study. *Environmental Research*. <https://doi.org/10.1016/j.envres.2015.09.009>.

- Tong, S., Wang, X. Y., Yu, W., Chen, D., & Wang, X. (2014). The impact of heatwaves on mortality in Australia: A multicity study. *BMJ Open*. <https://doi.org/10.1136/bmjopen-2013-003579>.
- Turner, L. R., Connell, D., & Tong, S. (2013). The effect of heat waves on ambulance attendances in Brisbane, Australia. *Prehospital and Disaster Medicine*. <https://doi.org/10.1017/S1049023X13008789>.
- Walker, B., & Salt, D. (2006). Resilience thinking: Sustaining ecosystems and people in a changing world. Washington, DC: Island Press.
- Wang, X.-Y., Barnett, A., Guo, Y.-M., Yu, W.-W., Shen, X.-M., & Tong, S.-L. (2014). Increased risk of emergency hospital admissions for children with renal diseases during heatwaves in Brisbane, Australia. *World Journal of Pediatrics: WJP*. <https://doi.org/10.1007/s12519-014-0469-x>.
- Whittaker, J., Handmer, J., & Mercer, D. (2012). Vulnerability to bushfires in rural Australia: A case study from East Gippsland, Victoria. *Journal of Rural Studies*. <https://doi.org/10.1016/j.jrurstud.2011.11.002>.
- WHO. (2013). *Review of evidence on health aspects of air pollution—REVIHAAP Project: Technical Report*. Copenhagen, Denmark: WHO.
- WHO. (2014a). *Burden of disease from Ambient Air pollution for 2012*. Retrieved from http://www.who.int/phe/health_topics/outdoorair/databases/FINAL_HAP_AAP_BoD_24March2014.pdf?ua=1.
- WHO. (2014b). *Quantitative risk assessment of the effects of climate change on selected causes of death, 2030s and 2050s*. WHO. ISBN: 978-92-4-150769-1.
- Williams, S., Nitschke, M., Weinstein, P., Pisaniello, D. L., Parton, K. A., & Bi, P. (2012). The impact of summer temperatures and heatwaves on mortality and morbidity in Perth, Australia 1994–2008. *Environment International*. <https://doi.org/10.1016/j.envint.2011.11.011>.
- Wilson, L. A., Gerard Morgan, G., Hanigan, I. C., Johnston, F. H., Abu-Rayya, H., Broome, R. ... Jalaludin, B. (2013). The impact of heat on mortality and morbidity in the Greater Metropolitan Sydney Region: A case crossover analysis. *Environmental Health: A Global Access Science Source*. <https://doi.org/10.1186/1476-069X-12-98>.
- Wreford, A., & Neil Adger, W. (2010). Adaptation in agriculture: Historic effects of heat waves and droughts on UK agriculture. *International Journal of Agricultural Sustainability*. <https://doi.org/10.3763/ijas.2010.0482>.
- Xiao, J., Spicer, T., Jian, L., Yun, G. Y., Shao, C., Nairn, J. ... Weeramanthri, T. S. (2017). Variation in population vulnerability to heat wave in Western Australia. *Frontiers in Public Health*. <https://doi.org/10.3389/fpubh.2017.00064>.
- Youssef, H., Liousse, C., Roblou, L., Assamoi, E. M., Salonen, R. O., Maesano, C., ... Annesi-Maesano, I. (2014). Non-accidental health impacts of wildfire smoke. *International Journal of Environmental Research and Public Health*. <https://doi.org/10.3390/ijerph11111772>.
- Zuo, J., Pullen, S., Palmer, J., Bennetts, H., Chileshe, N., & Ma, T. (2015). Impacts of heat waves and corresponding measures: A review. *Journal of Cleaner Production*. <https://doi.org/10.1016/j.jclepro.2014.12.078>.